

CBSE | CUET | JEE

www.mtg.in
March 2023
Pages 100 | ₹ 75

MATHEMATICS TODAY

India's #1
MATHEMATICS MONTHLY FOR
CBSE, CUET, JEE & OTHER PET EXAMS

ARE YOU READY?
FOR JEE 2023?

See inside page no. 47-54

PRACTICE PAPERS
JEE MAIN
WB JEE | CUET

2023: SOLVED PAPER
JEE MAIN
SESSION 1

PRACTICE PAPER 2023
CBSE
warm up!



mtg

Trust of more than
1 Crore Readers
Since 1982



2023030013622



VISIT...

LANZAROTE
Caliente.COM



SRM

INSTITUTE OF SCIENCE & TECHNOLOGY
(Deemed to be University u/s 3 of UGC Act, 1956)


SRMJEEE B.Tech 2023

April | June | July


LEARNERS BECOMING LEADERS SRM - A BRAND TO RECKON WITH!

SRMIST is a uniquely large pioneering multi-disciplinary institution

GLOBAL RECOGNITION



University
19th Rank



Offers UG/PG/PhD Programs in Engineering and Technology
Management | Science and Humanities | Law
Medicine & Health Sciences | Agricultural Sciences

www.srmist.edu.in

Helpline : 080 690 87000

Email : admissions.india@srmist.edu.in



Disclaimer : *The content available in MTG digital/physical books is intellectual property of MTG Learning Media Pvt Ltd. Paid content on this website after purchase is not entitled for free distribution on whatsapp, telegram, twitter, facebook, and all other social media and digital channels. The registered owner/admin/ individual of any digital channel will be liable for an infringement of the copyright for illegally using the material / copyrighted content of MTG Learning Media Pvt Ltd.*

ADMISSION OPEN



Programs Offered

- MAJMC • MBA • M.Sc / MA in Financial Economics • MCA • M.Com
- LL.M. • BAJMC (Hons.) • BBA (Honours) • BBA LL.B. (Hons.)
- B.Com (Hons.) • B.Sc. (Hons.) Data Science • BCA

Distinct Advantages

- Scholarship for Meritorious Students
- Activity-Based Self-Learning
- Employment Enhancement Programs
- Short-Term Live Projects
- Modern Media & Digital Labs
- Regular interaction with industry experts
- Excellent Industry - Academia Interface
- Faculty with industry & Academic Experience
- Case and Project Based Teaching Pedagogy
- Industry-oriented Curriculum & Pedagogy

The University Provides Excellent Placement

BGU has collaboration with institutes/ Universities in France, Poland, Slovenia and Malaysia

Visit us at www.bgu.ac.in Email: admission@bgu.ac.in
Toll-Free No: 1800-212-3001, Mobile: 7381058307, 9776129900
IDCO Plot No.2, Institutional Area, Gothapatna, Bhubaneswar - 751029

How to choose the right answer, fast?

₹ 850

₹ 850

₹ 850

₹ 999

Visit www.mtg.in for latest offers and to buy online!

The answer is practice...

Our team has seen that in NEET and JEE, Multiple Choice Questions (MCQs) are based on the NCERT syllabus. Largely !! With Objective NCERT at your FINGERTIPS, you can become a pro at handling MCQs. Practice to increase your accuracy and improve timing with a bank of approx. 15,000 questions, all framed from NCERT course books. Don't take our word, have a look what some of our readers / toppers have to say...

Features:

- NCERT NEET/JEE Trend Indicator
- NCERT Notes with HD Pages
- NCERT Based Topicwise MCQs
- NCERT Exemplar Problems MCQs
- NCERT Exam Scorer : A & R, Case Based & NV Type Questions, HOTS
- NCERT Based Exam Archive Questions
- Practice Papers & Model Test Papers

Tanishka Yadav, "I have used MTG NCERT Fingertips and Biology Today magazine. They provided me additional information that proved to be very useful in NEET preparation. MCQs given in the MTG NCERT Fingertips gave me a fair idea of what to expect in the exam."

Nalin Khandelwal, "MTG NCERT at Your Fingertips-Biology and MTG 31 Years NEET- AIPMT Chapterwise Solutions helped me a lot in my preparation. My advice for future aspirants would be to be thorough with NCERT by doing multiple readings till all the concepts in all subjects are clear. Also, solving previous years' papers is very helpful."

Bhavik Bansal, "I have followed MTG NCERT at Your Fingertips, AIIMS Explorer Assertion and Reason books and magazines Physics For You, Biology Today and Chemistry Today. I used to solve questions from CBSE Drill and NEET Ready from MTG Magazines."

Kalpna Kumari, "I have followed MTG NCERT at Your Fingertips, Assertion and Reason books and magazines Physics For You, Biology Today and Chemistry Today. In my opinion, full dedication and smartly planning your preparation are the most important components of an ideal preparation."

Mrinal Dewangan, "MTG Objective NCERT at your Fingertips is a great book and Explorers provided very accurate question papers similar to those asked in exams, which helped me a lot during preparation."

Hei Shah, "For MCQs I referred to 'MTG Objective NCERT at your Fingertips' which is an exceptional book in itself. Also I referred to all the MTG Books for previous years' question papers of different exams."




Scan now with your smartphone or tablet

Application to read QR codes required


Study right. Dig deep.

mtg


Build a solid foundation for success
in JEE Main



₹ 1050/-



₹ 1050/-



₹ 1050/-

Are you a do-it-yourself type of a student? Then for success in JEE Main, choose MTG's JEE Main combo, comprising coursebooks for Physics, Chemistry & Mathematics. This combo is all class XI and XII students need for a solid and deep understanding of concepts in these three key subjects.

FEATURES

- Based on latest pattern of JEE Main
- Covers the entire syllabus
- Full of graphic illustrations for deep understanding of concepts
- Exhaustive Theory with Illustrations and Self Check MCQs
- Chapterwise Concept Booster, Rank Booster and JEE Connect MCQs
- Chapterwise Numerical Value Type Questions
- JEE Archive-Solved Previous Years' Questions of JEE Main
- 3 Mock Test Papers based on Latest Pattern of JEE Main



Scan now with your
smartphone or tablet
Application to read
QR codes required

Note: Coursebooks are also available separately.


Available at all leading book shops throughout India. To buy online visit www.mtg.in.

For more information or for help in placing your order, call 0124-6601200 or e-mail: info@mtg.in



FULLY LOADED & COMPLETELY UPDATED

MTG's BITSAT Power Guide is not only the most exhaustive prep-tool, but also the only book available at present, updated as per the latest BITSAT syllabus for students aspiring for top rank in BITSAT 2023.



₹ 999/-

Get MTG's BITSAT Power Guide today for a real-world feel of BITSAT. Find out what's different about the BITSAT test, including its pattern of examination and key success factors. Be it with chapterwise MCQs or model test papers, check how good your chances are for glory in BITSAT 2023.

FEATURES

- Covers all 5 subjects - Physics, Chemistry, Mathematics, English & Logical Reasoning
- Chapterwise MCQs in each section for practice
- Past chapterwise BITSAT Qs (2022-2018)
- 10 Model Test Papers with detailed solutions

Visit www.MTG.in to buy online. Or visit a leading bookseller near you.
For more information, email info@mtg.in or call **1800-10-38673** (toll-free).

MATHEMATICS today

VOLUME 41

No. 3

March 2023

Managing Editor
Mahabir Singh

Editor
Anil Ahlawat

Corporate Office:

Plot 99, Sector 44 Institutional area, Gurugram - 122 003 (HR).
Tel : 0124-6601200 e-mail : info@mtg.in website : www.mtg.in

Regd. Office:

406, Taj Apartment, Near Safdarjung Hospital, New Delhi - 110029.

Contents

Competition Edge

JEE Main Solved Paper 2023 **9**

(Session-1, 24th Jan, Morning Shift)

JEE Main Practice Paper 2023 **16**

CUET (UG) Chapterwise Practice Paper 2023 **23**

Practice Paper

WB JEE Practice Paper 2023 **37**

Are You Ready for JEE? **47**

GK Corner **55**

Unique Career in Demand **57**

JEE Work Outs **58**

Quantitative Aptitude **65**

Brain Teaser **70**

Beat the Time Trap **73**

You Ask We Answer **76**

Quiz Club **91**

Class 11

Monthly Test Drive **77**

Straight Lines

Class 12

CBSE Warm Up! **80**

Practice Paper 2022-23

Monthly Test Drive **92**

Differential Equations

Subscribe online at www.mtg.in

SUBSCRIPTION PLAN				
Subject	Format	1 Year	2 Years	3 Years
Individual • Physics • Chemistry • Mathematics • Biology	Print issue	₹749/-	₹1349/-	₹1799/-
	Print + Digital	₹1150/-	₹1850/-	₹2400/-
Combo of 3 Magazines • Physics, Chemistry, Biology • Physics, Chemistry, Mathematics	Print issue	₹2200/-	₹4000/-	₹5400/-
	Print + Digital	₹3000/-	₹5000/-	₹6600/-
Combo of 4 Magazines • Physics, Chemistry, • Mathematics, Biology	Print issue	₹2800/-	₹5200/-	₹6300/-
	Print + Digital	₹3800/-	₹6399/-	₹7800/-
Courier Charges Add to your subscription amount for quicker & traceable delivery		₹800/-	₹960/-	₹1440/-

Send D.D/M.O in favour of MTG Learning Media (P) Ltd.
Payments should be made directly to : MTG Learning Media (P) Ltd,
Plot No. 99, Sector 44, Gurugram - 122003 (Haryana)
We have not appointed any subscription agent.


Printed and Published by Mahabir Singh on behalf of MTG Learning Media Pvt. Ltd. Printed at HT Media Ltd., B-2, Sector-63, Noida, UP-201307 and published at 406, Taj Apartment, Ring Road, Near Safdarjung Hospital, New Delhi - 110029.
Editor : Anil Ahlawat

Readers are advised to make appropriate thorough enquiries before acting upon any advertisement published in this magazine. Focus/ Infocus features are marketing incentives. MTG does not vouch or subscribe to the claims and representations made by advertisers. All disputes are subject to Delhi jurisdiction only.


Copyright© MTG Learning Media (P) Ltd.

All rights reserved. Reproduction in any form is prohibited.


Last Minute Check on your JEE Readiness



₹ 675/-



₹ 675/-



₹ 675/-

The Wait is Over! MTG Presents you all the 80 Fully Solved online Question Papers of JEE Main of past 4 years (2022-2019) conducted by NTA with Chapterwise graphical analysis showing the weightage of chapters and Mock Test Papers as per latest pattern of JEE Main. Get the maximum benefit from the book through personalised course on **Pedagogy app**. The best preparation for tomorrow is doing your best today so, go and get your copy.

Highlights

- Fully Solved Authentic 80 Online Papers
- 6895 Questions for Practice
- Chapterwise Tabular and Graphical Analysis Showing the Weightage of Chapters
- 5 MTPs as per the latest JEE Main pattern

Practice Online Tests to Predict Your Rank

Make It Happen, CRACK JEE Main

Get the complete feel of Online exam
by attempting tests on computers



Take the Final Leap Towards **JEE Main** Success!

Available at all leading book shops throughout the country. To buy online visit www.mtg.in.
For more information or for help in placing your order,
Call 0124-6601200 or email: info@mtg.in



Scan to explore & buy more
JEE Books on mtg.in

JEE MAIN 2023

SECTION-A (MULTIPLE CHOICE QUESTIONS)

- Let $y = y(x)$ be the solution of the differential equation $x^3 dy + (xy - 1) dx = 0, x > 0, y\left(\frac{1}{2}\right) = 3 - e$. Then $y(1)$ is equal to
(a) $2 - e$ (b) e (c) 1 (d) 3
- The area enclosed by the curves $y^2 + 4x = 4$ and $y - 2x = 2$ is
(a) $\frac{23}{3}$ (b) $\frac{22}{3}$ (c) 9 (d) $\frac{25}{3}$
- Let Ω be the sample space and $A \subseteq \Omega$ be an event. Given below are two statements
(S1) : If $P(A) = 0$, then $A = \phi$
(S2) : If $P(A) = 1$, then $A = \Omega$
Then
(a) both (S1) and (S2) are true
(b) only (S1) is true
(c) only (S2) is true
(d) both (S1) and (S2) are false
- The value of $\sum_{r=0}^{22} {}^{22}C_r {}^{23}C_r$ is
(a) ${}^{45}C_{24}$ (b) ${}^{45}C_{23}$ (c) ${}^{44}C_{23}$ (d) ${}^{44}C_{22}$
- Let α be a root of the equation $(a - c)x^2 + (b - a)x + (c - b) = 0$ where a, b, c are distinct real numbers such that the matrix $\begin{bmatrix} \alpha^2 & \alpha & 1 \\ 1 & 1 & 1 \\ a & b & c \end{bmatrix}$ is singular. Then, the value of $\frac{(a - c)^2}{(b - a)(c - b)} + \frac{(b - a)^2}{(a - c)(c - b)} + \frac{(c - b)^2}{(a - c)(b - a)}$ is
(a) 6 (b) 3 (c) 9 (d) 12
- $\lim_{t \rightarrow 0} \left(\frac{1}{1 \sin^2 t} + \frac{1}{2 \sin^2 t} + \dots + n \frac{1}{n \sin^2 t} \right)^{\sin^2 t}$ is equal to
(a) n^2 (b) $n^2 + n$ (c) n (d) $\frac{n(n+1)}{2}$
- The distance of the point $(-1, 9, -16)$ from the plane $2x + 3y - z = 5$ measured parallel to the line $\frac{x+4}{3} = \frac{2-y}{4} = \frac{z-3}{12}$ is
(a) $20\sqrt{2}$ (b) 31 (c) $13\sqrt{2}$ (d) 26
- Let N denote the number that turns up when a fair die is rolled. If the probability that the system of equations
 $x + y + z = 1$
 $2x + Ny + 2z = 2$
 $3x + 3y + Nz = 3$
has unique solution is $k/6$, then the sum of value of k and all possible values of N is
(a) 18 (b) 19 (c) 20 (d) 21
- Let $f(x) = \begin{cases} x^2 \sin\left(\frac{1}{x}\right), & x \neq 0 \\ 0, & x = 0 \end{cases}$
Then at $x = 0$
(a) f is continuous but not differentiable
(b) f is continuous but f' is not continuous
(c) f' is continuous but not differentiable
(d) f and f' both are continuous
- Let PQR be a triangle. The points A, B and C are the sides QR, RP and PQ respectively such that $\frac{QA}{AR} = \frac{RB}{BP} = \frac{PC}{CQ} = \frac{1}{2}$. Then $\frac{\text{Area}(\Delta PQR)}{\text{Area}(\Delta ABC)}$ is equal to
(a) 4 (b) 3 (c) 2 (d) $5/2$

11. For three positive integers p, q, r , $x^{pq^2} = y^{qr} = z^{p^2r}$ and $r = pq + 1$ such that $3, 3 \log_y x, 3 \log_z y, 7 \log_x z$ are in A.P. with common difference $1/2$. The $r - p - q$ is equal to
(a) 6 (b) 2 (c) 12 (d) -6
12. $\tan^{-1}\left(\frac{1+\sqrt{3}}{3+\sqrt{3}}\right) + \sec^{-1}\left(\sqrt{\frac{8+4\sqrt{3}}{6+3\sqrt{3}}}\right)$ is equal to
(a) $\frac{\pi}{2}$ (b) $\frac{\pi}{4}$ (c) $\frac{\pi}{6}$ (d) $\frac{\pi}{3}$
13. The equation $x^2 - 4x + [x] + 3 = x[x]$, where $[x]$ denotes the greatest integer function, has
(a) no solution
(b) exactly two solutions in $(-\infty, \infty)$
(c) a unique solution in $(-\infty, 1)$
(d) a unique solution in $(-\infty, \infty)$
14. Let $p, q \in \mathbb{R}$ and $(1 - \sqrt{3}i)^{200} = 2^{199}(p + iq)$, $i = \sqrt{-1}$. The $p + q + q^2$ and $p - q + q^2$ are roots of the equation
(a) $x^2 - 4x + 1 = 0$ (b) $x^2 + 4x + 1 = 0$
(c) $x^2 - 4x - 1 = 0$ (d) $x^2 + 4x - 1 = 0$
15. If A and B are two non-zero $n \times n$ matrices such that $A^2 + B = A^2B$, then
(a) $A^2 = I$ or $B = I$ (b) $A^2B = BA^2$
(c) $AB = I$ (d) $A^2B = I$
16. Let a tangent to the curve $y^2 = 24x$ meet the curve $xy = 2$ at the points A and B . Then the mid points of such line segments AB lie on a parabola with the
(a) length of latus rectum $3/2$
(b) length of latus rectum 2
(c) directrix $4x = 3$
(d) directrix $4x = -3$
17. The distance of the point $(7, -3, -4)$ from the plane passing through the points $(2, -3, 1)$, $(-1, 1, -2)$ and $(3, -4, 2)$ is
(a) $5\sqrt{2}$ (b) $4\sqrt{2}$ (c) 4 (d) 5
18. The compound statement $(\sim(P \wedge Q)) \vee ((\sim P) \wedge Q) \Rightarrow ((\sim P) \wedge (\sim Q))$ is equivalent to
(a) $((\sim P) \vee Q) \wedge ((\sim Q) \vee P)$
(b) $(\sim Q) \vee P$
(c) $((\sim P) \vee Q) \wedge (\sim Q)$ (d) $(\sim P) \vee Q$
19. The relation $R = \{(a, b) : \gcd(a, b) = 1, 2a \neq b, a, b \in \mathbb{Z}\}$ is
(a) reflexive but not symmetric
(b) transitive but not reflexive
(c) symmetric but not transitive
(d) neither symmetric nor transitive

20. Let $\vec{u} = \hat{i} - \hat{j} - 2\hat{k}$, $\vec{v} = 2\hat{i} + \hat{j} - \hat{k}$, $\vec{v} \cdot \vec{w} = 2$ and $\vec{v} \times \vec{w} = \vec{u} + \lambda \vec{v}$. Then $\vec{u} \cdot \vec{w}$ is equal to
(a) 1 (b) 2 (c) $3/2$ (d) $-2/3$

SECTION - B (NUMERICAL VALUE TYPE)

Attempt any 5 questions out of 10.

21. Let $\lambda \in \mathbb{R}$ and let the equation $Ebe|x^2| - 2|x| + |\lambda - 3| = 0$. Then the largest element in the set $S = \{x + \lambda : x \text{ is an integer solution of } E\}$ is _____.
22. A boy needs to select five courses from 12 available courses, out of which 5 courses are language courses. If he can choose at most two language courses, then the number of ways he can choose five courses is _____.
23. The value of $\frac{8}{\pi} \int_0^{\pi/2} \frac{(\cos x)^{2023}}{(\sin x)^{2023} + (\cos x)^{2023}} dx$ is _____.
24. The shortest distance between the lines $\frac{x-2}{3} = \frac{y+1}{2} = \frac{z-6}{2}$ and $\frac{x-6}{3} = \frac{1-y}{2} = \frac{z+8}{0}$ is equal to _____.
25. The 4th term of GP is 500 and its common ratio is $1/m$, $m \in \mathbb{N}$. Let S_n denote the sum of the first n terms of this GP. If $S_6 > S_5 + 1$ and $S_7 < S_6 + \frac{1}{2}$, then the number of possible values of m is _____.
26. Let C be the largest circle centred at $(2, 0)$ and inscribed in the ellipse $\frac{x^2}{36} + \frac{y^2}{16} = 1$. If $(1, \alpha)$ lies on C , then $10\alpha^2$ is equal to _____.
27. Let a tangent to the curve $9x^2 + 16y^2 = 144$ intersect the coordinate axes at the points A and B . Then, the minimum length of the line segment AB is _____.
28. Suppose $\sum_{r=0}^{2023} r^2 \cdot {}^{2023}C_r = 2023 \times \alpha \times 2^{2022}$. Then the value of α is _____.
29. The value of $12 \int_0^3 |x^2 - 3x + 2| dx$ is _____.
30. The number of 9 digit numbers, that can be formed using all the digits of the number 123412341 so that the even digits occupy only even places, is _____.

SOLUTIONS

1. (c): Given, $x^3 dy + (xy - 1)dx = 0$

$$\Rightarrow x^3 dy = (1 - xy)dx$$

$$\Rightarrow \frac{dy}{dx} = \frac{1-xy}{x^3} \Rightarrow \frac{dy}{dx} + \frac{y}{x^2} = \frac{1}{x^3}, \text{ which is linear, D.E.}$$

$$\text{Now, I.F.} = e^{\int \frac{1}{x^2} dx} = e^{-\frac{1}{x}} = \frac{1}{e^{1/x}}$$

$$\text{Solution is given by } ye^{-1/x} = \int \frac{1}{x^3} e^{-1/x} dx + c$$

$$\text{Putting } u = \frac{-1}{x} \Rightarrow du = \frac{1}{x^2} dx, \text{ we get}$$

$$ye^{-1/x} = \int -e^u u du + c = -[ue^u - \int e^u du] + c \\ = -[ue^u - e^u] + c = (1 - u)e^u + c$$

$$\Rightarrow ye^{-1/x} = \left(1 + \frac{1}{x}\right)e^{-1/x} + c \quad \dots(i)$$

$$\text{Put } x = \frac{1}{2} \text{ and } y = 3 - e; 3e^{-2} - e^{-1} = 3e^{-2} + c \Rightarrow c = -e^{-1}$$

$$\text{Equation (i) becomes, } ye^{-1/x} = e^{-1/x} + \frac{e^{-1/x}}{x} - e^{-1}$$

$$\text{Put } x = 1 \text{ to get } y(1),$$

$$\Rightarrow ye^{-1} = e^{-1} + e^{-1} - e^{-1} = e^{-1} \Rightarrow y = 1$$

2. (c): Given curves are


$$y^2 + 4x = 4 \text{ and } y - 2x = 2$$

The points of intersection of the given curves are

$$(0, 2) \text{ and } (-3, -4).$$

Required area

$$= \int_{-4}^2 \left[\left(\frac{4-y^2}{4} \right) - \left(\frac{y-2}{2} \right) \right] dy \\ = \int_{-4}^2 \frac{-y^2 - 2y + 8}{4} dy = \frac{1}{4} \left[-\frac{y^3}{3} - y^2 + 8y \right]_{-4}^2 \\ = \frac{1}{4} \left[\left(-\frac{8}{3} - 4 + 16 \right) - \left(\frac{64}{3} - 16 - 32 \right) \right] \\ = \frac{1}{4} \left[\left(\frac{28}{3} \right) - \left(-\frac{80}{3} \right) \right] = \frac{1}{4} \left(\frac{108}{3} \right) = \frac{108}{12} = 9 \text{ sq. units}$$



3. (a): Ω be the Sample space A be an event.

$$\text{If } P(A) = 0 \Rightarrow A = \phi$$

$$\text{If } P(A) = 1 \Rightarrow A = \Omega$$

Then both statements are true.

$$\mathbf{4} \quad \mathbf{(b)} : \sum_{r=0}^{22} {}^{22}C_r {}^{23}C_r = \sum_{r=0}^{22} {}^{22}C_r {}^{23}C_{23-r} \\ = {}^{22+23}C_{23} = {}^{45}C_{23}$$

5. (b) : Given matrix is singular,

$$\therefore \begin{vmatrix} \alpha^2 & \alpha & 1 \\ 1 & 1 & 1 \\ a & b & c \end{vmatrix} = 0$$

$$\Rightarrow \alpha^2(c-b) - \alpha(c-a) + 1(b-a) = 0$$

$$\text{It is singular if } \alpha = 1 \Rightarrow c-b-c+a+b-a=0$$

$$\text{Now, } \frac{(a-c)^2}{(b-a)(c-b)} + \frac{(b-a)^2}{(a-c)(c-b)} + \frac{(c-b)^2}{(a-c)(b-a)} \\ = \frac{(a-c)^3}{(a-c)(b-a)(c-b)} + \frac{(b-a)^3}{(a-c)(b-a)(c-b)} \\ + \frac{(c-b)^3}{(a-c)(b-a)(c-b)}$$

$$= \frac{(a-c)^3 + (b-a)^3 + (c-b)^3}{(a-c)(b-a)(c-b)}$$

$$= \frac{3(a-c)(b-a)(c-b)}{(a-c)(b-a)(c-b)} = 3 \quad (\because a-c+b-a+c-b=0)$$

$$\mathbf{6. (c):} \text{ Let } L = \lim_{t \rightarrow 0} \left(\frac{1}{1^{\sin^2 t}} + \frac{1}{2^{\sin^2 t}} + \dots + \frac{1}{n^{\sin^2 t}} \right)^{\sin^2 t} \\ \Rightarrow L = \lim_{t \rightarrow 0} n \left[\left(\frac{1}{n} \right)^{\csc^2 t} + \left(\frac{2}{n} \right)^{\csc^2 t} + \dots + \left(\frac{n-1}{n} \right)^{\csc^2 t} + 1 \right]^{\sin^2 t}$$

$$\Rightarrow L = n(1^{\sin^2(0)}) = n(1) = n$$

$$\left[0 < \frac{k}{n} < 1, \left(\frac{k}{n} \right)^\infty = 0 \text{ for } 1 \leq k < n \right]$$

7. (d) : Equation of line passing through the point

$$(-1, 9, -16) \text{ and parallel to line } \frac{x+4}{3} = \frac{2-y}{4} = \frac{z-3}{12} \text{ is } \\ \frac{x+1}{3} = \frac{y-9}{-4} = \frac{z+16}{12} = \lambda$$

$$\text{Any point on this line} = (3\lambda - 1, -4\lambda + 9, 12\lambda - 16)$$

Point of intersection line and plane

$$2(3\lambda - 1) + 3(-4\lambda + 9) - 1(12\lambda - 16) = 5$$

$$\Rightarrow 6\lambda - 2 - 12\lambda + 27 - 12\lambda + 16 = 5$$

$$\Rightarrow \lambda = 2 \therefore \text{Point} = (5, 1, 8)$$

$$\therefore \text{Distance} = \sqrt{(5+1)^2 + (9-1)^2 + (-16-8)^2}$$

$$= \sqrt{36 + 64 + 576} = 26 \text{ units}$$

8. (c): For unique solution, determinant $\neq 0$

$$\begin{vmatrix} 1 & 1 & 1 \\ 2 & N & 2 \\ 3 & 3 & N \end{vmatrix} \neq 0$$

$\Rightarrow N^2 - 5N + 6 \neq 0 \Rightarrow (N-2)(N-3) \neq 0 \Rightarrow N \neq 2, 3$
 $\therefore$ Possible values of N are 1, 4, 5, 6 and $\frac{k}{6} = \frac{4}{6} \Rightarrow k = 4$
 Required sum = $1 + 4 + 5 + 6 + 4 = 20$

9. (b) : $f(x) = \begin{cases} x^2 \sin\left(\frac{1}{x}\right), & x \neq 0 \\ 0, & x = 0 \end{cases}$

LHD = $\lim_{h \rightarrow 0} \frac{f(0-h) - f(0)}{-h} = \lim_{h \rightarrow 0^-} -\frac{h^2 \sin(1/h)}{-h} = 0$

RHD = $\lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} = \lim_{h \rightarrow 0^+} \frac{h^2 \sin(1/h)}{h} = 0$

$\Rightarrow f(x)$ is continuous and differentiable at $x = 0$

Now, $f'(x) = \begin{cases} 2x \sin(1/x) - \cos(1/x), & x \neq 0 \\ 0, & x = 0 \end{cases}$

$\lim_{x \rightarrow 0} f'(x) = \lim_{x \rightarrow 0} \left[2x \sin\left(\frac{1}{x}\right) - \cos\left(\frac{1}{x}\right) \right] = 0 - [-1, 1] \neq 0$

$f'(0) = 0$

Hence, $f'(x)$ is discontinuous at $x = 0$.

10. (b) : Let the position vector of P, Q, R be $\vec{0}, \vec{a}, \vec{b}$

$\Rightarrow$ P.V. of $A = \frac{2\vec{a} + \vec{b}}{3}$, P.V. of $B = \frac{2\vec{b}}{3}$

and P.V. of $C = \frac{\vec{a}}{3}$

$\therefore \vec{AB} = \frac{2\vec{b}}{3} - \left(\frac{2\vec{a} + \vec{b}}{3} \right) = \frac{\vec{b}}{3} - \frac{2\vec{a}}{3} = \frac{\vec{b} - 2\vec{a}}{3}$


$\vec{CA} = \frac{2\vec{a} + \vec{b}}{3} - \frac{\vec{a}}{3} = \frac{\vec{a} + \vec{b}}{3}$

Area of $\Delta PQR = \frac{1}{2} |\vec{PQ} \times \vec{PR}| = \frac{1}{2} |\vec{a} \times \vec{b}|$

Area of $\Delta ABC = \frac{1}{2} |\vec{CA} \times \vec{AB}|$

$= \frac{1}{2} \left| \left(\frac{\vec{a} + \vec{b}}{3} \right) \times \left(\frac{\vec{b} - 2\vec{a}}{3} \right) \right| = \frac{1}{2} \left| \frac{\vec{a} \times \vec{b}}{3} \right|$

$\therefore \frac{\text{Area}(\Delta PQR)}{\text{Area}(\Delta ABC)} = \frac{\frac{1}{2} |\vec{a} \times \vec{b}|}{\frac{1}{2} \left| \frac{\vec{a} \times \vec{b}}{3} \right|} = 3$



11. (b) : $pq^2 \log x = qr \log y = p^2 r \log z$

Now, $\frac{3 \log x}{\log y} - 3 = \frac{1}{2} \Rightarrow \frac{\log x}{\log y} = \frac{7}{6} = \frac{qr}{pq^2} = \frac{r}{pq}$

$\Rightarrow 7pq = 6r \Rightarrow 7(r-1) = 6r \Rightarrow r = 7$

and $\frac{3 \log y}{\log z} - 3 = 1 \Rightarrow \frac{\log y}{\log z} = \frac{4}{3} = \frac{p^2 r}{qr} = \frac{p^2}{q} \Rightarrow 3p^2 = 4q$

Also, $\frac{7 \log z}{\log x} - 3 = \frac{3}{2} \Rightarrow \frac{7 \log z}{\log x} = \frac{9}{2} = \frac{7pq^2}{p^2 r} = \frac{7q^2}{pr}$


$\Rightarrow \frac{7q^2}{p \times 7} = \frac{9}{2} \Rightarrow \frac{q^2}{p} = \frac{9}{2} \Rightarrow 2q^2 = 9p \Rightarrow 4q^4 = 81p^2$

$\Rightarrow 4q^4 = 27 \times 3p^2 = 27 \times 4q \Rightarrow q^3 = 27 \Rightarrow q = 3$

We have, $r = pq + 1 \Rightarrow 7 = 3p + 1 \Rightarrow p = 2$

$\therefore r - p - q = 7 - 2 - 3 = 2$

12. (d) : $\tan^{-1} \left(\frac{1+\sqrt{3}}{3+\sqrt{3}} \right) + \sec^{-1} \left(\sqrt{\frac{8+4\sqrt{3}}{6+3\sqrt{3}}} \right)$



$= \tan^{-1} \left(\frac{1+\sqrt{3}}{\sqrt{3}(\sqrt{3}+1)} \right) + \tan^{-1} \left(\frac{\sqrt{2+\sqrt{3}}}{\sqrt{6+3\sqrt{3}}} \right)$

$= \tan^{-1} \left(\frac{1}{\sqrt{3}} \right) + \tan^{-1} \left(\frac{\sqrt{2+\sqrt{3}}}{\sqrt{3}(2+\sqrt{3})} \right)$

$= \tan^{-1} \left(\frac{1}{\sqrt{3}} \right) + \tan^{-1} \left(\frac{1}{\sqrt{3}} \right) = \frac{\pi}{6} + \frac{\pi}{6} = \frac{\pi}{3}$

13. (d) : $x^2 - 4x + [x] + 3 = x[x]$

$\Rightarrow x^2 - x[x] - (x - [x]) + 3(x - 1) = 0$

$\Rightarrow x(x - [x]) - 1(x - [x]) - 3(x - 1) = 0$

$\Rightarrow (x - 1)(x - [x]) - 3(x - 1) = 0$

$\Rightarrow (x - 1)(x - [x] - 3) = 0$

$\Rightarrow (x - 1)(\{x\} - 3) = 0 \Rightarrow x = 1$ as $\{x\} \neq 3$

14. (a): Given, $(1 - \sqrt{3}i)^{200} = 2^{199}(p + iq)$

$\Rightarrow (2e^{-i\pi/3})^{200} = 2^{199}(p + iq)$

$\Rightarrow 2^{200} \left(\cos \frac{\pi}{3} - i \sin \frac{\pi}{3} \right) = 2^{199}(p + iq)$

$\Rightarrow 2 \left(\cos \frac{200\pi}{3} - i \sin \frac{200\pi}{3} \right) = p + iq$

$\therefore p = -1, q = -\sqrt{3}$

$\Rightarrow p + q + q^2 = -1 - \sqrt{3} + 3 = 2 - \sqrt{3}$

and $p - q + q^2 = -1 + \sqrt{3} + 3 = 2 + \sqrt{3}$

Sum of root = $2 + \sqrt{3} + 2 - \sqrt{3} = 4$

Product of root = $(2 + \sqrt{3})(2 - \sqrt{3}) = 4 - 3 = 1$

Required quadratic equation is $x^2 - 4x + 1 = 0$

15. (b) : $A^2 + B = A^2B$

$$\begin{aligned} \Rightarrow A^2 + B - A^2B &= 0 \Rightarrow A^2 - A^2B + B = 0 \\ \Rightarrow A^2(I - B) + B &= 0 \Rightarrow A^2(I - B) + B - I = -I \\ \Rightarrow (A^2 - I)(I - B) &= -I \Rightarrow (A^2 - I)(B - I) = I \\ \Rightarrow A^2B - A^2 - B &= 0 \Rightarrow A^2(B - I) = B \\ \Rightarrow A^2 &= B(B - I)^{-1} \Rightarrow A^2 = B(A^2 - I) \\ \Rightarrow A^2 &= BA^2 - B \Rightarrow A^2 + B = BA^2 \Rightarrow A^2B = BA^2 \end{aligned}$$

16. (c) : Let the equation of tangent to

$$y^2 = 24x \text{ is } ty = x + 6t^2$$

Now, $ty = x + 6t^2$ meet the curve $xy = 2$ at point A and B.

Let mid-point of AB be (h, k)

$$ty = \frac{2}{y} + 6t^2 \quad \left[\because x = \frac{2}{y} \right]$$

$$ty^2 - 6t^2y - 2 = 0; y_1 + y_2 = 6t$$

$$t \cdot \frac{2}{x} = x + 6t^2 \quad \left[\because y = \frac{2}{x} \right]$$

$$x^2 + 6t^2x - 2t = 0; x_1 + x_2 = -6t^2$$

Mid point P is $(-3t^2, 3t)$

$$\Rightarrow h = -3t^2, k = 3t \Rightarrow \left(\frac{h}{-3} \right) = \left(\frac{k}{3} \right)^2 \Rightarrow y^2 = -3x$$

17. (a) : Let A(2, -3, 1), B(-1, 1, -2) and C(3, -4, 2)
Cartesian equation of plane passing through A, B and C is

$$\begin{vmatrix} x-2 & y+3 & z-1 \\ -3 & 4 & -3 \\ 1 & -1 & 1 \end{vmatrix} = 0$$

$$\Rightarrow (x-2)(4-3) - (y+3)(-3+3) + (z-1)(3-4) = 0$$

$$\Rightarrow x-2-z+1=0 \Rightarrow x-z=1$$

$$\text{Required distance} = \frac{|1(7) - (-4) - 1|}{\sqrt{1+1}} = \frac{10}{\sqrt{2}} = 5\sqrt{2} \text{ sq. units}$$

18. (a) : Constructing the truth table.

P	Q	$\sim P$	$\sim Q$	$P \wedge Q$	$\sim(P \wedge Q)$	$\sim P \wedge Q$	$\sim(P \wedge Q)$	$(\sim P)$	$(\sim(P \wedge Q))$
T	T	F	F	T	F	F	F	F	T
T	F	F	T	F	T	F	T	F	F
F	T	T	F	F	T	T	T	F	F
F	F	T	T	F	T	F	T	T	T

P	Q	$(\sim P) \vee Q$	$(\sim Q) \vee P$	$((\sim P) \vee Q) \wedge ((\sim Q) \vee P)$
T	T	T	T	T
T	F	F	T	F
F	T	T	F	F
F	F	T	T	T

Thus, the statement given in question is equivalent to $((\sim P \vee Q) \wedge ((\sim Q) \vee P))$

19. (d) : Reflexive : $\gcd(a, a) = a \neq 1$ for $a \neq 1$.

Symmetric : Let $a = 2, b = 1; \gcd(a, b) = 1, b \neq 2a$

Thus, $(2, 1) \in R; \gcd(1, 2) = 1$. But $2 = 2 \times 1$.

Therefore $(1, 2) \notin R$. Hence, R is not symmetric.

Transitive : $(2, 3) \in R; (3, 8) \in R$

But $(2, 8) \notin R$. Hence R is not transitive

R is neither symmetric nor transitive

20. (a) : $\vec{v} \times \vec{w} = (\vec{u} + \lambda \vec{v}) = \hat{i} - \hat{j} - 2\hat{k} + \lambda(2\hat{i} + \hat{j} - \hat{k})$

$$= (2\lambda + 1)\hat{i} + (\lambda - 1)\hat{j} - (2 + \lambda)\hat{k}$$

$$\text{Now, } \vec{v} \cdot (\vec{v} \times \vec{w}) = 0 \Rightarrow (2\hat{i} + \hat{j} - \hat{k}) \cdot ((2\lambda + 1)\hat{i} + (\lambda - 1)\hat{j} - (2 + \lambda)\hat{k}) = 0$$

$$\Rightarrow 2(2\lambda + 1) + \lambda - 1 + \lambda + 2 = 0 \Rightarrow 6\lambda + 3 = 0 \Rightarrow \lambda = -\frac{1}{2}$$

$$\vec{w} \cdot (\vec{v} \times \vec{w}) = \vec{w} \cdot (\vec{u} + \lambda \vec{v}) = \vec{u} \cdot \vec{w} + \lambda \vec{v} \cdot \vec{w} = 0$$

$$\vec{u} \cdot \vec{w} = -\lambda \vec{v} \cdot \vec{w}; \vec{u} \cdot \vec{w} = \frac{1}{2} \times 2 = 1$$

21. (5) : $|x|^2 - 2|x| + |\lambda - 3| = 0$

$$(|x| - 1)^2 + |\lambda - 3| - 1 = 0$$

...(i)

Given that x is an integer $\Rightarrow |x| - 1$ is an integer

$(|x| - 1)^2 - 1$ is also an integer. Thus $|\lambda - 3|$ is also an integer.

Therefore λ is also an integer, $(|x| - 1)^2 \leq 1$

Case I : $|x| - 1 = 0 \Rightarrow x = \pm 1 \therefore |\lambda - 3| = 1 \Rightarrow \lambda = 4, 2$

Possible values of $\lambda + x = 5, 3, 1$.

Case II : $|x| - 1 = 1 \Rightarrow x = \pm 2 \therefore |\lambda - 3| = 0 \Rightarrow \lambda = 3$

Possible values of $x + \lambda = 5, 1$

Considering all possible cases maximum value of

$$\lambda + x = 5$$

22. (546) : 5 language courses 7 non language courses

He can select :

(i) 0 language courses and 5 non language courses in

$${}^5C_0 \times {}^7C_5 = 21 \text{ ways}$$

(ii) 1 language courses and 4 non language courses in

$${}^5C_1 \times {}^7C_4 = 175 \text{ ways}$$

(iii) 2 language courses and 3 non language courses in

$${}^5C_2 \times {}^7C_3 = 350 \text{ ways}$$

Required number of ways = $21 + 175 + 350 = 546$ ways.

23. (2) : Let $I = \int_0^{\pi/2} \frac{(\cos x)^{2023}}{(\cos x)^{2023} + (\sin x)^{2023}} dx$... (i)

$$I = \int_0^{\pi/2} \frac{(\sin x)^{2023}}{(\sin x)^{2023} + (\cos x)^{2023}} dx \quad \dots (ii)$$

$$\left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

Adding equation (i) and (ii), we get

$$2I = \int_0^{\pi/2} 1 dx = \frac{\pi}{2} \Rightarrow I = \frac{\pi}{4} \Rightarrow \frac{8}{\pi} I = 2$$

24. (14): $L_1: \frac{x-2}{3} = \frac{y+1}{2} = \frac{z-6}{2}$; $L_2: \frac{x-6}{3} = \frac{y-1}{-2} = \frac{z+8}{0}$

Shortest distance = $\frac{\begin{vmatrix} 6-2 & 1-(-1) & -8-6 \\ 3 & 2 & 2 \\ 3 & -2 & 0 \end{vmatrix}}{\begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 3 & 2 & 2 \\ 3 & -2 & 0 \end{vmatrix}} = \frac{\begin{vmatrix} 4 & 2 & -14 \\ 3 & 2 & 2 \\ 3 & -2 & 0 \end{vmatrix}}{|\hat{4i} + \hat{6j} - 12\hat{k}|} = \frac{4(0+4) - 2(0-6) - 14(-6-6)}{\sqrt{4^2 + 6^2 + 12^2}} = \frac{16+12+14 \times 12}{14} = 14 \text{ units.}$

25. (12): Let the first term be A

$T_4 = 500$; $\frac{A}{m^3} = 500$; $S_6 - S_5 > 1 \Rightarrow T_6 > 1 \Rightarrow \frac{A}{m^5} > 1$
 $\Rightarrow \frac{500}{m^2} > 1 \Rightarrow m^2 < 500 \Rightarrow m < \sqrt{500} \Rightarrow m \leq 22$
 $(\because m \in \mathbb{N})$

$S_7 - S_6 < \frac{1}{2} \Rightarrow T_7 < \frac{1}{2} \Rightarrow \frac{A}{m^6} < \frac{1}{2} \Rightarrow \frac{500}{m^3} < \frac{1}{2}$
 $m^3 > 1000 \Rightarrow m \geq 11$

Thus, $11 \leq m \leq 22 \therefore m$ can take 12 values

26. (118): Let P be the point where ellipse and circle touch each other.

Let P be $(6 \cos \theta, 4 \sin \theta)$

Equation of tangent to ellipse at point P is

$$\frac{x \cos \theta}{6} + \frac{y \sin \theta}{4} = 1$$

$$\Rightarrow y = \frac{4}{\sin \theta} \left(\frac{-x \cos \theta}{6} + 1 \right) \Rightarrow y = -\frac{2 \cos \theta}{3 \sin \theta} x + \frac{4}{\sin \theta}$$

Let $m_1 = \frac{-2 \cos \theta}{3 \sin \theta}$.

Slope of normal to circle at P = m_2 .

$$m_2 = \frac{(4 \sin \theta - 0)}{(6 \cos \theta - 2)}; m_1 m_2 = -1$$

$$\Rightarrow \frac{-2 \cos \theta}{3 \sin \theta} \times \frac{4 \sin \theta}{(6 \cos \theta - 2)} = -1 \Rightarrow 8 \cos \theta = 3(6 \cos \theta - 2)$$

$$\Rightarrow 6 = 10 \cos \theta \Rightarrow \cos \theta = \frac{3}{5} \Rightarrow \sin \theta = \frac{4}{5} \text{ So, } P \equiv \left(\frac{18}{5}, \frac{16}{5} \right)$$

$$r^2 = \left(\frac{18}{5} - 2 \right)^2 + \left(\frac{16}{5} \right)^2 = (1-2)^2 + \alpha^2 = \frac{64}{25} + \frac{256}{25} = 1 + \alpha^2$$

$$\alpha^2 = \frac{320}{25} - 1 = \frac{64}{5} - 1 = \frac{59}{5} \therefore 10\alpha^2 = 59 \times 2 = 118$$

27. (7): $\frac{x^2}{16} + \frac{y^2}{9} = 1$. Let $(4 \cos \theta, 3 \sin \theta)$ be the coordinates of the point at which tangent is drawn.

Equation of tangent $\frac{x \cos \theta}{4} + \frac{y \sin \theta}{3} = 1$

Let A = $(4 \sec \theta, 0)$. B = $(0, 3 \operatorname{cosec} \theta)$

(Tangent intersects coordinate axes at A and B)

$$AB^2 = 16 \sec^2 \theta + 9 \operatorname{cosec}^2 \theta = 25 + 16 \tan^2 \theta + 9 \cot^2 \theta$$

$$AB^2 \geq 25 + 2 \times \sqrt{16 \tan^2 \theta \times 9 \cot^2 \theta} = 49; AB \geq 7$$

$$\therefore AB_{\min} = 7$$

28. (1012): $\sum_{r=0}^{2023} r^2 \cdot {}^{2023}C_r = 2023 \times \alpha \times 2^{2022}$

$$\sum_{r=0}^n r^2 \cdot {}^nC_r = n(n+1) \cdot 2^{n-2}$$

Then, $\sum_{r=0}^{2023} r^2 \cdot {}^{2023}C_r = (2023)(2023+1) \cdot 2^{2023-2}$

$$= 2023 \times 2024 \times 2^{2021} = 2023 \times \alpha \times 2^{2022}$$

$$\therefore 2\alpha = 2024; \alpha = 1012$$

29. (22): Let $I = 12 \int_0^3 |x^2 - 3x + 2| dx = 12 \int_0^3 |(x-2)(x-1)| dx$

$$= 12 \times \left[\int_0^1 (x^2 - 3x + 2) dx - \int_1^2 (x^2 - 3x + 2) dx + \int_2^3 (x^2 - 3x + 2) dx \right]$$

$$= 12 \times \left[\left(\frac{x^3}{3} - \frac{3x^2}{2} + 2x \right)_0^1 + \left(\frac{x^3}{3} - \frac{3x^2}{2} + 2x \right)_2^3 - \left(\frac{x^3}{3} - \frac{3x^2}{2} + 2x \right)_1^2 \right] = 22$$

30. (60): Even numbers are 2, 4, 2, 4

odd numbers are 1, 3, 1, 3, 1. $\bar{O} \bar{E} \bar{O} \bar{E} \bar{O} \bar{E} \bar{O} \bar{E} \bar{O}$

Even places can be filled with even numbers in $\frac{4!}{2! 2!}$ ways

Similarly, odd places can be filled in $\frac{5!}{2! 3!}$ ways

$$\text{Required number of ways} = \frac{4!}{2! 2!} \times \frac{5!}{2! 3!} = 60$$



MONTHLY TEST DRIVE CLASS XI ANSWER KEY


1. (b) 2. (d) 3. (a) 4. (a) 5. (d)
 6. (d) 7. (a,c) 8. (a,d) 9. (a,b,d) 10. (a,c)
 11. (a,c) 12. (b,d) 13. (a,d) 14. (b) 15. (c)
 16. (c) 17. (0.16) 18. (5) 19. (8) 20. (2188)

mtg

10 Very Similar Practice Tests **JEE Main**

Visit
www.mtg.in
for latest offers
and to buy
online!

₹ 250/-



HIGHLIGHTS

- 10 Very Similar Practice Tests as per the latest pattern of NTA JEE Main (60 MCQs+ 30 Numerical Value Type Questions)
- OMR sheet provided at the end of each test
- Detailed solutions of each practice test included



Scan to
explore & buy more
JEE Books on mtg.in

Available at all leading book shops throughout India. To buy online visit www.mtg.in.
For more information or for help in placing your order,
call 0124-6601200, 1800-10-38673 (Toll Free) or e-mail info@mtg.in

Also available on
*Flipkart, Snapdeal, Amazon,
Infibeam on cash on delivery*

PRACTICE PAPER 2023 JEE MAIN

Exam Dates

Session-2

06, 08, 10, 11, 12
April 2023

SECTION-A (MULTIPLE CHOICE QUESTIONS)

- The number of different words formed by using the letters 4A's, 2C's, 1H, 1D, 1B, if both C's, do not occur together, are
(a) 7560 (b) 1680 (c) 60 (d) 5880
- If $y = \left[x + \sqrt{x^2 - 1} \right]^{15} + \left[x - \sqrt{x^2 - 1} \right]^{15}$,
then $(x^2 - 1) \frac{d^2 y}{dx^2} + x \frac{dy}{dx}$ is equal to
(a) $225y^2$ (b) $224y^2$ (c) $125y$ (d) $225y$
- The set of all values of λ for which the system of linear equations
 $2x_1 - 2x_2 + x_3 = \lambda x_1$;
 $2x_1 - 3x_2 + 2x_3 = \lambda x_2$; $-x_1 + 2x_2 = \lambda x_3$
has a non-trivial solution,
(a) contains two elements.
(b) contains more than two elements.
(c) is an empty set.
(d) is a singleton.
- The probabilities of three events A, B and C are given by $P(A) = 0.6$, $P(B) = 0.4$ and $P(C) = 0.5$. If $P(A \cup B) = 0.8$, $P(A \cap C) = 0.3$, $P(A \cap B \cap C) = 0.2$, $P(B \cap C) = \beta$ and $P(A \cup B \cup C) = \alpha$, where $0.85 \leq \alpha \leq 0.95$, then β lies in the interval
(a) $[0.35, 0.36]$ (b) $[0.25, 0.35]$
(c) $[0.20, 0.25]$ (d) $[0.36, 0.40]$
- Let $r \in \{p, q, \sim p, \sim q\}$ be such that the logical statement $r \vee (\sim p) \Rightarrow (p \wedge q) \vee r$ is a tautology. Then r is equal to
(a) p (b) q (c) $\sim p$ (d) $\sim q$
- The domain of the function $f(x) = \frac{1}{4 - x^2} + \log_{10}(x^3 - x)$ is
(a) $(-2, -1) \cup (-1, 0) \cup (2, \infty)$
(b) $(1, 2) \cup (2, \infty)$
(c) $(-1, 0) \cup (1, 2) \cup (2, \infty)$
(d) $(-1, 0) \cup (1, 2) \cup (3, \infty)$
- If a variate assumes the values 0, 1, 2, ..., n , with frequencies ${}^nC_0, {}^nC_1, {}^nC_2, \dots, {}^nC_n$, then mean square deviation about the value $x = 0$ is
(a) $\frac{n(n-1)}{n}$ (b) $\frac{n^2(n-1)}{4}$
(c) $\frac{n(n+1)}{4}$ (d) $\frac{n(n+1)}{2}$
- If $f(x) = \int_{x^2}^{x^3} \frac{dt}{\log t}$, $x > 0$ then
(a) $f(x)$ is maximum at $x = 1$
(b) $f(x)$ is an increasing function $\forall x \in R^+$ only
(c) $f(x)$ is minimum at $x = 1$
(d) $f(x)$ is neither maximum nor minimum at $x = 1$ but an increasing function $\forall x \in R$
- Let R_1 and R_2 be two relations defined as follows :
 $R_1 = \{(a, b) \in R^2 : a^2 + b^2 \in Q\}$ and
 $R_2 = \{(a, b) \in R^2 : a^2 + b^2 \notin Q\}$, where Q is the set of all rational numbers. Then
(a) Neither R_1 nor R_2 is transitive.
(b) R_2 is transitive but R_1 is not transitive.
(c) R_1 is transitive but R_2 is not transitive.
(d) R_1 and R_2 are both transitive.
- Let ABC be a triangle with $A(-3, 1)$ and $\angle ACB = \theta$, $0 < \theta < \pi/2$. If the equation of the median through B is $2x + y - 3 = 0$ and the equation of angle bisector of C is $7x - 4y - 1 = 0$, then $\tan \theta$ is equal to
(a) $3/4$ (b) $4/3$ (c) 2 (d) $1/2$
- If $\int \frac{d\theta}{\cos^2 \theta (\tan 2\theta + \sec 2\theta)} = \lambda \tan \theta + 2 \log_e |f(\theta)| + C$, where C is a constant of integration, then the ordered pair $(\lambda, f(\theta))$ is equal to
(a) $(-1, 1 - \tan \theta)$ (b) $(-1, 1 + \tan \theta)$
(c) $(1, 1 + \tan \theta)$ (d) $(1, 1 - \tan \theta)$
- If $A = \begin{bmatrix} \cos^2 \alpha & \sin \alpha \cos \alpha \\ \sin \alpha \cos \alpha & \sin^2 \alpha \end{bmatrix}$;

$$B = \begin{bmatrix} \cos^2 \beta & \sin \beta \cos \beta \\ \sin \beta \cos \beta & \sin^2 \beta \end{bmatrix} \text{ are such that, } AB \text{ is}$$

a null matrix, then which of the following should necessarily be an odd integral multiple of $\pi/2$?

- (a) α (b) β (c) $\alpha - \beta$ (d) $\alpha + \beta$

13. Let $\vec{a} = \hat{i} - \hat{j} + 2\hat{k}$ and let $\vec{b}$ be a vector such that $\vec{a} \times \vec{b} = 2\hat{i} - \hat{k}$ and $\vec{a} \cdot \vec{b} = 3$. Then the projection of $\vec{b}$ on the vector $\vec{a} - \vec{b}$ is

- (a) $\frac{2}{\sqrt{21}}$ (b) $2\sqrt{\frac{3}{7}}$ (c) $\frac{2}{3}\sqrt{\frac{7}{3}}$ (d) $\frac{2}{3}$

14. If α and β are the roots of the equation $x^2 + px + 2 = 0$ and $1/\alpha$ and $1/\beta$ are the roots of the equation $2x^2 + 2qx + 1 = 0$, then

$\left(\alpha - \frac{1}{\alpha}\right)\left(\beta - \frac{1}{\beta}\right)\left(\alpha + \frac{1}{\beta}\right)\left(\beta + \frac{1}{\alpha}\right)$ is equal to

- (a) $\frac{9}{4}(9 + q^2)$ (b) $\frac{9}{4}(9 - q^2)$
(c) $\frac{9}{4}(9 + p^2)$ (d) $\frac{9}{4}(9 - p^2)$

15. The value of $\lim_{x \rightarrow 1} \frac{(x^2 - 1)\sin^2(\pi x)}{x^4 - 2x^3 + 2x - 1}$ is

- (a) $\pi^2/6$ (b) $\pi^2/3$ (c) $\pi^2/2$ (d) π^2

16. The solution of the differential equation $ydx - (x + 2y^2)dy = 0$ is $x = f(y)$. If $f(-1) = 1$, then $f(1)$ is equal to

- (a) 4 (b) 3 (c) 2 (d) 1

17. The integral

$$\int_2^4 \frac{\log x^2}{\log x^2 + \log(36 - 12x + x^2)} dx \text{ is equal to}$$

- (a) 1 (b) 6 (c) 2 (d) 4

18. Let $z \in \mathbb{C}$, the set of complex numbers. Then the equation, $2|z + 3i| - |z - i| = 0$ represents

- (a) a circle with radius $8/3$.
(b) a circle with diameter $10/3$.
(c) an ellipse with length of major axis $16/3$.
(d) an ellipse with length of minor axis $16/9$.

19. Let $[x]$ denote the greatest integer less than or equal

to x . Then $\lim_{x \rightarrow 0} \frac{\tan(\pi \sin^2 x) + (|x| - \sin(x[x]))^2}{x^2}$

- (a) equals 0 (b) equals π
(c) equals $\pi + 1$ (d) does not exist

20. Let A be a 3×3 real matrix such that

$$A \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}; A \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix} \text{ and } A \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix}.$$

If $X = (x_1, x_2, x_3)^T$ and I is an identity matrix of order

$$3, \text{ then the system } (A - 2I)X = \begin{pmatrix} 4 \\ 1 \\ 1 \end{pmatrix} \text{ has}$$

- (a) no solution
(b) infinitely many solutions
(c) unique solution
(d) exactly two solutions

SECTION-B (NUMERICAL VALUE TYPE)

Attempt any 5 questions out of 10.

21. Let $y = y(x)$ be the solution of the differential equation $dy = e^{ax+y} dx$; $\alpha \in \mathbb{N}$. If $y(\log_e 2) = \log_e 2$ and $y(0) = \log_e \left(\frac{1}{2}\right)$, then the value of α is equal to _____.

22. If b is the first term of an infinite G.P. whose sum is 5, then b lies in the interval (a, b) , then $a + b$ is equal to _____.

23. Let the mean and variance of four numbers 3, 7, x and y ($x > y$) be 5 and 10 respectively. Then the mean of four numbers $3 + 2x$, $7 + 2y$, $x + y$ and $x - y$ is _____.

24. For real numbers α and β , consider the following system of linear equations:

$$x + y - z = 2, x + 2y + \alpha z = 1, 2x - y + z = \beta.$$

If the system has infinite solutions, then $\alpha + \beta$ is equal to _____.

25. If the least and the largest real values of α , for which the equation $z + \alpha|z - 1| + 2i = 0$ ($z \in \mathbb{C}$ and $i = \sqrt{-1}$) has a solution, are p and q respectively, then $4(p^2 + q^2)$ is equal to _____.

26. Let a and b respectively be the points of local maximum and local minimum of the function $f(x) = 2x^3 - 3x^2 - 12x$. If A is the total area of the region bounded by $y = f(x)$, the x -axis and the lines $x = a$ and $x = b$, then $4A$ is equal to _____.

27. Let $\vec{a} = \hat{i} - \alpha\hat{j} + \beta\hat{k}$, $\vec{b} = 3\hat{i} + \beta\hat{j} - \alpha\hat{k}$ and $\vec{c} = -\alpha\hat{i} - 2\hat{j} + \hat{k}$, where α and β are integers. If $\vec{a} \cdot \vec{b} = -1$ and $\vec{b} \cdot \vec{c} = 10$, then $(\vec{a} \times \vec{b}) \cdot \vec{c}$ is equal to _____.

28. If the foot of the perpendicular drawn from the point (1, 0, 3) on a line passing through $(\alpha, 7, 1)$ is

$$\left(\frac{5}{3}, \frac{7}{3}, \frac{17}{3}\right), \text{ then } \alpha \text{ is equal to } \underline{\hspace{2cm}}.$$

29. In a hurdle race, a runner has probability p of jumping over a specific hurdle. Given that in 5 trials, the runner succeeded 3 times, the conditional probability that the runner had succeeded in the first trial is _____.

30. If every solution of the equation $3 \cos^2 x - \cos x - 1 = 0$ is a solution of the equation $a \cos^2 2x + b \cos 2x - 1 = 0$. Then the value of $(a + b)$ is equal to _____.

SOLUTIONS

1. (d): Total number of words = Number of words in which particular things are always together + never comes together

$$\therefore \frac{9!}{4!2!} = \frac{8!}{4!} + \text{ways when C's do not come together}$$

[Consider 2C's as only 1C, then $n = 8$]

- $\therefore$ Number of ways when 2C's never come together

$$= \frac{9!}{4!2!} - \frac{8!}{4!} = 7560 - 1680 = 5880$$

2. (d)

3. (a): The system is $(2 - \lambda)x_1 - 2x_2 + x_3 = 0$

$$2x_1 - (3 + \lambda)x_2 + 2x_3 = 0$$

$$-x_1 + 2x_2 - \lambda x_3 = 0$$

For non-trivial solution, the determinant of the coefficient matrix must vanish. Then

$$\begin{vmatrix} 2-\lambda & -2 & 1 \\ 2 & -3-\lambda & 2 \\ -1 & 2 & -\lambda \end{vmatrix} = 0$$

$$\Rightarrow (2 - \lambda)\{\lambda(3 + \lambda) - 4\} + 2\{-2\lambda + 2\} + 1\{4 - (3 + \lambda)\} = 0$$

$$\Rightarrow (2 - \lambda)(\lambda^2 + 3\lambda - 4) - 4\lambda + 4 + 1 - \lambda = 0$$

$$\Rightarrow (2 - \lambda)(\lambda^2 + 3\lambda - 4) - 5\lambda + 5 = 0$$

$$\Rightarrow (2 - \lambda)(\lambda - 1)(\lambda + 4) - 5(\lambda - 1) = 0$$

$$\Rightarrow (\lambda - 1)(-\lambda^2 - 2\lambda + 8 - 5) = 0$$

$$\Rightarrow (\lambda - 1)(\lambda^2 + 2\lambda - 3) = 0 \Rightarrow (\lambda - 1)^2(\lambda + 3) = 0$$

Thus, $\lambda = 1, 1, -3$

$\therefore$ Set of all λ 's contain 2 elements.

4. (b): We have, $P(A) = 0.6$, $P(B) = 0.4$, $P(C) = 0.5$, $P(A \cup B) = 0.8$, $P(A \cap C) = 0.3$, $P(A \cap B \cap C) = 0.2$, $P(B \cap C) = \beta$ and $P(A \cup B \cup C) = \alpha$, where $0.85 \leq \alpha \leq 0.95$.

$$\text{Consider, } P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(B \cap C) - P(A \cap C) + P(A \cap B \cap C)$$

$$\Rightarrow \alpha = 0.6 + 0.4 + 0.5 - P(A \cap B) - \beta - 0.3 + 0.2$$

$$\Rightarrow \alpha = 1.4 - P(A \cap B) - \beta \quad \dots(i)$$

$$\text{Consider, } P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$\Rightarrow 0.8 = 0.6 + 0.4 - P(A \cap B)$$

$$\Rightarrow P(A \cap B) = 0.2$$

$$\dots(ii)$$

From (i) and (ii), we get

$$\alpha = 1.4 - 0.2 - \beta \Rightarrow \alpha = 1.2 - \beta$$

$$\text{Now, as } 0.85 \leq \alpha \leq 0.95; 0.85 \leq 1.2 - \beta \leq 0.95$$

$$\Rightarrow -0.35 \leq -\beta \leq -0.25 \Rightarrow 0.25 \leq \beta \leq 0.35$$

5. (c)

6. (c): Given, $f(x) = \frac{1}{4-x^2} + \log_{10}(x^3 - x)$

For existence of the given function, $4 - x^2 \neq 0$

$$\therefore x \in \mathbf{R} - \{-2, 2\}$$

$$\text{Also, } x^3 - x > 0 \Rightarrow x(x-1)(x+1) > 0$$

The sign scheme is

$$\therefore x \in (-1, 0) \cup (1, \infty)$$

$$\therefore \text{Required domain} = x \in (-1, 0) \cup (1, 2) \cup (2, \infty)$$

7. (c): Mean square deviation about 0 = $\frac{\sum f_i x_i^2}{\sum f_i}$

$$= \frac{\sum_{r=0}^n r^2 {}^n C_r}{\sum_{r=0}^n (r(r-1) + r) {}^n C_r} = \frac{\sum_{r=0}^n r^2 {}^n C_r}{2^n}$$

$$= \frac{1}{2^n} \left[\sum_{r=0}^n r(r-1) {}^n C_r + \sum_{r=0}^n r {}^n C_r \right]$$

$$= \frac{1}{2^n} \left[\sum_{r=1}^n r(r-1) \frac{n}{r} {}^{n-1} C_{r-1} + \sum_{r=1}^n r \cdot \frac{n}{r} {}^{n-1} C_{r-1} \right]$$

$$= \frac{1}{2^n} \left[\sum_{r=2}^n r(r-1) \frac{n}{r} \cdot \frac{n-1}{r-1} {}^{n-2} C_{r-2} + n \sum_{r=1}^n {}^{n-1} C_{r-1} \right]$$

$$= \frac{1}{2^n} \left[n(n-1) \sum_{r=2}^n {}^{n-2} C_{r-2} + n \sum_{r=1}^n {}^{n-1} C_{r-1} \right]$$

$$= \frac{n(n-1)2^{n-2} + n \cdot 2^{n-1}}{2^n} = \frac{n(n-1)}{4} + \frac{n}{2} = \frac{n(n+1)}{4}$$

8. (d): $f(x) = \int_{x^2}^{x^3} \frac{1}{\log t} dt$

$$\therefore f(x) = \frac{3x^2}{3 \log x} - \frac{2x}{2 \log x} \quad (\text{using Leibnitz theorem})$$

$$\Rightarrow f'(x) = \frac{x^2 - x}{\log x} = \frac{x(x-1)}{\log x}$$

$$\therefore \text{For critical points } x(x-1) = 0 \Rightarrow x = 0, x = 1$$

Consider $x = 1$ as $x > 0$

Now, for $x < 1$, $f'(x) = (-ve) (-ve) = \text{positive } (> 0)$

and for $x > 1$, $f'(x) = (+ve) (+ve) = \text{positive } (> 0)$
 $\therefore f'(x)$ does not change its sign in the immediate nbd of $x = 1$. So $x = 1$ is neither the point of maxima nor minima. But $f'(x) > 0 \forall x < 1$ & $x > 1$.
 $\therefore f(x)$ is an increasing function $\forall x \in R$.

9. (a)

10. (b): Let M be the midpoint AC .

Let coordinates of $M \equiv (\alpha, 3 - 2\alpha)$ and $C \equiv \left(\beta, \frac{7\beta - 1}{4}\right)$.

Now, $\frac{\beta - 3}{2} = \alpha \Rightarrow 2\alpha = \beta - 3$... (i)

and $\frac{7\beta - 1}{4} + 1 = 6 - 4\alpha \Rightarrow 7\beta + 3 = 24 - 16\alpha$

$\Rightarrow 7\beta + 3 = 24 - 8 \times (\beta - 3)$ [Using (i)]

$\Rightarrow 15\beta = 48 - 3 \Rightarrow 15\beta = 45$

$\Rightarrow \beta = 3$ and $\frac{7\beta - 1}{4} = 5$


$\therefore$ Coordinates of $C \equiv (3, 5)$

Now, slope of $AC = \frac{5 - 1}{3 - 3} = \frac{2}{3}$

Slope of angle bisector of angle $C = 7/4$

Now, $\tan \frac{\theta}{2} = \frac{\left| \frac{7 - 2}{4 - 3} - \frac{1}{1 + \frac{14}{12}} \right|}{1 + \frac{14}{12}} = \frac{1}{2}$

$\therefore \tan \theta = \frac{2 \tan \frac{\theta}{2}}{1 - \tan^2 \frac{\theta}{2}} = \frac{2 \times \frac{1}{2}}{1 - \frac{1}{4}} = \frac{4}{3}$



$$\begin{aligned} 11. (b): \text{ Let } I &= \int \frac{\sec^2 \theta}{\left(\frac{2 \tan \theta}{1 - \tan^2 \theta} \right) + \left(\frac{1 + \tan^2 \theta}{1 - \tan^2 \theta} \right)} d\theta \\ &= \int \frac{\sec^2 \theta (1 - \tan^2 \theta)}{(1 + \tan \theta)^2} d\theta = \int \frac{\sec^2 \theta (1 - \tan \theta)}{1 + \tan \theta} d\theta \end{aligned}$$

Putting $\tan \theta = t \Rightarrow \sec^2 \theta d\theta = dt$, we get

$$I = \int \left(\frac{1-t}{1+t} \right) dt = \int \left(-1 + \frac{2}{1+t} \right) dt$$

$$= -t + 2 \log_e |(1+t)| + C = -\tan \theta + 2 \log_e |(1 + \tan \theta)| + C$$

$\therefore \lambda = -1$ and $f(\theta) = 1 + \tan \theta$

12. (c)

13. (a): Given, $\vec{a} = \hat{i} - \hat{j} + 2\hat{k}$, $\vec{a} \times \vec{b} = 2\hat{i} - \hat{k}$

and $\vec{a} \cdot \vec{b} = 3$

Projection of $\vec{b}$ on $(\vec{a} - \vec{b})$ is $= \frac{\vec{b} \cdot (\vec{a} - \vec{b})}{|\vec{a} - \vec{b}|}$

Now, $|\vec{a} \times \vec{b}|^2 + (\vec{a} \cdot \vec{b})^2 = |\vec{a}|^2 |\vec{b}|^2$

$$\Rightarrow 5 + 9 = 6 |\vec{b}|^2 \Rightarrow \frac{7}{3} = |\vec{b}|^2 \Rightarrow |\vec{b}| = \sqrt{\frac{7}{3}}$$

$$|\vec{a} - \vec{b}|^2 = |\vec{a}|^2 + |\vec{b}|^2 - 2(\vec{a} \cdot \vec{b}) = 6 + \frac{7}{3} - 2 \times 3 = \frac{7}{3}$$

$$\Rightarrow |\vec{a} - \vec{b}| = \sqrt{\frac{7}{3}}$$

Projection $\vec{b}$ on $(\vec{a} - \vec{b})$

$$= \frac{\vec{b} \cdot \vec{a} - |\vec{b}|^2}{|\vec{a} - \vec{b}|} = \frac{3 - \frac{7}{3}}{\sqrt{\frac{7}{3}}} = \frac{2}{3} \times \sqrt{\frac{3}{7}} = \frac{2}{\sqrt{21}}$$

14. (d)

$$15. (d): \lim_{x \rightarrow 1} \frac{(x^2 - 1) \sin^2 \pi x}{x^4 - 2x^3 + 2x - 1}$$

$$= \lim_{x \rightarrow 1} \frac{(x+1)(x-1) \frac{\sin^2(\pi x - \pi)}{\pi^2 (x-1)^2} \pi^2 (x-1)^2}{x^4 - 2x^3 + 2x - 1}$$

[$\because \sin^2(\theta - \pi) = \sin^2 \theta$]

$$= \lim_{x \rightarrow 1} \frac{\pi^2 (x+1)(x-1)^3}{x^4 - 2x^3 + 2x - 1} \left[\because \lim_{x \rightarrow 1} \left(\frac{\sin(\pi(x-1))}{\pi(x-1)} \right)^2 = 1 \right]$$

$$= 2\pi^2 \lim_{x \rightarrow 1} \frac{(x-1)^3}{x^4 - 2x^3 + 2x - 1} \quad \left(\frac{0}{0} \text{ form} \right)$$

$$= 2\pi^2 \lim_{x \rightarrow 1} \frac{3(x-1)^2}{4x^3 - 6x^2 + 2} \quad [\text{Using L'Hospital's rule}]$$

$$= 2\pi^2 \lim_{x \rightarrow 1} \frac{6(x-1)}{12x^2 - 12x} = 2\pi^2 \lim_{x \rightarrow 1} \frac{6(x-1)}{12x(x-1)}$$

$$= 2\pi^2 \times \frac{6}{12} = \pi^2$$

16. (b): We have, $y - (x + 2y^2) \frac{dy}{dx} = 0$

$$\Rightarrow y = (x + 2y^2) \frac{dy}{dx} \Rightarrow y \frac{dx}{dy} = x + 2y^2$$

$$\Rightarrow \frac{dx}{dy} + \left(-\frac{1}{y} \right) x = 2y \quad \dots (i)$$

$$\text{I.F.} = e^{\int -\frac{1}{y} dy} = e^{-\ln y} = y^{-1} = \frac{1}{y}$$

$\therefore$ The solution of the equation (i) is

$$x \left(\frac{1}{y} \right) = \int (2y) \times \frac{1}{y} dy + c \Rightarrow \frac{x}{y} = 2y + c \quad \dots (ii)$$

When $x = 1$ and $y = -1$, we get $c = 1$

The equation (ii) becomes $\frac{x}{y} = 2y + 1$... (iii)

Put $y = 1$ in (iii), we get $x = 2 + 1 = 3$

17. (a): Let $I = \int_2^4 \frac{\log x^2}{\log x^2 + \log(x-6)^2} dx$
 $= \int_2^4 \frac{\log x}{\log x + \log(6-x)} dx$ (use $\log x^2 = 2\log x$, $x > 0$) ... (i)

Using $\int_a^b f(x) dx = \int_a^b f(a+b-x) dx$, the above rewrite as
 $I = \int_2^4 \frac{\log(6-x)}{\log x + \log(6-x)} dx$... (ii)

Adding (i) & (ii), $2I = \int_2^4 \frac{\log x + \log(6-x)}{\log x + \log(6-x)} dx = \int_2^4 dx = 2$
 $\therefore I = 1$.

18. (a): We have, $2|z + 3i| - |z - i| = 0$
 $\Rightarrow 2|x + i(y + 3)| = |x + i(y - 1)|$ ($\because z = x + iy$)
 $\Rightarrow 2\sqrt{x^2 + (y + 3)^2} = \sqrt{x^2 + (y - 1)^2}$
 $\Rightarrow 4(x^2 + (y + 3)^2) = x^2 + (y - 1)^2$
 $\Rightarrow 3x^2 = y^2 - 2y + 1 - 4y^2 - 24y - 36$
 $\Rightarrow 3x^2 + 3y^2 + 26y + 35 = 0 \Rightarrow x^2 + y^2 + \frac{26}{3}y + \frac{35}{3} = 0$

This is the equation of circle with radius,

$$r = \sqrt{0^2 + \left(\frac{13}{3}\right)^2 - \frac{35}{3}} = \sqrt{\frac{64}{9}} = \frac{8}{3}$$

19. (d): Here,

R.H.L. = $\lim_{x \rightarrow 0^+} \frac{\tan(\pi \sin^2 x) + (|x| - \sin(x[x]))^2}{x^2}$
 $= \lim_{x \rightarrow 0^+} \frac{\tan(\pi \sin^2 x) + x^2}{x^2}$ ($\because x \rightarrow 0^+ \Rightarrow [x] = 0$)
 $= \lim_{x \rightarrow 0^+} \frac{\tan(\pi \sin^2 x)}{\pi \sin^2 x} \cdot \frac{\pi \sin^2 x}{x^2} + 1 = \pi + 1$

Now, L.H.L. = $\lim_{x \rightarrow 0^-} \frac{\tan(\pi \sin^2 x) + (-x + \sin x)^2}{x^2}$
 $(\because x \rightarrow 0^- \Rightarrow [x] = -1)$
 $= \lim_{x \rightarrow 0^-} \frac{\tan(\pi \sin^2 x)}{\pi \sin^2 x} \cdot \frac{\pi \sin^2 x}{x^2} + \left(-1 + \frac{\sin x}{x}\right)^2 = \pi$

$\therefore$ R.H.L. $\neq$ L.H.L.

So, limit does not exist.

20. (b): Let $A = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$; $A \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} a+b \\ d+e \\ g+h \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$

$$A \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} a+c \\ d+f \\ g+i \end{bmatrix} = \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}; A \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} c \\ f \\ i \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$$

$c = 1, f = 1, i = 2, a = -2, d = -1, g = -1,$
 $b = 3, e = 2, h = 1$

$$A - 2I = \begin{bmatrix} -2 & 3 & 1 \\ -1 & 2 & 1 \\ -1 & 1 & 2 \end{bmatrix} - 2 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} -4 & 3 & 1 \\ -1 & 0 & 1 \\ -1 & 1 & 0 \end{bmatrix}$$

$$(A - 2I)X = \begin{bmatrix} -4 & 3 & 1 \\ -1 & 0 & 1 \\ -1 & 1 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -4x_1 + 3x_2 + x_3 \\ -x_1 + x_3 \\ -x_1 + x_2 \end{bmatrix} = \begin{bmatrix} 4 \\ 1 \\ 1 \end{bmatrix}$$

$$\Rightarrow -4x_1 + 3x_2 + x_3 = 4$$

$$-x_1 + x_3 = 1$$

$$-x_1 + x_2 = 1$$

From above equations it can be said that it has infinitely solutions.

21. (2): The given differential equation is,

$$dy = (e^{\alpha x + y}) dx$$

$$\Rightarrow \int \frac{dy}{e^y} = \int e^{\alpha x} dx \Rightarrow -e^{-y} = \frac{e^{\alpha x}}{\alpha} + c$$

But $y(\log_e 2) = \log_e 2$ [Given]

$$\therefore \frac{-1}{e^{\log_e 2}} = \frac{e^{\alpha \log_e 2}}{\alpha} + c \Rightarrow c = \frac{-1}{2} - \frac{1}{\alpha} \cdot 2^\alpha \quad \dots (i)$$

Also, $y(0) = \log_e \left(\frac{1}{2}\right)$

$$\Rightarrow \frac{-1}{e^{\log_e 1/2}} = \frac{1}{\alpha} + c \Rightarrow c = -2 - \frac{1}{\alpha} \quad \dots (ii)$$

From (i) and (ii), we have $\frac{-1}{2} - \frac{1}{\alpha} \cdot 2^\alpha = -2 - \frac{1}{\alpha}$

$$\Rightarrow \frac{1}{\alpha} [1 - 2^\alpha] = -2 + \frac{1}{2} = \left(\frac{1 - 2^2}{2}\right) \Rightarrow \alpha = 2$$

22. (10): Let $b, br, br^2, \dots$ be an infinite G.P. with first term b , common ratio r and sum = 5.

$$\therefore 5 = \frac{b}{1-r}; |r| < 1 \Rightarrow \frac{b}{5} = 1-r \Rightarrow r = 1 - \frac{b}{5}$$

Now, $|r| < 1 \Rightarrow \left|1 - \frac{b}{5}\right| < 1 \Rightarrow -1 < 1 - \frac{b}{5} < 1$

$$\Rightarrow 0 < -\frac{b}{5} + 2 < 2 \Rightarrow 0 < \frac{10-b}{5} < 2$$

$$\Rightarrow 0 < -b + 10 < 10 \Rightarrow b < 10 \text{ and } b > 0 \text{ or } b \in (0, 10)$$

23. (12) : We have, $\frac{3+7+x+y}{4} = 5$
 $\Rightarrow x+y=10$... (i)

and also, we have $\frac{3^2+7^2+x^2+y^2}{4} - 25 = 10$
 $\therefore x^2+y^2=82$... (ii)

From (i), we have $(x^2+y^2) + 2xy = 100$
 $\Rightarrow xy=9 \Rightarrow y=9/x$ [Using (i)]

Thus, we have $x + \frac{9}{x} = 10$ [Using (i)]

$\Rightarrow x^2 - 10x + 9 = 0 \Rightarrow (x-9)(x-1) = 0 \Rightarrow x=1, 9$

Now, $x=1$ implies $y=9$ (rejected as $x > y$)

and $x=9$ implies $y=1$ (accepted)

$\Rightarrow 3+2x=21; 7+2y=9; x+y=10; x-y=8$

$\therefore$ Required mean = $\frac{21+9+10+8}{4} = 12$

24. (5) : $\therefore$ The system of equations has infinite solutions.

$\therefore$ Det $A = 0$, where A is the coefficient matrix.

$\Rightarrow \begin{vmatrix} 1 & 1 & -1 \\ 1 & 2 & \alpha \\ 2 & -1 & 1 \end{vmatrix} = 0$

$\Rightarrow 1(2+\alpha) - (1-2\alpha) - 1(-1-4) = 0 \Rightarrow \alpha = -2$

Also, $(\text{adj } A) \cdot B = 0$, where $B = \begin{bmatrix} 2 \\ 1 \\ \beta \end{bmatrix}$

Now, $\text{adj } A = \begin{bmatrix} 0 & 0 & 0 \\ -5 & 3 & 1 \\ -5 & 3 & 1 \end{bmatrix}$

$\therefore (\text{adj } A) \cdot B = \begin{bmatrix} 0 & 0 & 0 \\ -5 & 3 & 1 \\ -5 & 3 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \\ \beta \end{bmatrix}$

$\Rightarrow \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ \beta-7 \\ \beta-7 \end{bmatrix} \Rightarrow \beta-7=0 \Rightarrow \beta=7$

$\therefore \alpha + \beta = -2 + 7 = 5$

25. (10) : The given equation is $z + \alpha|z-1| + 2i = 0$

Putting $z = x + iy$, we get $x + iy + \alpha|x+iy-1| + 2i = 0$

$\Rightarrow x + \alpha\sqrt{(x-1)^2 + y^2} + i(y+2) = 0 + 0i$

$\Rightarrow y+2=0$ and $x + \alpha\sqrt{(x-1)^2 + y^2} = 0$

$\Rightarrow y = -2$ and $\alpha^2 = \frac{x^2}{x^2 - 2x + 5}$

Now, $\frac{x^2}{x^2 - 2x + 5} \in \left[0, \frac{5}{4}\right]$



PUBLIC NOTICE

JEE Main Session 2

Inviting Online Applications for Joint Entrance Examination (Main) – 2023 Session 2 - Reg.

In continuation of the Public Notice dated: **15 December 2022**, the National Testing Agency is now inviting online Application Forms for **Joint Entrance Examination (Main) – 2023 Session 2**. The schedule for the same is given below :

Examination / Session	Dates of Examination	Submission of Application Forms Online	Last Date for Payment of Application Fee Online
JEE (Main) – 2023 Session 2	06, 08, 10, 11, and 12 April 2023 (Reserve dates – 13, 15 April 2023)	14 February to 12 March 2023 (up to 09:00 P.M.)	12 March 2023 (up to 11:50 P.M.)

The dates of Advance Intimation of Examination City, downloading of Admit Cards, and declaration of Result will be displayed on JEE (Main) portal in due course.

The candidates who have applied and successfully paid the Examination Fee for JEE (Main) – 2023 Session 1 and wish to appear for JEE (Main) – 2023 Session 2, are required to log in with their previous Application Number and Password as provided in Session 1. They may only choose the **Paper, Medium of the Examination, State code of Eligibility and cities for Session 2 and pay the Examination Fees.**

The candidates are not allowed to fill more than one Application Form. Any candidate with more than one Application Number will be treated as UFM (Unfair Means), even if found at a later stage, and strict action will be taken against that Candidate.

The candidates who have already registered for Session 1 are not required to register again.

Those candidates who have not applied earlier, need to apply afresh as per the schedule given. For details, candidates may refer to the table below.

1. Existing candidates i.e. candidates registered for Session 1 and Paid Fee be considered for Session 2	Already registered candidates in Session 1 i.e. existing candidates for Session 2 : Candidates are allowed to opt for the following: i. Course (Paper) ii. Medium of Question Paper iii. State code of Eligibility iv. Examination Cities v. Fee Payment
2. For new candidates, the process for registration and application form filling in Session 2 will remain the same as followed in Session 1.	For new candidates in Session 2 i.e. not registered including Fee not paid for Session 1 : Process for registration and Application will remain the same as followed for Session 1.

The Candidates are advised to keep visiting the official websites of NTA (www.nta.ac.in) and (<https://jeemain.nta.nic.in/>) for the latest updates.

For further clarification related to JEE (Main) – 2023, the candidates can also contact **011- 40759000/011-69227700** or email at jeemain@nta.ac.in.

$$\therefore \alpha^2 \in \left[0, \frac{5}{4}\right] \Rightarrow \alpha \in \left[-\frac{\sqrt{5}}{2}, \frac{\sqrt{5}}{2}\right]$$

$$\therefore p = -\frac{\sqrt{5}}{2}; q = \frac{\sqrt{5}}{2}$$

$$\Rightarrow 4(p^2 + q^2) = 4\left(\frac{5}{4} + \frac{5}{4}\right) = 10$$

26. (114) : We have, $f(x) = 2x^3 - 3x^2 - 12x$

$$\Rightarrow f'(x) = 6x^2 - 6x - 12 = 6(x-2)(x+1)$$

$$\text{Now, } f'(x) = 0 \Rightarrow x = -1, 2$$

$$\text{and } f''(x) = 12x - 6 \Rightarrow f''(2) > 0 \Rightarrow b = 2$$

$$\text{and } f''(-1) < 0 \Rightarrow a = -1$$

$$\text{Consider, } f(x) = 0 \Rightarrow x(2x^2 - 3x - 12) = 0$$

$$\Rightarrow x = 0, \frac{3 \pm \sqrt{105}}{4}$$

$$\text{Now, } \frac{3 - \sqrt{105}}{4} < -1 \text{ and } \frac{3 + \sqrt{105}}{4} > 2$$

The only root lying between -1 & 2 is $x = 0$.

$$\text{Let, } A_1 = \left| \int_{-1}^0 (2x^3 - 3x^2 - 12x) dx \right|$$

$$= \left| \left[\frac{x^4}{2} - x^3 - 6x^2 \right]_{-1}^0 \right| = \left| 0 - \left(\frac{1}{2} + 1 - 6 \right) \right|$$

$$= 6 - \frac{3}{2} = \frac{9}{2} \text{ sq. units and}$$

$$A_2 = \left| \int_0^2 (2x^3 - 3x^2 - 12x) dx \right| = \left| \left[\frac{x^4}{2} - x^3 - 6x^2 \right]_0^2 \right|$$

$$= |8 - 8 - 24| = 24$$

$$\therefore A = A_1 + A_2 = 24 + \frac{9}{2} = \frac{57}{2} \text{ sq. units}$$

$$\Rightarrow 4A = 57 \times 2 = 114 \text{ sq. units}$$

27. (9) : We have,

$$\vec{a} \cdot \vec{b} = -1 \Rightarrow 3 - \alpha\beta - \alpha\beta = -1 \Rightarrow \alpha\beta = 2 \quad \dots(i)$$

$$\vec{b} \cdot \vec{c} = 10 \Rightarrow -3\alpha - 2\beta - \alpha = 10$$

$$\Rightarrow 4\alpha^2 + 2\beta + 10 = 0 \Rightarrow 4\alpha^2 + 4 + 10\alpha = 0 \text{ (Using (i))}$$

$$\Rightarrow (\alpha + 2)(4\alpha + 2) = 0$$

$$\Rightarrow \alpha = -2, \alpha = -\frac{1}{2} \text{ (Not possible, as } \alpha \in \mathbb{Z})$$

$$\therefore \text{From (i), } \beta = -1$$

$$\text{Now, } \vec{a} = \hat{i} + 2\hat{j} - \hat{k}, \vec{b} = 3\hat{i} - \hat{j} + 2\hat{k},$$

$$\vec{c} = 2\hat{i} - 2\hat{j} + \hat{k} \therefore (\vec{a} \times \vec{b}) \cdot \vec{c} = \begin{vmatrix} 1 & 2 & -1 \\ 3 & -1 & 2 \\ 2 & -2 & 1 \end{vmatrix}$$

$$= 1(-1 + 4) - 2(3 - 4) - 1(-6 + 2) = 3 + 2 + 4 = 9$$

28. (4) : Let $P \equiv (1, 0, 3)$ and $Q \equiv \left(\frac{5}{3}, \frac{7}{3}, \frac{17}{3}\right)$


Since, PQ is perpendicular to line l .

$$\therefore \left(1 - \frac{5}{3}\right)\left(\alpha - \frac{5}{3}\right) + \left(\frac{-7}{3}\right)\left(7 - \frac{7}{3}\right)$$

$$+ \left(3 - \frac{17}{3}\right)\left(1 - \frac{17}{3}\right) = 0$$

$$\Rightarrow \frac{-2\alpha}{3} + \frac{10}{9} - \frac{98}{9} + \frac{112}{9} = 0$$

$$\Rightarrow \frac{2\alpha}{3} = \frac{24}{9} \Rightarrow \alpha = 4$$



29. (0.6) : Let A denote the event that the runner succeeds exactly 3 times out of five and B denote the event that the runner succeeds on the first trial.

$$\text{Required probability} = P(B|A) = \frac{P(B \cap A)}{P(A)}$$

But $P(B \cap A) = P$ (succeeding in the first trial and exactly twice in other four trials)

$$= p({}^4C_2 p^2 (1-p)^2) = 6p^3 (1-p)^2$$

$$\text{and } P(A) = {}^5C_3 p^3 (1-p)^2 = 10p^3 (1-p)^2$$

$$\text{Thus, } P(B|A) = \frac{6p^3 (1-p)^2}{10p^3 (1-p)^2} = \frac{3}{5}$$

30. (13) : From 1st equation, $\cos x = \frac{1 \pm \sqrt{13}}{6}$

$$\text{Now, } \cos 2x = 2 \cos^2 x - 1 = \frac{2}{36} (1 + 13 \pm 2\sqrt{13}) - 1$$

$$= \frac{14 \pm 2\sqrt{13} - 18}{18} = \frac{\pm 2\sqrt{13} - 4}{18} = \frac{\pm \sqrt{13} - 2}{9}$$

$$\therefore \cos 2x_1 = \frac{\sqrt{13} - 2}{9} \text{ and } \cos 2x_2 = -\frac{(\sqrt{13} + 2)}{9}$$

Now from 2nd equation, $a \cos^2 2x + b \cos 2x - 1 = 0$

$$\therefore \cos 2x_1 \cdot \cos 2x_2 = -\frac{1}{a}$$

$$\Rightarrow -\frac{1}{a} = -\left(\frac{13 - 4}{81}\right) = -\frac{1}{9} \Rightarrow a = 9$$

$$\text{and } \cos 2x_1 + \cos 2x_2 = -\frac{b}{a} = -\frac{b}{9}$$

$$\Rightarrow -\frac{b}{9} = -\frac{4}{9} \Rightarrow b = 4 \therefore a + b = 13.$$



Recipe for Success

“If there is no struggle, there is no progress.”

– Frederick Douglass

CHAPTERWISE PRACTICE PAPER 2023

CUET (UG)

Exam on
21st to 31st
May 2023

General Instructions

This practice paper contains two sections i.e. Section A and Section B [B1 and B2].

Section A has 15 questions covering both i.e. Mathematics/Applied Mathematics which is compulsory for all candidates.

Section B1 has 30 questions from Mathematics out of which 20 questions need to be attempted.

Section B2 has 30 questions purely from Applied Mathematics out of which 20 questions need to be attempted.

SECTION A

1. If the points (a_1, b_1) , (a_2, b_2) and $(a_1 + a_2, b_1 + b_2)$ are collinear, then

- (a) $a_1 b_2 = a_2 b_1$ (b) $a_1 + a_2 = b_1 + b_2$
(c) $a_2 b_2 = a_1 b_1$ (d) $a_1 + b_1 = a_2 + b_2$

2. Let $A = \begin{bmatrix} 2 & 2 \\ 2 & 2 \end{bmatrix}$ and $B = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$. Consider the following statements.

Statement-I : $(A + B)^2 = A^2 + 2AB + B^2$ for the given matrices.

Statement-II : $AB = BA$ for the above matrices.

In the light of above statement, choose the correct answer from the options given below

- (a) Both statement I and statement I are true
(b) Both statement I and statement II are false
(c) Statement I is correct but statement II is false
(d) Statement I is incorrect but statement II is true
3. Match the following columns :

Column-I		Column-II	
P.	A is a real skew-symmetric matrix such that $A^2 + I = 0$, then AA' is equal to	1.	A
Q.	A is a matrix such that $A^2 = A$ if $(I + A)^n = I + \lambda A$, then	2.	O
R.	If for a matrix A, $A^2 = A$ and $B = I - A$, then $AB + BA + I - (I - A)^2$ equals	3.	$2^n - 1$
S.	If $A' = A$ and $B' = B$, then $(AB - BA)' - (BA - AB)$ equals	4.	I

- (a) P-1, Q-2, R-4, S-3 (b) P-2, Q-1, R-3, S-4
(c) P-3, Q-4, R-2, S-1 (d) P-4, Q-3, R-1, S-2
4. The maximum value of the function $f(x) = x^{-x}$, $x > 0$, is
- (a) $1/e$ (b) $e^{1/e}$ (c) e^{-e} (d) e

5. If tangent to the curve $y^2 = x^3$ at its point (m^2, m^3) is also normal to the curve at (M^2, M^3) , then what is the value of mM ?
- (a) $-4/9$ (b) $-2/9$ (c) $-1/3$ (d) 1

6. If $f'(x) = (x - 1)^3(x - 2)^8$, then

Statement-I : $f(x)$ has neither maximum nor minimum at $x = 2$.

Statement-II : $f'(x)$ changes sign from negative to positive at $x = 2$.

In the light of above statements, choose the correct answer from the options given below.

- (a) Both statement I and statement I are true
(b) Both statement I and statement II are false
(c) Statement I is correct but statement II is false
(d) Statement I is incorrect but statement II is true

7. For the differential equation, $\frac{dy}{dx} + \frac{y}{x} = \frac{1}{\sqrt{1+x^2}}$,

I. Integrating factor is x

II. $y = \frac{\sqrt{1+x^2}}{x} + \frac{c}{x}$ III. $y = \frac{x}{\sqrt{1+x^2}} + cx$

IV. Integrating factor is $1/x$

- (a) All are correct
(b) Only I and II are correct
(c) Only II and III are correct
(d) Only III is correct
8. $\int_0^{1.5} [x^2] dx$ is equal to ($[\cdot]$ denotes greatest integer function)
- (a) 2 (b) $2 - \sqrt{2}$ (c) $2 + \sqrt{2}$ (d) $\sqrt{2}$
9. The area bounded between the parabolas $x^2 = \frac{y}{4}$ and $x^2 = 9y$ and the straight line $y = 2$, is

- (a) $\frac{20\sqrt{2}}{3}$ sq. units (b) $10\sqrt{2}$ sq. units
(c) $20\sqrt{2}$ sq. units (d) $\frac{10\sqrt{2}}{3}$ sq. units

10. If $I = \int \frac{\sin x + \sin^3 x}{\cos 2x} dx = A \cos x + B \log|f(x)| + C$, then:

I. $A = \frac{1}{4}, B = -\frac{1}{\sqrt{2}}, f(x) = \frac{\sqrt{2} \cos x - 1}{\sqrt{2} \cos x + 1}$

II. $A = \frac{1}{2}, B = -\frac{3}{4\sqrt{2}}, f(x) = \frac{\sqrt{2} \cos x - 1}{\sqrt{2} \cos x + 1}$

III. $A = -\frac{1}{2}, B = \frac{3}{\sqrt{2}}, f(x) = \frac{\sqrt{2} \cos x + 1}{\sqrt{2} \cos x - 1}$

IV. $A = \frac{1}{2}, B = \frac{3}{4\sqrt{2}}, f(x) = \frac{\sqrt{2} \cos x + 1}{\sqrt{2} \cos x - 1}$

V. $A = \frac{-1}{2}, B = \frac{-3}{4\sqrt{2}}, f(x) = \frac{\sqrt{2} \cos x + 1}{\sqrt{2} \cos x - 1}$

- (a) All are correct
(b) Only I and II are correct
(c) II and IV are correct
(d) Only II and V are correct
11. If X is a random-variable with probability distribution given below:

X	0	1	2	3
$P(X)$	k	$3k$	$3k$	k

The value of k and its variance are

- (a) $1/8, 22/27$ (b) $1/8, 23/27$
(c) $1/8, 24/27$ (d) $1/8, 3/4$
12. Match the differential equation given in Column-II with their respective solution given in Column-I.

Column-I	Column-II
P. $y = ae^{3x} + be^{5x}$	1. $a + b[(y')^2 + yy''] = 0$
Q. $ax^2 + by^2 = 1$	2. $y'' + 16y = 0$
R. $y = a \sin(4x + b)$	3. $(y^2 - 2y + 2)y'' + (2y - 2)(y')^2 + e^y y' = 0$
S. $xe^y - y^2 = 2$	4. $y'' - 8y' + 15y = 0$

- (a) P - 2, Q - 3, R - 4, S - 1
(b) P - 3, Q - 4, R - 1, S - 2
(c) P - 4, Q - 1, R - 2, S - 3
(d) P - 1, Q - 2, R - 3, S - 4
13. If the chance that a ship arrives safely at a port is $9/10$, find the chance that out of 5 expected ships, atleast 4 will arrive safely at the port.
- (a) $\frac{91854}{100000}$ (b) $\frac{32805}{100000}$

- (c) $\frac{59049}{100000}$ (d) $\frac{26244}{100000}$

14. In solving the LPP : "minimize $f = 6x + 10y$ subject to constraints $x \geq 6, y \geq 2, 2x + y \geq 10, x \geq 0, y \geq 0$ " redundant constraints are

- (a) $x \geq 6, y \geq 2$
(b) $2x + y \geq 10, x \geq 0, y \geq 0$
(c) $x \geq 6$ (d) none of these

15. Corner points of the feasible region for an LPP are $(0, 2), (3, 0), (6, 0), (6, 8)$ and $(0, 5)$.

Let $F = 4x + 6y$ be the objective function.

The minimum value of F occurs at

- (a) $(0, 2)$ only (b) $(3, 0)$ only
(c) the mid-point of the line segment joining the points $(0, 2)$ and $(3, 0)$ only
(d) any point on the line segment joining the points $(0, 2)$ and $(3, 0)$

SECTION B1 (MATHEMATICS)

16. Which one of the following relations on R is an equivalence relation?

- (a) $aR_1b \Rightarrow |a| = |b|$ (b) $aR_2b \Rightarrow a \geq b$
(c) $aR_3b \Rightarrow a$ divides b (d) $aR_4b \Rightarrow a < b$

17. Solve $\sin^{-1}2x + \cos^{-1}2x + 2\tan^{-1}x = \pi$.

- (a) 1 (b) 0
(c) -1 (d) None of these

18. The only integral root of the equation

$$\begin{vmatrix} 2-y & 2 & 3 \\ 2 & 5-y & 6 \\ 3 & 4 & 10-y \end{vmatrix} = 0$$
 is

- (a) $y = 3$ (b) $y = 2$
(c) $y = 1$ (d) none of these

19. If $x + y = \tan^{-1}y$ and $\frac{d^2y}{dx^2} = f(y)\frac{dy}{dx}$, then $f(y) =$

- (a) $\frac{-2}{y^3}$ (b) $\frac{2}{y^3}$ (c) $\frac{1}{y}$ (d) $\frac{-1}{y}$

20. Given $0 \leq x \leq \frac{1}{2}$, then the value of

$$\tan \left[\sin^{-1} \left\{ \frac{x}{\sqrt{2}} + \frac{\sqrt{1-x^2}}{\sqrt{2}} \right\} - \sin^{-1}x \right] \text{ is}$$

- (a) 1 (b) $\sqrt{3}$ (c) -1 (d) $1/\sqrt{3}$

21. Match the following :

Column-I	Column-II
P. The total number of onto functions from the set $\{1, 2, 3, 4\}$ to the set $\{3, 4, 7\}$ is	1. 720

Q.	It set A and B contain 4 distinct elements, then total number of distinct function from A to B is	2.	5040
R.	If set A has 5 elements and set B has 6 elements, then number of injections that can be formed from A to B is	3.	256
S.	The number of bijective functions from set A to itself, when A contain 7 elements is	4.	36

- (a) P - 3, Q - 4, R - 2, S - 1
 (b) P - 4, Q - 3, R - 1, S - 2
 (c) P - 2, Q - 1, R - 3, S - 4
 (d) P - 1, Q - 2, R - 4, S - 3

22. The value of the integral $\int_2^4 \left(\frac{\log t}{t} \right) dt$ is equal to

- (a) $\frac{1}{2} (\log 2)^2$ (b) $\frac{5}{2} (\log 2)^2$
 (c) $\frac{3}{2} (\log 2)^2$ (d) $(\log 2)^2$

23. The area (in sq. units) bounded by the circle $x^2 + y^2 = 8$, the parabola $x^2 = 2y$ and the line $y = x$ in $y \geq 0$ is

- (a) $\frac{2}{3} + 2\pi$ (b) $\frac{2}{3} - 2\pi$ (c) $\frac{2}{3} + \pi$ (d) $\frac{2}{3} - \pi$

24. The solution of the differential equation

$$\frac{dy}{dx} + \frac{y}{2} \sec x = \frac{\tan x}{2y}, \text{ where } 0 \leq x < \frac{\pi}{2},$$

and $y(0) = 1$, is given by

- (a) $y^2 = 1 + \frac{x}{\sec x + \tan x}$ (b) $y = 1 + \frac{x}{\sec x + \tan x}$
 (c) $y = 1 - \frac{x}{\sec x + \tan x}$ (d) $y^2 = 1 - \frac{x}{\sec x + \tan x}$

25. The equation of the plane containing the line $2x - y + z - 3 = 0$, $3x + y + z = 5$ and at a distance of $1/\sqrt{6}$ from the point $(2, 1, -1)$ is

- I. $x + y + z - 3 = 0$
 II. $2x - y - z - 3 = 0$
 III. $2x - y + z - 3 = 0$
 IV. $62x + 29y + 19z - 105 = 0$

- (a) All are correct
 (b) Only II and IV are correct
 (c) Only III and IV are correct
 (d) Only I and II are correct

26. Match Column-I with Column-II and select the correct answer by choosing an appropriate option.

Column-I		Column-II	
P.	The slope of the curve $2y^2 = ax^2 + b$ at $(1, -1)$ is -1 , then	1.	$a - b = 2$
Q.	If (a, b) be the point on the curve $9y^2 = x^3$ where normal to the curve makes equal intercepts with the axes, then	2.	$a - b = 7/2$
R.	If the tangent at any point $(1, 2)$ on the curve $y = ax^2 + bx + 7/2$ be parallel to the normal at $(-2, 2)$ on the curve $y = x^2 + 6x + 10$, then	3.	$a - b = 4/3$
		4.	$a + b = 20/3$
		5.	$5a + 2b = 0$

- (a) P-1, Q-3,4, R-2, 5 (b) P-2, Q-3, 4, R-1, 5
 (c) P-3, Q-1, 5, R-2, 4 (d) P-4, Q-3, R-1, 2

27. **Statement-I** : A fair coin is tossed fixed number of times. If probability of getting 7 heads is equal to probability of getting 9 heads, then probability of getting 2 heads is equal to $\frac{15}{2^{12}}$.

Statement-II : If ${}^nC_x = {}^nC_y$, then either

$$x = y \text{ or } x + y = n \text{ and } P(X = r) = {}^nC_r p^r q^{n-r}.$$

In the light of above statements, choose the correct answer from the options given below.

- (a) Both statement I and statement I are true
 (b) Both statement I and statement II are false
 (c) Statement I is correct but statement II is false
 (d) Statement I is incorrect but statement II is true

28. Solve $\frac{dy}{dx} = (4x + y + 1)^2$

- (a) $\frac{1}{2} \tan^{-1} \left(\frac{4x+y}{2} \right) = x + c$
 (b) $\tan^{-1} \left(\frac{4x+y}{2} \right) = x + c$
 (c) $\frac{1}{2} \tan^{-1} \left(\frac{4x+y+1}{2} \right) = x + c$
 (d) $\tan^{-1} \left(\frac{4x+y-1}{2} \right) = x - c$

29. **Statement-I** : The area bounded by parabola $y = x^2 - 4x + 3$ and $y = 0$ is $4/3$ sq. units.

Statement-II : The area bounded by curves $y = f(x) \geq 0$ and $y = 0$ between ordinates $x = a$ and

$$x = b \text{ (where } b > a \text{) is } \int_a^b f(x) dx.$$

In the light of above statements, choose the correct answer from the options given below.

- (a) Both statement I and statement II are true
 (b) Both statement I and statement II are false
 (c) Statement I is correct but statement II is false
 (d) Statement I is incorrect but statement II is true

30. If $\vec{a}$ and $\vec{b}$ are two vectors and angle between them is θ , then

- I. $|\vec{a} \times \vec{b}|^2 + (\vec{a} \cdot \vec{b})^2 = |\vec{a}|^2 |\vec{b}|^2$
 II. $|\vec{a} \times \vec{b}| = (\vec{a} \cdot \vec{b})$, if $\theta = \pi/4$
 III. $\vec{a} \times \vec{b} = (\vec{a} \cdot \vec{b})\hat{n}$ (where $\hat{n}$ is a normal unit vector), if $\theta = \pi/4$
 IV. $(\vec{a} \times \vec{b}) \cdot (\vec{a} + \vec{b}) = 0$
 (a) All are correct
 (b) Only I and II are correct
 (c) II, III and IV are correct
 (d) I, II and IV are correct

31. Vectors $\vec{a}$ and $\vec{b}$ are inclined at an angle $\theta = 60^\circ$. If $|\vec{a}| = 1, |\vec{b}| = 2$, then $[(\vec{a} + 3\vec{b}) \times (3\vec{a} - \vec{b})]^2$ is equal to
 (a) 225 (b) 275 (c) 325 (d) 300

32. If $y = x \tan y$, then $\frac{dy}{dx} =$

- I. $\frac{\tan y}{x - x^2 - y^2}$ II. $\frac{y}{x - x^2 - y^2}$
 III. $\frac{\tan y}{y - x}$ IV. $\frac{\tan x}{x - y^2}$
 V. $\frac{\tan y}{1 - x \sec^2 y}$
 (a) All are correct (b) Only I and II are correct
 (c) II and V are correct (d) I, II and V are correct

33. The value of $\int \frac{dx}{\sqrt{x+3}\sqrt[3]{x}}$ is

- (a) $3\sqrt{x} + 3(\sqrt[3]{x}) - 6\sqrt[6]{x} + \log(\sqrt[6]{x} + 1) + C$
 (b) $2\sqrt{x} + 6(\sqrt[6]{x}) - 6\log(\sqrt[6]{x} + 1) + C$
 (c) $2\sqrt{x} - 3(\sqrt[3]{x}) + 6(\sqrt[6]{x}) - 6\log(\sqrt[6]{x} + 1) + C$
 (d) None of the above

34. Two tailors P and Q earn ₹350 and ₹450 per day respectively. Tailor P can stitch 6 shirts and 3 trousers while tailor Q can stitch 7 shirts and 3 trousers per day. How many days should each of them work, if it is desired to produce at least 51 shirts and 24 trousers at a minimum labour cost?

- (a) Minimize $Z = 350x + 450y$
 Subject to, $6x + 7y \geq 51, 3x + 3y \geq 24, x \geq 0, y \geq 0$

- (b) Minimize $Z = 350x + 450y$

Subject to, $6x + 7y \geq 51, 3x + 3y \geq 24, x \geq 0, y \geq 0$

- (c) Minimize $Z = 350x + 450y$

Subject to, $6x + 7y \geq 51, 3x + 3y \geq 24, x \geq 0, y \leq 0$

- (d) Minimize $Z = 350x + 450y$

Subject to, $6x + 7y \geq 51, 3x + 3y \geq 24, x \leq 0, y \leq 0$

35. If a, b, c are the roots of the equation $x^3 - 3x^2 + 3x + 7 = 0$,


then the value of $\begin{vmatrix} 2bc - a^2 & c^2 & b^2 \\ c^2 & 2ac - b^2 & a^2 \\ b^2 & a^2 & 2ab - c^2 \end{vmatrix}$ is

- (a) 9 (b) 27 (c) 81 (d) 0

Case Based MCQs

Case I : Read the following passage and answer the questions from 36 to 40.

Shreya got a rectangular parallelepiped shaped box and spherical ball inside it as return gift. Sides of the box are $x, 2x$, and $x/3$, while radius of the ball is r .



36. If S represents the sum of volume of parallelepiped and sphere, then S can be written as

- (a) $\frac{4x^3}{3} + \pi r^2$ (b) $\frac{2x^2}{3} + \frac{4}{3}\pi r^2$
 (c) $\frac{2x^3}{3} + \frac{4}{3}\pi r^3$ (d) $\frac{2}{3}x + \frac{4}{3}\pi r$

37. If sum of the surface areas of box and ball are given to be constant k^2 , then x is equal to

- (a) $\sqrt{\frac{k^2 - 4\pi r^2}{6}}$ (b) $\sqrt{\frac{k^2 - 4\pi r}{6}}$
 (c) $\sqrt{\frac{k^2 - 4\pi}{6}}$ (d) none of these

38. The radius of the ball, when S is minimum, is

- (a) $\sqrt{\frac{k^2}{54 + \pi}}$ (b) $\sqrt{\frac{k^2}{54 + 4\pi}}$
 (c) $\sqrt{\frac{k^2}{64 + 3\pi}}$ (d) $\sqrt{\frac{k^2}{4\pi + 3}}$

39. Relation between length of the box and radius of the ball can be represented as

- (a) $x = 2r$ (b) $x = r/2$ (c) $x = r/3$ (d) $x = 3r$

Success Guaranteed in CUET (UG)

Common University Entrance Test Under-Graduate Programmes
with **MTG CUET (UG) Practice Papers**


From the academic session 2022-23, admissions to Undergraduate Programmes in all central universities in the country will take place through Central University Entrance Test (CUET).

To help students with their preparation for the above test MTG is coming up with the book CUET 2023 which caters the Science Stream for the test. The MTG CUET (UG) – 2023 title covers 15 Practice papers which include following subjects in each practice paper :

Section IA Languages : English and Hindi

Section II Domain : Physics, Chemistry, Mathematics and Biology

Section III General Test : General Knowledge, Current Affairs, General Mental Ability, Numerical Ability, Quantitative Reasoning, Logical and Analytical Reasoning



₹ 800/-



Available at all leading book shops throughout the country.
For more information or for help in placing your order:
Visit www.mtg.in to buy online!
Call 0124-6601200 or e-mail: info@mtg.in

Visit
www.mtg.in
for latest offers
and to buy
online!

40. Minimum value of S is

- (a) $\frac{k^2}{2(3\pi + 54)^{2/3}}$ (b) $\frac{k}{(3\pi + 54)^{3/2}}$
 (c) $\frac{k^3}{3(4\pi + 54)^{1/2}}$ (d) none of these

Case II : Read the following passage and answer the questions from 41 to 45.

A factory has three machines A, B and C to manufacture bolts. Machine A manufacture 30%, machine B manufacture 20% and machine C manufacture 50% of the bolts respectively. Out of their respective outputs 5%, 2% and 4% are defective. A bolt is drawn at random from total production and it is found to be defective.



41. Probability that defective bolt drawn is manufactured by machine A, is
 (a) $\frac{4}{13}$ (b) $\frac{5}{13}$ (c) $\frac{6}{13}$ (d) $\frac{9}{13}$
 42. Probability that defective bolt drawn is manufactured by machine B, is
 (a) 0.3 (b) 0.1 (c) 0.2 (d) 0.4
 43. Probability that defective bolt drawn is manufactured by machine C, is
 (a) $\frac{16}{39}$ (b) $\frac{17}{39}$ (c) $\frac{20}{39}$ (d) $\frac{15}{39}$
 44. Probability that defective bolt is not manufactured by machine B, is
 (a) $\frac{35}{39}$ (b) $\frac{61}{39}$
 (c) $\frac{41}{39}$ (d) none of these
 45. Probability that defective bolt is not manufactured by machine C, is
 (a) 0.03 (b) 0.09 (c) 0.5 (d) 0.9

SECTION B2 (APPLIED MATHEMATICS)

16. The interval in which the function $f(x) = 2x^3 + 3x^2 - 12x + 5$ is strictly increasing is
 (a) $[-2, 1]$ (b) $(-\infty, -2] \cup [1, \infty)$
 (c) $(-\infty, -2) \cup (0, \infty)$ (d) None of these
 17. If $y = \frac{\ln x}{x}$, then $\frac{d^2 y}{dx^2}$ at $(x = 1)$ is equal to
 (a) 3 (b) -3 (c) 2 (d) -2

18. Two vessels A and B contain milk and water in the ratio 7 : 5 and 17 : 7 respectively. In what ratio mixtures from two vessels should be mixed to get a new mixture containing milk and water in the ratio 5 : 3?
 (a) 2 : 1 (b) 1 : 2 (c) 3 : 2 (d) 2 : 3

19. Price relative of rice is 110 in 2019 compared to 2018. If rice costs ₹ 16.50 per kg in 2019, what did it cost in 2018.

- (a) ₹ 14 per kg (b) ₹ 14.5 per kg
 (c) ₹ 15 per kg (d) ₹ 15.5 per kg

20. Match the following columns:

Column-I		Column-II	
(P)	Average Cost (AC)	(1)	$\frac{d}{dx} C(x)$
(Q)	Marginal Cost (MC)	(2)	$\frac{1}{x} (MC - AC)$
(R)	Marginal Average Cost (MAC)	(3)	$\frac{C(x)}{x}$
(S)	Marginal Revenues	(4)	$\frac{d}{dx} R(x)$

- (a) P-1, Q-3, R-2, S-4 (b) P-1, Q-3, R-4, S-2
 (c) P-3, Q-2, R-1, S-4 (d) P-3, Q-1, R-2, S-4

21. If matrix $A = \begin{bmatrix} a & 5 & c \\ -5 & b & 0 \\ d & 0 & 0 \end{bmatrix}$ is skew symmetric, then

the value of $2a + b + c + d$ is equal to

- (a) 0 (b) 1 (c) 10 (d) -10

22. If the corner points of the feasible region for an L.P.P. are (0, 3), (5, 0), (6, 0), (2, 3) and (0, 4), then minimum value of objective function $F = 3x + 5y$ occurs at
 (a) (0, 3) only
 (b) Both (0, 3) and (5, 0)
 (c) Mid point of (0, 3) and (5, 0)
 (d) Every point of the line segment joining (0, 3) and (5, 0)

23. The probability distribution of a discrete random variable X is given below:

X	20	15	10	5
$P(X)$	1/10	1/5	1/2	1/5

Then $E(X)$ is equal to

- (a) 5 (b) 11
 (c) 15 (d) None of these

24. $3^{16} \bmod 4$ is equal to
(a) 1 (b) 2 (c) 12 (d) 64
25. In what ratio must a shopkeeper mix two types of oranges worth ₹ 30 per kg and ₹ 45 per kg respectively so as to get a mixture at ₹ 40 per kg?
(a) 3 : 2 (b) 1 : 2 (c) 2 : 1 (d) 2 : 3
26. Solve for x : $3(2 - x) \geq 2(1 - x)$
(a) $x \geq 4$ (b) $x \geq 5$ (c) $x \leq 4$ (d) $x \leq 3$
27. Consider the following statements:
Statement I: The measurable characteristic of sample is called parameter.
Statement II: The measurable characteristic of population is called statistic.
In the light of above statements, choose the correct answer from the options give below.
(a) Both Statement I and Statement II are correct.
(b) Both Statement I and Statement II are incorrect.
(c) Statement I is correct but statement II is incorrect.
(d) Statement I is incorrect but Statement II is correct.
28. If Laspeyre's index number = 196 and Paasche's index number = 64, then Fisher's ideal index number is
(a) 28 (b) 56 (c) 106 (d) 112
29. Which of the following is not a component of time series.
(a) Seasonal (b) Cyclical
(c) Irregular (d) Variance
30. The critical points for the function $f(x) = x^3 - 3x^2 + 3x + 4$ are
(a) 1, $1/3$ (b) 1, 3 (c) 1 only (d) 3 only
31. In the given two columns Column II represent the interval of set given in Column I. Match the following columns by choosing appropriate option.

Column I		Column II	
(P)	$\{x : x \in \mathbf{R}, x \leq 2\}$	(1)	$[2, \infty)$
(Q)	$\{y : y \in \mathbf{R}, y \geq 2\}$	(2)	$[-2, 2]$
(R)	$\{z : z \in \mathbf{R}, -2 \leq z \leq 2\}$	(3)	$(-\infty, 2]$
(S)	$\{a : a \in \mathbf{R}, a > 2\}$	(4)	$(2, \infty)$

- (a) P-3, Q-1, R-2, S-4 (b) P-1, Q-3, R-2, S-4
(c) P-4, Q-1, R-2, S-3 (d) P-3, Q-4, R-2, S-1
32. Rakesh and Suresh invest in a business in the ratio 4 : 5. If 10% of the profit goes for charity purpose and Rakesh's share of profit is ₹ 18000. Find the total profit.

- (a) ₹ 54000 (b) ₹ 45000
(c) ₹ 36000 (d) None of these
33. If the total cost function is given by $(9x - 6x^2 + 5x^3)$, then marginal average cost function (MAC) is given by
(a) $15x^2 - 12x + 9$ (b) $30x - 12$
(c) $-6 + 10x$ (d) None of these
34. A firm anticipates an expenditure of ₹ 500000 for plant modernization at end of 10 years from now. How much should the company deposit at the end of each year into a sinking fund earning interest 5% per annum? (Given: $(1.05)^{10} = 1.629$)
(a) ₹ 39745 (b) ₹ 32435
(c) ₹ 37725 (d) ₹ 42135
35. Consider the following statements:
I. A non singular matrix has a non zero determinant
II. If A is an involuntary matrix then $A^2 = A$
III. A matrix has all elements zero except its diagonal is triangular.
IV. For a orthogonal matrix, $A = A^T$
V. We can not calculate the determinant of a non square matrix.
Which of the above statements is/are true.
(a) I only (b) I, III and V
(c) I, II, IV and V (d) I and V
36. Which of the following is not a type of statistical inferences?
(a) Point Estimation (b) Interval Estimation
(c) Data Estimation (d) Hypothesis Testing
37. Solve the following linear programming problem:
Maximize $Z = 60x + 15y$ subject to the constraints $x + y \leq 50$, $3x + y \leq 90$, $x, y \geq 0$. The maximum value of Z is
(a) 1700 (b) 1800 (c) 1750 (d) 1850
38. A dining table costing ₹ 36000 has a useful life of 15 years. If annual depreciation is ₹ 2000, find its scrap value.
(a) ₹ 6000 (b) ₹ 5400 (c) ₹ 4500 (d) ₹ 4200
39. Suppose that $x = t - t^4$, $y = t^2 - t^3$
Statement I: $\frac{dy}{dx} = \frac{2t - 3t^2}{1 - 4t^3}$
Statement II: $\frac{dy}{dx}$ at (0, 0) is $\frac{1}{2}$ or 0
(a) Both Statement I and Statement II are correct.
(b) Both Statement I and Statement II are incorrect.
(c) Statement I is correct but statement II is incorrect.
(d) Statement I is incorrect but Statement II is correct.

40. Two pipes A and B together can fill a tank in 10 minutes. If pipe A take 15 minutes less than B to fill the tank alone, then find the time taken by pipe B to fill the tank alone.

(a) 30 min (b) 26 min
(c) 24 min (d) None of these

41. Find the mean, and standard deviation of the probability distribution:

X	2	3	4
$P(X)$	0.2	0.5	0.3

(a) 3.1, 0.49 (b) 3.1, 0.7 (c) 2.9, 0.49 (d) 2.9, 0.7

42. Find the remainder when

$672 + 541 + 383 + 295 + 101 + 86$ is divided by 3.

(a) 0 (b) 1
(c) 2 (d) Can't determined

43. A boat can row upstream at 10km/h and downstream at 18 km/h. Then

I. Speed of boat in still water is 14 km/h.
II. Speed of boat in still water is 8 km/h.
III. Speed of the stream is 4 km/h.
IV. Speed of the stream is 7 km/h.

Which of the above statement is/are true?

(a) I and III (b) I and IV
(c) II and III (d) II and IV

44. A person wishes to purchase a house for ₹ 5200000 with a down payment of ₹ 700000 and balance in EMI for 25 years. If bank charges 9% per annum compounded monthly, then calculate the EMI.

(Given $(1.0075)^{300} = 9.4084$)

(a) ₹35264 (b) ₹36257
(c) ₹ 37764 (d) ₹ 39241

45. Which of the following is test of adequacy of index numbers?

(a) Time reversal test (b) Factor reversal test
(c) Circular test (d) All of these

SOLUTIONS

SECTION A

1. (a) : The given points are collinear.

$$\therefore \frac{1}{2} \begin{vmatrix} a_1 & b_1 & 1 \\ a_2 & b_2 & 1 \\ a_1 + a_2 & b_1 + b_2 & 1 \end{vmatrix} = 0$$

Applying $R_2 \rightarrow R_2 - R_1$, $R_3 \rightarrow R_3 - R_1$, we get

$$\begin{vmatrix} a_1 & b_1 & 1 \\ a_2 - a_1 & b_2 - b_1 & 0 \\ a_2 & b_2 & 0 \end{vmatrix} = 0$$

Expanding along C_3 , we get $b_2(a_2 - a_1) - a_2(b_2 - b_1) = 0$
 $\Rightarrow -a_1b_2 + a_2b_1 = 0 \Rightarrow a_1b_2 = a_2b_1$

$$2. (a) : \text{Here, } AB = \begin{bmatrix} 2 & 2 \\ 2 & 2 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 0+2 & 2+0 \\ 0+2 & 2+0 \end{bmatrix} = \begin{bmatrix} 2 & 2 \\ 2 & 2 \end{bmatrix}$$

$$\text{And } BA = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 2 & 2 \\ 2 & 2 \end{bmatrix} = \begin{bmatrix} 0+2 & 0+2 \\ 2+0 & 2+0 \end{bmatrix} = \begin{bmatrix} 2 & 2 \\ 2 & 2 \end{bmatrix}$$

Thus, $AB = BA$.

Consequently $(A + B)^2 = A^2 + AB + BA + B^2$
 $= A^2 + AB + AB + B^2 = A^2 + 2AB + B^2$.

3. (d)

4. (b) : $f(x) = x^{-x}$, $x > 0$

$$\Rightarrow f(x) = e^{-x \log x} \Rightarrow f'(x) = e^{-x \log x} \times \frac{d}{dx}(-x \log x)$$

$$\Rightarrow f'(x) = x^{-x}(-1 - \log x) = -x^{-x}(1 + \log x)$$

For critical points, put $f'(x) = 0$

$$\Rightarrow -x^{-x}(1 + \log x) = 0 \Rightarrow 1 + \log x = 0$$

$[\because x^{-x} \neq 0, \text{ for } x > 0]$

$$\Rightarrow \log x = -1 \Rightarrow x = e^{-1} \text{ is the only critical point.}$$

$$\text{Now, } f''(x) = -x^{-x} \left[\frac{1}{x} - (1 + \log x)^2 \right]$$

$$\therefore f''(e^{-1}) = -(e^{-1})^{-e^{-1}} [e - (1 - 1)^2] = -e^{1/e} \times e = -e^{1+1/e}$$

$$\Rightarrow f''(e^{-1}) < 0$$

$[\because e^x > 0 \text{ for all } x > 0]$

So, $x = e^{-1}$ is the point of maxima.

$$\therefore \text{Maximum value} = (e^{-1})^{-e^{-1}} = e^{1/e}$$

5. (a) : We have, $y^2 = x^3 \Rightarrow 2y \frac{dy}{dx} = 3x^2 \Rightarrow \frac{dy}{dx} = \frac{3x^2}{2y}$

Given that, slope of tangent at (m^2, m^3)

= slope of normal at (M^2, M^3)

$$\Rightarrow \left(\frac{dy}{dx} \right)_{(m^2, m^3)} = \frac{-1}{\left(\frac{dy}{dx} \right)_{(M^2, M^3)}}$$


$$\Rightarrow \frac{3m^4}{2m^3} = -\frac{2M^3}{3M^4} \Rightarrow mM = -\frac{4}{9}$$

6. (c) : It is clear from figure that $f'(x)$ has no sign change at $x = 2$.

Hence, $f(x)$ has neither maximum nor minimum at $x = 2$.

7. (b) : Given $\frac{dy}{dx} + \frac{y}{x} = \frac{1}{\sqrt{1+x^2}}$

It is linear differential equation of the form $\frac{dy}{dx} + Py = Q$, where P and Q are functions of x



Here, I.F. = $e^{\int \frac{1}{x} dx} = x$

∴ Required solution is

$$y \cdot x = \int \frac{x}{\sqrt{1+x^2}} dx + c = \frac{1}{2} \int \frac{2x}{\sqrt{1+x^2}} dx + c$$

$$\Rightarrow y \cdot x = \sqrt{1+x^2} + c \Rightarrow y = \frac{\sqrt{1+x^2}}{x} + \frac{c}{x}$$

8. (b): $\int_0^{1.5} [x^2] dx = \int_0^1 [x^2] dx + \int_1^{\sqrt{2}} [x^2] dx + \int_{\sqrt{2}}^{1.5} [x^2] dx$

$$= \int_0^1 0 dx + \int_1^{\sqrt{2}} 1 dx + \int_{\sqrt{2}}^{1.5} 2 dx = 0 + [x]_1^{\sqrt{2}} + [2x]_{\sqrt{2}}^{1.5}$$

$$= \sqrt{2} - 1 + 3 - 2\sqrt{2} = 2 - \sqrt{2}$$


9. (a): Required area

$$= 2 \int_0^2 \left(3\sqrt{y} - \frac{\sqrt{y}}{2} \right) dy$$

$$= 2 \left[\frac{3y^{3/2}}{3/2} - \frac{1}{2} \cdot \frac{y^{3/2}}{3/2} \right]_0^2$$

$$= 2 \left[2(2)^{3/2} - \frac{1}{3}(2)^{3/2} - 0 \right]$$

$$= 2 \left(4\sqrt{2} - \frac{2\sqrt{2}}{3} \right) = \frac{20}{3} \sqrt{2} \text{ sq. units.}$$



10. (c): Since the integral is odd w.r.t. sine, we put $\cos x = t$, so $\cos 2x = 2t^2 - 1$ and $dt = -\sin x dx$.

$$\therefore I = \int \frac{t^2 - 2}{2t^2 - 1} dt = \frac{1}{2} \int \frac{2t^2 - 4}{2t^2 - 1} dt = \frac{1}{2} \int dt - \frac{3}{2} \int \frac{dt}{2t^2 - 1}$$

$$= \frac{1}{2} t - \frac{3}{2\sqrt{2}} \cdot \frac{1}{2} \log \left| \frac{\sqrt{2}t - 1}{\sqrt{2}t + 1} \right| + C$$

$$= \frac{1}{2} \cos x - \frac{3}{4\sqrt{2}} \log \left| \frac{\sqrt{2} \cos x - 1}{\sqrt{2} \cos x + 1} \right| + C$$

So $A = \frac{1}{2}, B = -\frac{3}{4\sqrt{2}}, f(x) = \frac{\sqrt{2} \cos x - 1}{\sqrt{2} \cos x + 1}$

or $A = \frac{1}{2}, B = \frac{3}{4\sqrt{2}}, f(x) = \frac{\sqrt{2} \cos x + 1}{\sqrt{2} \cos x - 1}$

11. (d): We know, $\Sigma P(X) = 1$

$$\Rightarrow k + 3k + 3k + k = 1 \Rightarrow 8k = 1 \Rightarrow k = 1/8$$

Now,

X	$P(X)$	$XP(X)$	$X^2P(X)$
0	1/8	0	0
1	3/8	3/8	3/8
2	3/8	6/8	12/8
3	1/8	3/8	9/8
Total		12/8	24/8

$$\text{Variance}(X) = \Sigma X^2 P(X) - [\Sigma X P(X)]^2$$

$$= \frac{24}{8} - \frac{144}{64} = \frac{192 - 144}{64} = \frac{48}{64} = \frac{3}{4}$$

12. (c)

13. (a): Here, $p = \frac{9}{10}, q = 1 - p = 1 - \frac{9}{10} = \frac{1}{10}$

Probability that atleast 4 will arrive safely at the port
= Probability that exactly 4 ships arrive safely +
Probability that exactly 5 ships arrive safely

$$= {}^5C_4 \times \left(\frac{9}{10} \right)^4 \left(\frac{1}{10} \right) + {}^5C_5 \left(\frac{9}{10} \right)^5$$

$$= 5 \times \frac{9^4}{10^5} + \frac{9^5}{10^5} = \frac{14 \times 9^4}{10^5} = \frac{91854}{100000}$$

14. (b): When $x \geq 6$ and $y \geq 2$, then $2x + y \geq 2 \times 6 + 2$,
i.e., $2x + y \geq 14$.

Hence, $x \geq 0, y \geq 0$ and $2x + y \geq 10$ are automatically
satisfied by every point of the region

$$\{(x, y) : x \geq 6\} \cap \{(x, y) : y \geq 2\}$$

15. (d): Construct the following table of values of
objective function :

Corner Point	Value of $F = 4x + 6y$
(0, 2)	$4 \times 0 + 6 \times 2 = 12$
(3, 0)	$4 \times 3 + 6 \times 0 = 12$
(6, 0)	$4 \times 6 + 6 \times 0 = 24$
(6, 8)	$4 \times 6 + 6 \times 8 = 72$
(0, 5)	$4 \times 0 + 6 \times 5 = 30$

← minimum

← maximum

Since the minimum value (F) = 12 occurs at two distinct
corner points, it occurs at every point of the segment
joining these two points.

SECTION B1 (MATHEMATICS)

16. (a): Reflexive : $a \in R, aR_1 a \Rightarrow |a| = |a|$

∴ R_1 is reflexive.

Symmetric : $a, b \in R$, then $aR_1 b \Rightarrow |a| = |b|$

$$\Rightarrow |b| = |a| \Rightarrow bR_1 a$$

∴ R_1 is symmetric

Transitive : Let $a, b, c \in R$, then $aR_1 b \Rightarrow |a| = |b|$... (i)

and $bR_1 c \Rightarrow |b| = |c|$... (ii)

From (i) and (ii), $|a| = |c| \Rightarrow aR_1 c$

∴ R_1 is transitive $\Rightarrow R_1$ is an equivalence relation.

17. (d): Given trigonometrical equation is

$$\sin^{-1} 2x + \cos^{-1} 2x + 2 \tan^{-1} x = \pi$$

$$\Rightarrow \frac{\pi}{2} + 2 \tan^{-1} x = \pi \Rightarrow 2 \tan^{-1} x = \frac{\pi}{2}$$

$\Rightarrow \tan^{-1} x = \frac{\pi}{4} \Rightarrow x = \tan\left(\frac{\pi}{4}\right) = 1$ but $\sin^{-1} 2$ is not defined $x = 1$.

18. (c) : Since, $\begin{vmatrix} 2-y & 2 & 3 \\ 2 & 5-y & 6 \\ 3 & 4 & 10-y \end{vmatrix} = 0$

$\Rightarrow (2-y)[(5-y)(10-y) - 24] - 2[20 - 2y - 18] + 3[8 - 15 + 3y] = 0$

$\Rightarrow -y^3 + 17y^2 - 43y + 27 = 0$

Only $y = 1$ satisfies this equation.

19. (b) : $x + y = \tan^{-1} y$

$\Rightarrow 1 + \frac{dy}{dx} = \frac{1}{1+y^2} \frac{dy}{dx} \Rightarrow \frac{dy}{dx} = -\frac{1}{y^2} - 1$

$\frac{d^2 y}{dx^2} = -\frac{(-2)}{y^3} \frac{dy}{dx} = \frac{2}{y^3} \frac{dy}{dx}$

20. (a) : Let $y = \tan\left[\sin^{-1}\left\{\frac{x}{\sqrt{2}} + \frac{\sqrt{1-x^2}}{\sqrt{2}}\right\} - \sin^{-1} x\right]$

Putting $x = \sin\theta$, we get

$y = \tan\left[\sin^{-1}\left[\frac{\sin\theta}{\sqrt{2}} + \frac{\sqrt{1-\sin^2\theta}}{\sqrt{2}}\right] - \sin^{-1}(\sin\theta)\right]$

$= \tan\left[\sin^{-1}\left[\frac{\sin\theta + \cos\theta}{\sqrt{2}}\right] - \theta\right]$

$= \tan\left[\sin^{-1}\left\{\sin\theta\cos\frac{\pi}{4} + \cos\theta\sin\frac{\pi}{4}\right\} - \theta\right]$

$= \tan\left[\sin^{-1}\left\{\sin\left(\theta + \frac{\pi}{4}\right)\right\} - \theta\right] = \tan\frac{\pi}{4} = 1$

21. (b) : (P) Since, we know that if A and B are two sets consisting of m and n elements respectively such that $1 \leq n \leq m$, then number of onto function from A to B is

$\sum_{r=1}^n (-1)^{n-r} {}^nC_r r^m$ if $m \geq n$. Here, $m = 4$, $n = 3$

$\therefore$ Total number of onto functions

$= \sum_{r=1}^3 (-1)^{3-r} {}^3C_r r^4 = {}^3C_1 - {}^3C_2 \cdot 2^4 + {}^3C_3 \cdot 3^4$

$= 3 - 48 + 81 = 36$

(Q) We know that if A and B are two sets consisting of m and n elements respectively, then total number of functions from A to B is n^m . Here, $n = m = 4$

$\therefore$ Total number of functions $= 4^4 = 256$

(R) We know that if A and B are two sets consisting of m and n elements respectively then total number of injections that can be formed are ${}^nP_m$, $n \geq m$.

Here, $m = 5$ and $n = 6$

$\therefore$ Total number of one one functions from

$A \text{ to } B = {}^6P_5 = \frac{6!}{(6-5)!} = \frac{6 \times 5 \times 4 \times 3 \times 2 \times 1}{1!} = 720$

(S) We know that if A and B are two sets consisting of m and n elements respectively, then number of bijections that can be formed are $m!$, if $m = n$.

$\therefore$ Total number of bijective functions from set A to itself $= 7! = 7 \times 6 \times 5 \times 4 \times 3 \times 2 \times 1 = 5040$


22. (c) : Let $\log t = x \Rightarrow \frac{1}{t} dt = dx$

$\int_{\log 2}^{\log 4} x dx = \left[\frac{x^2}{2}\right]_{\log 2}^{\log 4} = \frac{1}{2}[(\log 2^2)^2 - (\log 2)^2]$

$= \frac{1}{2}[(\log 2^2 - \log 2)(\log 2^2 + \log 2)]$

$= \frac{1}{2}[(\log 2)(3 \log 2)] = \frac{3}{2}(\log 2)^2$

23. (a) : We have, $x^2 + y^2 = 8$, $x^2 = 2y$ and $y = x$



Required area = area of shaded region $AOBC$

$= \int_{-2}^2 \sqrt{8-x^2} dx - \int_{-2}^0 \frac{x^2}{2} dx - \int_0^2 x dx$

$= 2 \int_0^2 \sqrt{8-x^2} dx - \left[\frac{x^3}{2 \times 3}\right]_{-2}^0 - \left[\frac{x^2}{2}\right]_0^2$

$\left[\because \sqrt{8-x^2} \text{ is an even function}\right]$

$= 2 \left[\frac{x}{2} \sqrt{8-x^2} + \frac{8}{2} \sin^{-1} \frac{x}{2\sqrt{2}} \right]_0^2 - \frac{1}{6}(0+8) - \frac{1}{2}(4-0)$

$= 2 \left[\frac{2}{2}(2) + 4 \left(\frac{\pi}{4} \right) - 0 - 0 \right] - \frac{4}{3} - 2 = \left(2\pi + \frac{2}{3} \right) \text{sq. units}$

24. (d) : We have, $\frac{dy}{dx} + \frac{y}{2} \sec x = \frac{\tan x}{2y}$

$\Rightarrow 2y \frac{dy}{dx} + y^2 \sec x = \tan x \quad \dots(i)$

Put $y^2 = t \Rightarrow 2y \frac{dy}{dx} = \frac{dt}{dx}$

$\therefore$ (i) becomes, $\frac{dt}{dx} + t \sec x = \tan x$

I.F. $= e^{\int \sec x dx} = e^{\log(\sec x + \tan x)} = \sec x + \tan x$

∴ Solution is given by

$$t(\sec x + \tan x) = \int \tan x(\sec x + \tan x) dx$$

$$= \int \tan x \sec x dx + \int \tan^2 x dx$$

$$\Rightarrow t(\sec x + \tan x) = \sec x + \tan x - x + c$$

$$\Rightarrow t = 1 - \left(\frac{x-c}{\sec x + \tan x} \right) \Rightarrow y^2 = 1 - \left(\frac{x-c}{\sec x + \tan x} \right)$$

$$\text{Now, } y(0) = 1 \Rightarrow 1 = 1 - (0 - c) \Rightarrow c = 0$$

$$\therefore \text{ Particular solution is } y^2 = 1 - \frac{x}{\sec x + \tan x}$$

25. (c) : The plane is

$$2x - y + z - 3 + \lambda(3x + y + z - 5) = 0 \quad \dots(i)$$

$$\Rightarrow (2 + 3\lambda)x + (\lambda - 1)y + (\lambda + 1)z - (5\lambda + 3) = 0$$

$$\text{Its distance from } (2, 1, -1) \text{ is } \frac{1}{\sqrt{6}}. \quad [\text{Given}]$$

$$\therefore \frac{|4 + 6\lambda + \lambda - 1 - \lambda - 1 - 5\lambda - 3|}{\sqrt{(2 + 3\lambda)^2 + (\lambda - 1)^2 + (\lambda + 1)^2}} = \frac{1}{\sqrt{6}}$$

$$\Rightarrow 6(\lambda - 1)^2 = 11\lambda^2 + 12\lambda + 6$$

$$\Rightarrow (5\lambda + 24)\lambda = 0 \Rightarrow \lambda = 0, \lambda = -\frac{24}{5}$$

$$\text{For } \lambda = 0, 2x - y + z - 3 = 0$$

$$\text{For } \lambda = -\frac{24}{5}, 5(2x - y + z - 3) - 24(3x + y + z - 5) = 0$$

$$\Rightarrow 62x + 29y + 19z - 105 = 0.$$

26. (a) : (P) Here, $2y^2 = ax^2 + b$

$$\Rightarrow 4y \frac{dy}{dx} = 2ax \Rightarrow \left[\frac{dy}{dx} \right]_{(1,-1)} = \frac{a}{-2} = -1 \quad [\text{Given}]$$

$$\Rightarrow a = 2$$

$$\text{Also, } (1, -1) \text{ lies on } 2y^2 = ax^2 + b$$

$$\Rightarrow 2 = a + b \Rightarrow b = 0 \Rightarrow a - b = 2$$

(Q) ∴ (a, b) lies on the curve $9y^2 = x^3$

$$\therefore 9b^2 = a^3 \quad \dots(i)$$

$$\text{Now, } 18y \frac{dy}{dx} = 3x^2 \text{ and } \frac{dy}{dx} = \frac{x^2}{6y} \Rightarrow \left[\frac{dy}{dx} \right]_{(a,b)} = \frac{a^2}{6b}$$

Since, the normal to the curve at (a, b) makes equal intercepts with the coordinate axes. Therefore, slope of normal = ± 1 .

$$\Rightarrow -\frac{1}{\left[\frac{dy}{dx} \right]_{(a,b)}} = \pm 1 \Rightarrow \left[\frac{dy}{dx} \right]_{(a,b)} = \pm 1 \Rightarrow \frac{a^2}{6b} = \pm 1$$

$$\Rightarrow a^2 = \pm 6b \Rightarrow a^4 = 36b^2 = 36 \left(\frac{a^3}{9} \right) \quad [\text{From (i)}]$$

$$\therefore a = 0, 4 \Rightarrow b = 0, \pm 8/3.$$

But the line making equal intercepts with the coordinate axes can not pass through the origin.

Hence, the required points are $(4, 8/3)$ and $(4, -8/3)$

$$\therefore a - b = 4/3 \text{ and } a - b = 20/3 \text{ and } a + |b| = 4 + 8/3 = 20/3$$

$$\text{(R) We have, } y = ax^2 + bx + 7/2 \Rightarrow \frac{dy}{dx} = 2ax + b$$

$$\therefore (1, 2) \text{ lies on } y = ax^2 + bx + 7/2$$

$$\therefore 2 = a + b + 7/2 \Rightarrow a + b + 3/2 = 0 \quad \dots(ii)$$

$$\left[\frac{dy}{dx} \right]_{(1,2)} = 2a(1) + b = 2a + b \quad \dots(iii)$$

$$\text{Also, we have } y = x^2 + 6x + 10 \Rightarrow \frac{dy}{dx} = 2x + 6$$

$$\therefore -\frac{1}{\left[\frac{dy}{dx} \right]_{(-2,2)}} = -\frac{1}{2(-2)+6} = -\frac{1}{2} \quad \dots(iii)$$

$$\text{Given, } 2a + b = -1/2 \quad \dots(iv)$$

Solving (i) and (iv), we get $a = 1, b = -5/2$

$$\Rightarrow a - b = 7/2 \text{ and } 5a + 2b = 0$$

27. (d) : Let n be the number of times a fair coin is tossed and X be the number of times head occurs.

$$\therefore p = \frac{1}{2} = q, \text{ Now, } P(7) = P(9)$$

$$\Rightarrow {}^nC_7 \left(\frac{1}{2} \right)^7 \left(\frac{1}{2} \right)^{n-7} = {}^nC_9 \left(\frac{1}{2} \right)^9 \left(\frac{1}{2} \right)^{n-9}$$

$$\Rightarrow {}^nC_7 = {}^nC_9 \Rightarrow n = 16$$

$$\therefore P(2) = {}^{16}C_2 \left(\frac{1}{2} \right)^2 \left(\frac{1}{2} \right)^{14} = \frac{15}{2^{13}}$$

∴ Statement-I is false but Statement-II is true.

$$\text{28. (c) : Putting } 4x + y + 1 = t \Rightarrow 4 + \frac{dy}{dx} = \frac{dt}{dx}$$

$$\Rightarrow \frac{dy}{dx} = \frac{dt}{dx} - 4$$

$$\text{Given equation becomes } \frac{dt}{dx} - 4 = t^2 \Rightarrow \frac{dt}{t^2 + 4} = dx$$

$$\text{Integrating both sides, } \int \frac{dt}{4 + t^2} = \int dx$$

$$\Rightarrow \frac{1}{2} \tan^{-1} \frac{t}{2} = x + c \Rightarrow \frac{1}{2} \tan^{-1} \left(\frac{4x + y + 1}{2} \right) = x + c$$

29. (a) : We have, $y = x^2 - 4x + 3$


$$\therefore \text{ Required area } = \int_1^3 -(x^2 - 4x + 3) dx$$

$$= - \left[\frac{x^3}{3} - \frac{4x^2}{2} + 3x \right]_1^3$$

$$= \frac{4}{3} \text{ sq. units}$$

∴ Statement-I is true.

Obviously, Statement-II is true.



30. (a)

31. (d): We have, $(\vec{a} + 3\vec{b}) \times (3\vec{a} - \vec{b})$
 $= 3(\vec{a} \times \vec{a}) - (\vec{a} \times \vec{b}) + 9(\vec{b} \times \vec{a}) - 3(\vec{b} \times \vec{b})$
 $= -(\vec{a} \times \vec{b}) - 9(\vec{a} \times \vec{b}) \quad [\because \vec{a} \times \vec{a} = 0 \text{ and } -(\vec{a} \times \vec{b}) = \vec{b} \times \vec{a}]$
 $= -10(\vec{a} \times \vec{b})$
 $\therefore [(\vec{a} + 3\vec{b}) \times (3\vec{a} - \vec{b})]^2 = 100(\vec{a} \times \vec{b})^2$

Now, $|\vec{a} \times \vec{b}| = |\vec{a}||\vec{b}|\sin 60^\circ = 1 \times 2 \times \frac{\sqrt{3}}{2} = \sqrt{3}$
 $\therefore 100(\vec{a} \times \vec{b})^2 = 100 \times (\sqrt{3})^2 = 300$

32. (c)

33. (c): Put $I = \int \frac{dx}{\sqrt{x+3}\sqrt{x}}$;

Now, put $x = t^6 \Rightarrow dx = 6t^5 dt$

$\Rightarrow I = \int \frac{6t^5}{t^3 + t^2} dt = 6 \int \frac{t^3}{t+1} dt = 6 \int \left(t^2 - t + 1 - \frac{1}{t+1} \right) dt$
 $= 2t^3 - 3t^2 + 6t - 6 \log(t+1) + C$
 $= 2\sqrt{x} - 3(\sqrt[3]{x}) + 6(\sqrt[6]{x}) - 6 \log(\sqrt[6]{x} + 1) + C$

34. (b): Let P and Q work for x and y days respectively.

$\therefore$ Total labour cost = $350x + 450y$

	P	Q	Minimum requirements
Shirts	6	7	51
Trousers	3	3	24

Constraints are $6x + 7y \geq 51$, $3x + 3y \geq 24$, $x \geq 0$, $y \geq 0$

$\therefore$ Given problem can be formulated as

Minimize $Z = 350x + 450y$

Subject to, $6x + 7y \geq 51$, $3x + 3y \geq 24$, $x \geq 0$, $y \geq 0$

35. (d)

36. (c): Let S be the sum of volume of parallelopiped and sphere, then

$S = x(2x) \left(\frac{x}{3} \right) + \frac{4}{3} \pi r^3 = \frac{2x^3}{3} + \frac{4}{3} \pi r^3$

37. (a): Since, sum of surface area of box and sphere is given to be constant k^2 .

$\therefore 2 \left(x \times 2x + 2x \times \frac{x}{3} + \frac{x}{3} \times x \right) + 4\pi r^2 = k^2$

$\Rightarrow 6x^2 + 4\pi r^2 = k^2$

$\Rightarrow x^2 = \frac{k^2 - 4\pi r^2}{6} \Rightarrow x = \sqrt{\frac{k^2 - 4\pi r^2}{6}}$

38. (b)

39. (d): Since, $x^2 = \frac{k^2 - 4\pi r^2}{6} = \frac{1}{6} \left[k^2 - 4\pi \left(\frac{k^2}{54 + 4\pi} \right) \right]$

$= \frac{9k^2}{54 + 4\pi} = 9 \left(\frac{k^2}{54 + 4\pi} \right) = 9r^2 = (3r)^2$
 $\Rightarrow x = 3r$

40. (c): Minimum value of S is given by

$\frac{2}{3}(3r)^3 + \frac{4}{3}\pi r^3 = 18r^3 + \frac{4}{3}\pi r^3 = \left(18 + \frac{4}{3}\pi \right) r^3$
 $= \left(18 + \frac{4}{3}\pi \right) \left(\frac{k^2}{54 + 4\pi} \right)^{3/2} = \frac{1}{3} \frac{k^3}{(54 + 4\pi)^{1/2}}$

(41-45): Let E_1, E_2, E_3 be the events of drawing a bolt produced by machine A, B and C, respectively.

Then, $P(E_1) = \frac{30}{100} = \frac{3}{10}$, $P(E_2) = \frac{20}{100} = \frac{1}{5}$

and $P(E_3) = \frac{50}{100} = \frac{1}{2}$

Also, let E be the event of drawing a defective bolt.

Then, $P(E|E_1) = \frac{5}{100} = \frac{1}{20}$, $P(E|E_2) = \frac{2}{100} = \frac{1}{50}$,

$P(E|E_3) = \frac{4}{100} = \frac{1}{25}$

41. (b)

42. (b): Probability that the defective bolt is manufactured by machine B = $P(E_2|E)$

$$= \frac{P(E|E_2) \cdot P(E_2)}{P(E|E_1) \cdot P(E_1) + P(E|E_2) \cdot P(E_2) + P(E|E_3) \cdot P(E_3)}$$

$$= \frac{\frac{1}{50} \times \frac{1}{5}}{\left(\frac{1}{20} \times \frac{3}{10} \right) + \left(\frac{1}{50} \times \frac{1}{5} \right) + \left(\frac{1}{25} \times \frac{1}{2} \right)} = \frac{1}{250} \times \frac{1000}{39} = \frac{4}{39} \approx 0.1$$

43. (c)

44. (a): Probability that the defective bolt is not manufactured by machine B i.e., it is manufactured by machine A or C = $\frac{15}{39} + \frac{20}{39} = \frac{35}{39}$.

45. (c): Required probability = $\frac{15}{39} + \frac{4}{39} = \frac{19}{39} \approx 0.5$.

SECTION B2 (APPLIED MATHEMATICS)

16. (b): $f(x) = 2x^3 + 3x^2 - 12x + 5$; $f'(x) = 6x^2 + 6x - 12$
 $= 6(x^2 + x - 2) = 6(x+2)(x-1)$

For critical point $f'(x) = 0$

$\Rightarrow 6(x+2)(x-1) = 0$

$\Rightarrow x = -2 \text{ and } x = 1$

Now, check the value of function $f(x)$ for different values of x, we get

Function $f'(x)$ is negative between $x = -2$ and 1

$\therefore f(x)$ is increasing in $(-\infty, -2] \cup [1, \infty)$

17. (b)

18. (a)

19. (c): When index number is 110 then cost of rice is ₹ 16.50 per kg.

So, when index number is 1, then cost of rice will be

$$₹ \frac{16.50}{110} \text{ per kg}$$

When index number is 100, then cost of rice will be

$$₹ \frac{16.50}{110} \times 100 \text{ per kg i.e., 15 per kg}$$

i.e., cost of rice in 2018 will be ₹ 15 per kg

[∵ We assume that index number of base year is 100]

20. (d): Average Cost (AC) = $\frac{C(x)}{x}$

Marginal Cost = $\frac{d}{dx} C(x)$

Marginal Average Cost = $\frac{1}{x} (MC - AC)$

Marginal Revenue = $\frac{d}{dx} R(x)$

21. (a): $A = \begin{bmatrix} a & 5 & c \\ -5 & b & 0 \\ d & 0 & 0 \end{bmatrix}$ is skew symmetric

∴ Diagonal element must be zero $\Rightarrow a = b = 0$

Also $a_{ij} = -a_{ji}$ for $1 \leq i \leq 3$ and $1 \leq j \leq 3. \Rightarrow a_{31} = -a_{13}$
 $\Rightarrow d = -c$

Now, $2a + b + c + d = c + d$ [$\because a = b = 0$]
 $= c - c$ [$\because d = -c$]
 $= 0$

22. (d): The value of function $F = 3x + 5y$ is minimum at (0, 3) and (5, 0)

So, every point on the line segment joining (0, 3) and (5, 0) will give minimum value.

23. (b): $E(x) = \sum x_i p(x_i)$
 $= 20 \times \frac{1}{10} + 15 \times \frac{1}{5} + 10 \times \frac{1}{2} + 5 \times \frac{1}{5} = 2 + 3 + 5 + 1 = 11$

24. (a): $3^2 = 9 = 1 \pmod{4}$
 $3^{16} = (3^2)^8 = 1^8 \pmod{4} \Rightarrow 3^{16} \equiv 1 \pmod{4}$
 $\therefore 39 \pmod{4} \equiv 1$

25. (b): Here,

The C.P. of cheaper oranges 'c' = ₹30 per kg,

The C.P. of dearer oranges 'd' = ₹45 per kg

and the mean price of the mixture 'm' = ₹40 per kg

So, $\frac{\text{quantity of cheaper oranges}}{\text{quantity of dearer oranges}} = \frac{d - m}{m - c}$

$$= \frac{45 - 40}{40 - 30} = \frac{5}{10} = \frac{1}{2}$$

Hence, the required ratio is 1 : 2.



PUBLIC NOTICE

CUET 2023

National Testing Agency has been mandated by the **Ministry of Education and UGC** to conduct Common University Entrance Test (CUET) for admission into **Undergraduate Programmes in Central Universities under Ministry of Education, and other participating Universities / Institutions / Organizations / Autonomous Colleges.**

CUET (UG) will provide a **single window opportunity** to students seeking admission in any of the **Central Universities (CUs)** or other participating organizations (including **State Universities, Deemed and Private Universities**) across the Country.

For the Academic Session **2023-24**, **CUET (UG) – 2023** will be conducted in the **Computer Based Test (CBT) mode** as per the details given below :

Online Submission of Application Form	09 February to 12 March 2023 (Up to 09:00 P.M.)
Last date of successful transaction of fee through Credit / Debit Card / Net-Banking / UPI	12 March 2023 (up to 11:50 P.M.)
Correction in Particulars	15 March to 18 March 2023 (Up to 11:50 P.M.)
Announcement of the City of Examination	30 April 2023
Downloading of Admit Cards from the NTA website	Second week of the May 2023
Date of Examination	21 May 2023 onwards
Display of Recorded Responses and Answer Keys	To be announced later on the website
Website(s)	www.nta.ac.in , https://cuetsamarth.ac.in/
Declaration of Result on the NTA website	To be announced later on the website

The details of the programmes offered by Central Universities (CUs) and other participating Universities / Organizations are available on their respective websites as well as on <https://cuetsamarth.ac.in/> **Candidates must carefully refer to the eligibility requirements of the programmes offered by various Central Universities and participating Universities / Organizations in this regard.**

The CUET (UG) - 2023 will be conducted in **13 languages** i.e. **English, Hindi, Assamese, Bengali, Gujarati, Kannada, Malayalam, Marathi, Odia, Punjabi, Tamil, Telugu, and Urdu.**

For CUET (UG) - 2023, the following is the scheme of examination :

Section	Subjects/ Tests	Questions to be Attempted
Section 1A - Languages	There are 13 different languages. Any of these languages may be chosen.	40 questions to be attempted
Section 1B - Languages	There are 20 Languages. Any of these languages may be chosen.	out of 50 in each language
Section 2 - Domain Specific Subject	There are 27 Domain specific subjects being offered under this section. A candidate may choose any subject as desired by the applicable University/ Organization.	35/40 Questions to be attempted out of 45/50
Section 3 - General Test	For any such undergraduate programme / programmes being offered by Universities where a General Test is being used for admission.	50 Questions to be attempted out of 60

Note:

- From the above subjects / languages, the **candidate can choose maximum of 10 subjects from all three Section.**
- Examination will be conducted on multiple days in three shifts, depending on the number of Candidates and Subject choices.**

In case any candidate faces difficulty in applying for **CUET (UG) – 2023**, he/she may contact **011 - 40759000 / 011 - 69227700** or e-mail at cuetsamarth@nta.ac.in. The candidates are also advised to be in touch with the NTA website(s) www.nta.ac.in and <https://cuetsamarth.ac.in/> for the latest updates regarding the exam.

26. (c) : Given $3(2-x) \geq 2(1-x) \Rightarrow 6-3x \geq 2-2x$
 $\Rightarrow 6-3x+2x \geq 2 \Rightarrow 6-x \geq 2 \Rightarrow -x \geq 2-6$
 $\Rightarrow -x \geq -4 \Rightarrow x \leq 4$

27. (b) : Both statements are incorrect as the measurable characteristic of population is called parameter and measurable characteristic of sample is called statistic.

28. (d) : Fisher's ideal index number
 $= \sqrt{\text{Laspeyre's Index} \times \text{Paache's Index}} = \sqrt{196 \times 64}$
 $= 14 \times 8 = 112$

29. (d) : Variance is not a time series component, it refers to the spread of a data set.

30. (c) : $f(x) = x^3 - 3x^2 + 3x + 4$; $f'(x) = 3x^2 - 6x + 3$.
 For critical point, put $f'(x) = 0$ i.e., $3x^2 - 6x + 3 = 0$
 $\Rightarrow 3(x^2 - 2x + 1) = 0 \Rightarrow (x-1)^2 = 0$
 $\Rightarrow x = 1$ is the critical point

31. (a)

32. (b) : Let the total profit be ₹x.

$\therefore$ 10% of profit goes in charity,
 $\therefore$ Profit shares by both the partner = 90% of
 $x = \frac{90x}{100} = \frac{9x}{10}$.

Now, Rakesh share of profit = $\frac{4}{9} \times \frac{9x}{10} \Rightarrow 18000 = \frac{2}{5}x$
 $\Rightarrow x = 45000$. Hence, the total profit is ₹45000.

33. (c) : $C(x) = 9x - 6x^2 + 5x^3$

Average cost (AC) = $\frac{\text{Total Cost}}{\text{Output}} = \frac{C(x)}{x} = 9 - 6x + 5x^2$

Marginal average cost = $\frac{d}{dx}(AC) = -6 + 10x$.

34. (a)

35. (d) : Only I and V are true statement.

36. (c) : Data estimation is not a type of statistical inference.

37. (b)

38. (a) : Given original cost of dining table = ₹36000
 Useful life = 15 years, Annual depreciation = ₹2000
 Let the scarp value of the dining table be ₹S, the using
 annual depreciation = $\frac{\text{original cost} - \text{scrap value}}{\text{useful life}}$
 we get $2000 = \frac{36000 - S}{15} \Rightarrow 30000 = 36000 - S \Rightarrow S = 6000$
 Hence, the scarp value of dining table is ₹6000.

39. (c) : $\frac{dx}{dt} = 1 - 4t^3$ and $\frac{dy}{dt} = 2t - 3t^2$

$\therefore \frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{2t - 3t^2}{1 - 4t^3}$

Now, at $x = 0, y = 0$ we get, $t - t^4 = 0$ and $t^2 - t^3 = 0$
 $t(1 - t^3) = 0$ and $t^2(1 - t) = 0 \Rightarrow t = 0$ and 1 is common value.

$\therefore \frac{dy}{dx} \Big|_{t=0} = 0$ and $\frac{dy}{dx} \Big|_{t=1} = \frac{2-3}{1-4} = \frac{-1}{-3} = \frac{1}{3}$

$\therefore \frac{dy}{dx}$ at (0, 0) is 0 or $\frac{1}{3}$.

40. (a)

41. (b) : We construct the following table :

x_i	P_i	$P_i x_i$	$P_i x_i^2$
2	0.2	0.4	0.8
3	0.5	1.5	4.5
4	0.3	1.2	4.8
Total		3.1	10.1

$\therefore$ Mean = $\mu = 3.1$

Variance = $\sum P_i x_i^2 - \mu^2 = 10.1 - (3.1)^2 = 10.1 - 9.61 = 0.49$

Standard deviation = $\sqrt{0.49} = 0.7$

42. (c) : We find $(672 + 541 + 383 + 295 + 101 + 86) \bmod 3$
 Since $672 \equiv 0 \pmod{3}$, $541 \equiv 1 \pmod{3}$, $383 \equiv 2 \pmod{3}$,
 $295 \equiv 1 \pmod{3}$, $101 \equiv 2 \pmod{3}$ and $86 \equiv 2 \pmod{3}$.
 So, $(672 + 541 + 383 + 295 + 101 + 86) \bmod 3$
 $\equiv (0 + 1 + 2 + 1 + 2 + 2) \bmod 3 \equiv 8 \pmod{3} \equiv 2 \pmod{3}$.
 Hence, the remainder when $(672 + 541 + 383 + 295 + 101 + 86)$ is divided by 3 is 2.

43. (a) : Given, upstream speed $v = 10$ km/h
 and downstream speed $u = 18$ km/h
 then speed of boat in still water

$x = \frac{u+v}{2} = \frac{18+10}{2} = 14$ km/h and speed of stream

$y = \frac{u-v}{2}$ and $\frac{18-10}{2} = 4$ km/h.

44. (c) : Cost of house = ₹ 5200000

Down payment = ₹ 700000 $\therefore$ Balance = ₹ 4500000

So, $P = ₹ 4500000$, $i = \frac{9}{12 \times 100} = 0.0075$

and $n = 25 \times 12 = 300$

$\therefore \text{EMI} = \frac{4500000 \times 0.0075 \times (1.0075)^{300}}{(1.0075)^{300} - 1}$
 $= \frac{33750 \times 9.4084}{9.4084} = \frac{33750 \times 9.4084}{8.4084} = 37763.84$

Hence, EMI = ₹ 37764

45. (d) : All of the given tests are the test of adequacy of index numbers.




WB JEE

PRACTICE PAPER 2023

Category-I (Q.1 to 50)

(Carry 1 mark each. Only one option is correct. Negative marks : $\frac{1}{4}$)

- For real numbers x and y , we write $xRy \Leftrightarrow x - y + \sqrt{2}$ is an irrational number. Then, the relation R is
(a) Reflexive (b) Symmetric (c) Transitive (d) None of these
- 28 is divided into 4 parts which are in A.P. The ratio of the product of the first and third to the product of the second and fourth is 8 : 15. The largest part is
(a) 6 (b) 8 (c) 10 (d) 12
- The summation of two unit vectors is a third unit vector, then the modulus of the difference of these unit vectors is
(a) $\sqrt{3}$ (b) $1 - \sqrt{3}$ (c) $1 + \sqrt{3}$ (d) $-\sqrt{3}$
- If $1 - i$ is a root of the equation $x^2 + ax + b = 0$, where $a, b \in R$, then find the values of a and b .
(a) 2, 2 (b) -2, 2 (c) -2, -2 (d) 1, 2
- Two cards are drawn at random one by one without replacement from a pack of 52 playing cards. Find the probability that both the cards are black.
(a) $\frac{21}{104}$ (b) $\frac{25}{102}$ (c) $\frac{23}{102}$ (d) $\frac{25}{104}$
- The number of solutions of $\cos 2\theta = \sin \theta$ in $(0, 2\pi)$ is
(a) 1 (b) 2 (c) 3 (d) 4
- If $\log_{10} 2 = 0.30103$, then $\log_{10} 50 =$
(a) 2.30103 (b) 2.69897 (c) 1.69897 (d) 0.69897
- If $(4, 0)$ is a point on the circle $x^2 + ax + y^2 = 0$, then the centre of the circle is at
(a) $(-2, 0)$ (b) $(0, 2)$ (c) $(2, 0)$ (d) $(1, 0)$
- The integrating factor of the differential equation $(x + 3y^2) \frac{dy}{dx} = y$ is
(a) y (b) $-y$ (c) $1/y$ (d) $-1/y$
- How many numbers lying between 100 and 1000 can be formed with the digits 0, 1, 2, 3, 4, 5, if the repetition of the digits is not allowed?
(a) 90 (b) 50 (c) 40 (d) 100
- If the set A contains 5 elements, then the number of elements in the power set $P(A)$ is
(a) 32 (b) 25 (c) 16 (d) 8
- The focus of a parabolic mirror as shown in figure is at a distance of 5 cm from its vertex. If the mirror is 45 cm deep, find the distance AB .
(a) 30 cm (b) 45 cm (c) 60 cm (d) 90 cm
- If $\vec{a} = \hat{i} + \hat{j} + \hat{k}, \vec{b} = 2\hat{i} - 4\hat{k}, \vec{c} = \hat{i} + \lambda\hat{j} + 3\hat{k}$ are coplanar, then the value of λ is
(a) $5/2$ (b) $3/5$ (c) $7/3$ (d) None of these
- If A, B, C, D be angles of a cyclic quadrilateral taken in order, then $\cos A + \cos B + \cos C + \cos D =$
(a) 0 (b) 1 (c) -1 (d) $1/2$
- Find the value of $\tan(\alpha + \beta)$, given that $\cot \alpha = \frac{1}{2}, \alpha \in \left(\pi, \frac{3\pi}{2}\right)$ and $\sec \beta = \frac{-5}{3}, \beta \in \left(\frac{\pi}{2}, \pi\right)$.
(a) $1/11$ (b) $2/11$ (c) $5/11$ (d) $3/11$
- The sum to 20 terms of the series $1 \times 3^2 + 2 \times 5^2 + 3 \times 7^2 + \dots$, is
(a) 18800 (b) 188010 (c) 188020 (d) 188090
- If α and β are the roots of the equation $x^2 + 2x + 4 = 0$, then $\frac{1}{\alpha^3} + \frac{1}{\beta^3}$ is equal to
(a) $-1/2$ (b) $1/2$ (c) 32 (d) $1/4$



18. In a $\triangle ABC$, a , c and A are given and b_1 , b_2 are two values of the third side b such that $b_2 = 2b_1$. Then $\sin A =$

- (a) $\sqrt{\frac{9c^2 - a^2}{8a^2}}$ (b) $\sqrt{\frac{9a^2 - c^2}{8c^2}}$
 (c) $\sqrt{\frac{8a^2 - 9c^2}{8a^2}}$ (d) none of these

19. The differential equation satisfied by $y = \frac{A}{x} + B$ is (A, B are parameters)

- (a) $x^2 y_1 = y$ (b) $xy_1 + 2y_2 = 0$
 (c) $xy_2 + 2y_1 = 0$ (d) none of these

20. Find a if the 17th and 18th terms of the expansion $(2 + a)^{50}$ are equal.

- (a) 1 (b) 2 (c) 3 (d) 4

21. $\int_{-\pi/4}^{\pi/4} \frac{dx}{1 + \cos 2x}$ is equal to

- (a) 1 (b) 2 (c) 3 (d) 4

22. If $x^m y^n = (x + y)^{m+n}$, then $\frac{dy}{dx}$ is equal to

- (a) $\frac{x+y}{xy}$ (b) xy (c) $\frac{x}{y}$ (d) $\frac{y}{x}$

23. The function $f(x) = xe^{1-x}$

- (a) strictly increases in the interval $(1/2, 2)$
 (b) increases in the interval $(0, \infty)$
 (c) decreases in the interval $(0, 2)$
 (d) strictly decreases in the interval $(1, \infty)$

24. There are 7 greeting cards, each of a different colour and 7 envelopes of same 7 colours as that of the cards. The number of ways in which the cards can be put in envelopes, so that exactly 4 of the cards go into envelopes of respective colour, is

- (a) ${}^7C_3$ (b) $2 \cdot {}^7C_3$
 (c) $3! \cdot {}^4C_4$ (d) $3! \cdot {}^7C_3 \cdot {}^4C_3$

25. If $\Delta = \begin{vmatrix} a+1 & a+2 & a+4 \\ a+3 & a+5 & a+8 \\ a+7 & a+10 & a+14 \end{vmatrix}$, then value of Δ is

- (a) 2 (b) -2 (c) 4 (d) -4

26. The equation of circle of radius $\sqrt{17}$ units, with centre on the positive side of x -axis and through the point $(0, 1)$ is

- (a) $x^2 + y^2 - 8x - 1 = 0$
 (b) $x^2 + y^2 + 8x - 1 = 0$
 (c) $x^2 + y^2 - 9y + 1 = 0$
 (d) $2x^2 + 2y^2 - 3x + 2y = 4$

27. Let $g(x) = x^2 - 4x - 5$, then

- (a) g is one-one on R (b) g is not one-one on R
 (c) g is bijective on R (d) None of these

28. Let the eccentricity of the hyperbola $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ be reciprocal to that of the ellipse $x^2 + 9y^2 = 9$, then the ratio $a^2 : b^2$ equals

- (a) 8 : 1 (b) 1 : 8 (c) 9 : 1 (d) 1 : 9

29. $\sin\left[\frac{\pi}{3} - \sin^{-1}\left(-\frac{1}{2}\right)\right]$ is equal to

- (a) 1/2 (b) 1/3 (c) -1 (d) 1

30. The area between the curve $y = 4 + 3x - x^2$ and x -axis is

- (a) $\frac{125}{6}$ sq. units (b) $\frac{125}{3}$ sq. units
 (c) $\frac{125}{2}$ sq. units (d) none of these

31. If $(\vec{a} \times \vec{b})^2 + (\vec{a} \cdot \vec{b})^2 = 676$ and $|\vec{b}| = 2$, then $|\vec{a}|$ is equal to

- (a) 13 (b) 26
 (c) 39 (d) None of these

32. For $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$, then $14A^{-1}$ is given by

- (a) $14 \begin{bmatrix} 2 & -1 \\ 1 & 3 \end{bmatrix}$ (b) $\begin{bmatrix} 4 & -2 \\ 2 & 6 \end{bmatrix}$
 (c) $2 \begin{bmatrix} 2 & -1 \\ 1 & -3 \end{bmatrix}$ (d) $2 \begin{bmatrix} -3 & -1 \\ 1 & -2 \end{bmatrix}$

33. If A and B are two independent events, then the probability of occurrence of atleast one of A and B is given by

- (a) $1 - P(A)P(B)$ (b) $1 - P(A)P(B')$
 (c) $1 - P(A')P(B')$ (d) $1 - P(A')P(B)$

34. If $\frac{z_2}{z_1}$ is pure imaginary, then $\left| \frac{6z_1 - 8z_2}{4z_2 + 3z_1} \right| =$

- (a) 1 (b) $\sqrt{2}$ (c) 2 (d) 4

35. If in a triangle ABC , $\cos A \cos B + \sin A \sin B \sin C = 1$, then $a : b : c =$

- (a) $1 : 1 : \sqrt{2}$ (b) $1 : \sqrt{3} : 1$
 (c) $1 : \sqrt{2} : 1$ (d) $\sqrt{2} : \sqrt{3} : 1$

36. The value of $\int_0^\infty \frac{dx}{(x^2 + 4)(x^2 + 9)}$ is

- (a) $\pi/60$ (b) $\pi/20$ (c) $\pi/40$ (d) $\pi/80$

37. $\lim_{x \rightarrow 0} \frac{\pi^x - 1}{\sqrt{1+x} - 1}$

- (a) does not exist (b) equals $\log_e(\pi^2)$
(c) equals 1 (d) lies between 10 and 11

38. Find the 4th term from the end in the expansion of $\left(\frac{x^3}{2} - \frac{2}{x^2}\right)^9$.

- (a) $\frac{674}{x}$ (b) $\frac{670}{x^3}$ (c) $\frac{672}{x^3}$ (d) $\frac{-670}{x^3}$

39. Find the point of intersection of the lines

$$\frac{x+1}{-3} = \frac{y-3}{2} = \frac{z+2}{1} \text{ and } \frac{x}{1} = \frac{y-7}{-3} = \frac{z+7}{2}.$$

- (a) $(-2, 1, 3)$ (b) $(2, 1, -3)$
(c) $(2, 3, 1)$ (d) $(2, -1, 3)$

40. The mean and variance of n observations $x_1, x_2,$

$x_3, \dots, x_n$ are 5 and 0 respectively. If $\sum_{i=1}^n x_i^2 = 400$, then the value of n is equal to

- (a) 80 (b) 25 (c) 20 (d) 16

41. The solution of the differential equation

$$\frac{dy}{dx} = \frac{e^x + x^2}{e^y} \text{ is}$$

- (a) $e^x = \frac{y^3}{3} + e^y + c$ (b) $e^y = \frac{x^2}{3} + e^x + c$

- (c) $e^y = \frac{x^3}{3} + e^x + c$ (d) None of these

42. A straight line through the point $(3, -2)$ is inclined at an angle 60° to the line $\sqrt{3}x + y = 1$. If it intersects the x -axis, then its equation will be

- (a) $y + x\sqrt{3} + 2 + 3\sqrt{3} = 0$
(b) $y - x\sqrt{3} + 2 + 3\sqrt{3} = 0$
(c) $y - x\sqrt{3} - 2 - 2\sqrt{3} = 0$
(d) $y - x\sqrt{3} + 2 - 3\sqrt{3} = 0$

43. If x and ' a ' are real, then the value of ' a ' for which

$$x^2 - \frac{3ax}{2} + 1 - a^2 \text{ is positive is}$$

- (a) $-\frac{4}{25}p$ (b) $\frac{4}{25}$ (c) $|a| > \frac{4}{5}$ (d) $|a| < \frac{4}{5}$

44. If $P(n)$ is a statement such that $P(3)$ is true. Also $P(k)$ is true $\Rightarrow P(k+1)$ is true for all $k \geq 3$, then $P(n)$ is true

- (a) for all $n \in N$ (b) for all $n \geq 3, n \in N$
(c) for all $n > 4, n \in N$ (d) none of these

45. If A and B are non-empty sets, then

$(A \cap B) \cup (A - B)$ is equal to

- (a) B (b) A (c) A' (d) B'

46. The second order derivative of $a \sin^3 t$ with respect to $a \cos^3 t$ at $t = \pi/4$ is

- (a) 2 (b) $\frac{1}{12a}$ (c) $\frac{4\sqrt{2}}{3a}$ (d) $\frac{3a}{4\sqrt{2}}$

47. A function $f(x)$ is defined as follows for real x

$$f(x) = \begin{cases} 1-x^2 & , \text{ for } x < 1 \\ 0 & , \text{ for } x = 1 \\ 1+x^2 & , \text{ for } x > 1 \end{cases} \text{, then}$$

- (a) $f(x)$ is not continuous at $x = 1$
(b) $f(x)$ is continuous at $x = 1$
(c) $f(x)$ is continuous at all real numbers
(d) none of these

48. $\int \sin^{-1} x dx$ is equal to

- (a) $\cos^{-1} x + C$ (b) $x \sin^{-1} x + \sqrt{1-x^2} + C$
(c) $\frac{1}{\sqrt{1-x^2}} + C$ (d) $x \sin^{-1} x - \sqrt{1-x^2} + C$

49. If e_1 is the eccentricity of the conic $9x^2 + 4y^2 = 36$ and e_2 is the eccentricity of conic $9x^2 - 4y^2 = 36$, then which of the following is true?

- (a) $e_1^2 + e_2^2 = 2$ (b) $3 < e_1^2 + e_2^2 < 4$
(c) $e_1^2 + e_2^2 > 4$ (d) None of these

50. The determinant $\begin{vmatrix} y+k & y & y \\ y & y+k & y \\ y & y & y+k \end{vmatrix}$ is equal to

- (a) $k(3y + k^2)$ (b) $3y + k^3$
(c) $3y + k^2$ (d) $k^2(3y + k)$

Category-II (Q.51 to 65)

(Carry 2 marks each. Only one option is correct. Negative marks : $\frac{1}{2}$)

51. A square piece of tin of side 18 cm is to be made into a box without top by cutting a square from each corner and folding up the flaps to form a box. What should be the side of the square to be cut off so that the volume of the box is maximum? Also, find this maximum volume.

- (a) 2 cm, 430 cm³ (b) 5 cm, 425 cm³
(c) 3 cm, 432 cm³ (d) 3 cm, 423 cm³

52. If $\tan^{-1}\left(\frac{x+1}{x-1}\right) + \tan^{-1}\left(\frac{x-1}{x}\right) = \tan^{-1}(-7)$, then

the value of x is

- (a) 0 (b) -2 (c) 1 (d) 2

53. Find the standard deviation for the following data :

x_i	3	8	13	18	23
f_i	7	10	15	10	6

- (a) 6.12 (b) 5.12 (c) 3.12 (d) 7.12

54. Evaluate : $\int_0^{\pi/2} \frac{1}{2 \cos x + 4 \sin x} dx$

- (a) $\sqrt{5} \log \left(\frac{3+\sqrt{5}}{2} \right)$ (b) $\frac{1}{\sqrt{5}} \log \left(\frac{3-\sqrt{5}}{5} \right)$
(c) $\frac{1}{\sqrt{5}} \log \left(\frac{3+\sqrt{5}}{2} \right)$ (d) None of these

55. If $y^{y^{y^{\dots \infty}}} = \log_e(x + \log_e(x + \dots))$, then $\frac{dy}{dx}$ at $(x = e^2 - 2, y = \sqrt{2})$ is

- (a) $\frac{\log \left(\frac{e}{2} \right)}{2\sqrt{2}(e^2 - 1)}$ (b) $\frac{\log 2}{2\sqrt{2}(e^2 - 1)}$
(c) $\frac{\sqrt{2} \log \frac{e}{2}}{(e^2 - 1)}$ (d) None of these

56. Coloured balls are distributed in three bags as shown in the following table :

Bag	Colour of balls		
	Black	White	Red
I	1	2	3
II	2	4	1
III	4	5	3

A bag is selected at random and then two balls are drawn from the selected bag. They happen to be black and red. What is the probability that they have come from bag I?

- (a) $\frac{231}{551}$ (b) $\frac{232}{551}$ (c) $\frac{233}{550}$ (d) $\frac{231}{550}$

57. If ω is an imaginary cube root of unity and

$$\begin{vmatrix} x + \omega^2 & \omega & 1 \\ \omega & \omega^2 & 1 + x \\ 1 & x + \omega & \omega^2 \end{vmatrix} = 0, \text{ then one of the values of } x \text{ is}$$

- (a) 1 (b) 0 (c) -1 (d) 2

58. The vector equation of the plane passing through the point $(-1, 2, -5)$ and parallel to the vectors $4\hat{i} - \hat{j} + 3\hat{k}$ and $\hat{i} + \hat{j} - \hat{k}$ is

- (a) $\vec{r} \cdot (-2\hat{i} + 7\hat{j} + 5\hat{k}) = 13$
(b) $\vec{r} \cdot (-2\hat{i} + 7\hat{j} + 5\hat{k}) = -13$

(c) $\vec{r} \cdot (-2\hat{i} + 7\hat{j} + 5\hat{k}) = -9$

(d) $\vec{r} \cdot (-2\hat{i} + 7\hat{j} + 5\hat{k}) = 9$

59. If the tangent drawn at a point $(t^2, 2t)$ on the parabola $y^2 = 4x$ is same as normal drawn at $(\sqrt{5} \cos \alpha, 2 \sin \alpha)$ on the ellipse $\frac{x^2}{5} + \frac{y^2}{4} = 1$, then which of following is not true?

- (a) $t = \pm 1/\sqrt{5}$ (b) $\alpha = -\tan^{-1} 2$
(c) $\alpha = \tan^{-1} 2$ (d) None of these

60. If a, b, c be the $p^{\text{th}}, q^{\text{th}}$ and r^{th} terms respectively of an A.P. and G.P. both, then the product of the roots of equation $(a^b b^c c^a)x^2 - (abc)x + (a^c b^a c^b) = 0$ equals

- (a) -1 (b) 2 (c) abc (d) 1

61. Find the sum of the sequence 7, 77, 777, 7777, ... to n terms.

- (a) $\frac{7}{9} \left[\frac{10(10^n - 1)}{9} - n \right]$ (b) $\frac{7}{9} \left[\frac{10(10^n + 1)}{9} - n \right]$
(c) $\frac{10^n - 1}{9}$ (d) None of these

62. Let $L = \lim_{x \rightarrow 0} \frac{a - \sqrt{a^2 - x^2} - \frac{x^2}{4}}{x^4}$, $a > 0$. If L is finite, then

- (a) $a = 2$ (b) $a = 1$
(c) $a = 1/3$ (d) None of these

63. Find the smaller of the two areas enclosed by the curves $x^2 + y^2 = 4$ and $y^2 = 3(2x - 1)$.

- (a) $(4\pi - \sqrt{3})$ sq. units (b) $\frac{1}{2}(2\pi + \sqrt{3})$ sq. units
(c) $\frac{1}{3}(4\pi - \sqrt{3})$ sq. units (d) None of these

64. Let ρ_1 and ρ_2 be two equivalence relations defined on a non-void set S . Then

- (a) both $\rho_1 \cap \rho_2$ and $\rho_1 \cup \rho_2$ are equivalence relations.
(b) $\rho_1 \cap \rho_2$ is equivalence relation but $\rho_1 \cup \rho_2$ is not so.
(c) $\rho_1 \cup \rho_2$ is equivalence relation but $\rho_1 \cap \rho_2$ is not so.
(d) neither $\rho_1 \cap \rho_2$ nor $\rho_1 \cup \rho_2$ is equivalence relation.

MONTHLY TEST DRIVE CLASS XII ANSWER KEY

1. (d) 2. (a) 3. (a) 4. (d) 5. (b)
6. (b) 7. (b,c) 8. (a,b,d) 9. (a,b) 10. (a,b,c)
11. (a,b,c) 12. (a,c) 13. (a,b,c) 14. (a) 15. (b)
16. (d) 17. (5) 18. (1) 19. (3) 20. (2)

65. Evaluate : $\int \frac{x^2}{(x-1)^3(x+1)} dx$

- (a) $\frac{1}{8} \log \left| \frac{x-1}{x+1} \right| + \frac{3}{4(x-1)} + \frac{1}{4(x-1)^2} + C$
 (b) $\frac{1}{8} \log \left| \frac{x+1}{x-1} \right| - \frac{3}{4(x-1)} - \frac{1}{4(x+1)^2} + C$
 (c) $\frac{1}{8} \log \left| \frac{x-1}{x+1} \right| - \frac{3}{4(x-1)} - \frac{1}{4(x-1)^2} + C$
 (d) None of these

Category - III (Q.66 to 75)

(Carry 2 marks each. One or more options are correct. No negative marks)

66. If maximum and minimum values of the

determinant $\begin{vmatrix} 1 + \cos^2 x & \sin^2 x & \cos 2x \\ \cos^2 x & 1 + \sin^2 x & \cos 2x \\ \cos^2 x & \sin^2 x & 1 + \cos 2x \end{vmatrix}$ are

α and β , then

- (a) $\alpha^2 + \beta^{101} = 10$ (b) $\alpha^3 - \beta^{99} = 26$
 (c) $2\alpha^2 - 18\beta^{11} = 0$ (d) $\alpha^3 + 2\beta^2 = 0$

67. The internal bisector of $\angle A$ of triangle ABC meets side BC at D . A line drawn through D perpendicular to AD meets the side AC at E and the side AB at F . If a, b, c , are the sides of triangle ABC , then

- (a) AE is H.M. of b and c
 (b) $AD = \frac{2bc}{b+c} \cos \frac{A}{2}$ (c) $EF = \frac{4bc}{b+c} \sin \frac{A}{2}$
 (d) triangle AEF is isosceles

68. A and B are independent events. The probability that both A and B occur is $1/20$ and the probability that neither of them occurs is $3/5$. The probability of occurrence of A is

- (a) $1/2$ (b) $1/10$ (c) $1/4$ (d) $1/5$

69. If $\frac{\cos x - \cos \alpha}{\cos x - \cos \beta} = \frac{\sin^2 \alpha \cos \beta}{\sin^2 \beta \cos \alpha}$, then

- (a) $\cos x = \frac{\cos \alpha + \cos \beta}{1 + \cos \alpha \cos \beta}$
 (b) $\cos x = \frac{\cos \alpha + \cos \beta}{1 - \cos \alpha \cos \beta}$
 (c) $\tan \frac{x}{2} = \tan \frac{\alpha}{2} \tan \frac{\beta}{2}$
 (d) $\tan \frac{x}{2} = -\tan \frac{\alpha}{2} \tan \frac{\beta}{2}$

70. Equation of a tangent to the hyperbola $5x^2 - y^2 = 5$ and which passes through an external point $(2, 8)$ is

- (a) $3x - y + 2 = 0$ (b) $3x + y - 14 = 0$
 (c) $23x - 3y - 22 = 0$ (d) $3x - 23y + 178 = 0$

71. Given a real valued function f such that

$$f(x) = \begin{cases} \frac{\tan^2 x}{(x^2 - [x]^2)}, & \text{for } x > 0 \\ 1, & \text{for } x = 0 \\ \sqrt{\{x\} \cot \{x\}}, & \text{for } x < 0 \end{cases}$$

where $[x]$ is the integral part and $\{x\}$ is the fractional part of x , then

- (a) $\lim_{x \rightarrow 0^+} f(x) = 1$ (b) $\lim_{x \rightarrow 0^-} f(x) = \sqrt{\cot 1}$
 (c) $\cot^{-1} \left(\lim_{x \rightarrow 0^-} f(x) \right)^2 = 1$
 (d) f is continuous at $x = 0$

72. Let $f(x)$ be twice differentiable function such that $f''(x) > 0$ in $[0, 1]$. Then

- (a) $f(0) + f(1) = 2f(c)$, $0 < c < 1$
 (b) $f(0) + f(1) = 2f\left(\frac{1}{2}\right)$
 (c) $f(0) + f(1) > 2f\left(\frac{1}{2}\right)$
 (d) $f(0) + f(1) < 2f\left(\frac{1}{2}\right)$

73. Which of the following is/are the point/points that is/are at a distance of 12 units from the point whose position vector is $(8\hat{i} + 10\hat{j} - 8\hat{k})$ on the line which is parallel to $(2\hat{i} + \hat{j} + 2\hat{k})$?

- (a) $16\hat{i} + 14\hat{j}$ (b) $6\hat{j} - 16\hat{k}$
 (c) $16\hat{i} + 18\hat{j} - 4\hat{k}$ (d) None of these

74. Three vectors $\vec{a}, \vec{b}, \vec{c}$ are such that $\vec{a} \times \vec{b} = 4(\vec{a} \times \vec{c})$ and $|\vec{a}| = |\vec{b}| = 1$ and $|\vec{c}| = \frac{1}{4}$. If the angle between $\vec{b}$ and $\vec{c}$ is $\frac{\pi}{3}$, then $\vec{b}$ is

- (a) $\vec{a} + 4\vec{c}$ (b) $\vec{a} - 4\vec{c}$ (c) $4\vec{c} - \vec{a}$ (d) $2\vec{c} - \vec{a}$

75. Tangent is drawn at any point $P(x, y)$ on a curve, which passes through $(1, 1)$. The tangent cuts X -axis and Y -axis at A and B respectively. If $AP : BP = 3 : 1$, then

- (a) the differential equation of the curve is

$$3x \frac{dy}{dx} + y = 0$$

(b) the differential equation of the curve is

$$3x \frac{dy}{dx} - y = 0$$

(c) the curve passes through $\left(\frac{1}{8}, 2\right)$

(d) the normal at $(1, 1)$ is $x + 3y = 4$

SOLUTIONS

1. (a) : Reflexive : For any $x \in R$, we have $x - x + \sqrt{2} = \sqrt{2}$ is an irrational number $\Rightarrow xRx$ for all x . So, R is reflexive.

Symmetric : R is not symmetric, because $\sqrt{2}R1$ but $1 \not R \sqrt{2}$.

Transitive : R is not transitive also, because $\sqrt{2}R1$ and $1R2\sqrt{2}$ but $\sqrt{2} \not R 2\sqrt{2}$.

2. (c) : Since the sum is 28, we take the A.P. as $7 - 3d, 7 - d, 7 + d, 7 + 3d$.

It is given that
$$\frac{(7-3d)(7+d)}{(7-d)(7+3d)} = \frac{8}{15}$$

$$\Rightarrow 15[49 - 14d - 3d^2] = 8[49 + 14d - 3d^2]$$

$$\Rightarrow 3d^2 + 46d - 49 = 0 \Rightarrow d = 1$$

$\therefore$ The four parts are 4, 6, 8, 10.

3. (a) : Since, $\vec{a} + \vec{b} = \vec{c} \Rightarrow (\vec{a} + \vec{b})^2 = \vec{c}^2$

$$\Rightarrow |\vec{a}|^2 + |\vec{b}|^2 + 2|\vec{a}||\vec{b}|\cos\theta = |\vec{c}|^2$$

$$\Rightarrow 2(1 + \cos\theta) = 1 \Rightarrow \cos\theta = -\frac{1}{2}$$

Now, $|\vec{a} - \vec{b}|^2 = |\vec{a}|^2 + |\vec{b}|^2 - 2|\vec{a}||\vec{b}|\cos\theta$

$$= 1 + 1 - 2 \cdot \left(-\frac{1}{2}\right) = 3 \Rightarrow |\vec{a} - \vec{b}| = \sqrt{3}$$

4. (b) : Since complex roots always occur in conjugate pair.

$\therefore$ Other root is $1 + i$.

Sum of roots = $\frac{-a}{1} = (1 - i) + (1 + i) \Rightarrow a = -2$

Product of roots = $\frac{b}{1} = (1 - i)(1 + i) \Rightarrow b = 2$

5. (b) : Let E and F be the events that first and second card drawn are black, respectively.

Then, $P(E) = P(\text{black card in first draw}) = \frac{26}{52} = \frac{1}{2}$

When second card is drawn without replacement, the probability that the second card is black is the conditional probability of event F occurring when event E has already occurred.

So, $P(F|E) = \frac{25}{51}$

By multiplication rule of probability, we have

$$P(E \cap F) = P(E) \cdot P(F|E) = \frac{1}{2} \cdot \frac{25}{51} = \frac{25}{102}$$

6. (c) : $\cos 2\theta = \sin\theta \Rightarrow 1 - 2\sin^2\theta - \sin\theta = 0$

$$\Rightarrow 2\sin^2\theta + 2\sin\theta - \sin\theta - 1 = 0$$

$$\Rightarrow 2\sin\theta(\sin\theta + 1) - 1(\sin\theta + 1) = 0$$

$$\Rightarrow \sin\theta = -1, 1/2$$

In $(0, 2\pi)$; $\sin\theta = -1$ for $\theta = 3\pi/2$

and $\sin\theta = \frac{1}{2}$ for $\theta = \frac{\pi}{6}, \frac{5\pi}{6}$

$\therefore$ Total 3 solutions exist in $(0, 2\pi)$.

7. (c) : Consider $\log_{10} 50 = \log_{10} \frac{100}{2}$

$$= \log_{10} 10^2 - \log_{10} 2 = 2 \log_{10} 10 - \log_{10} 2 = 2 - (0.30103) = 1.69897$$

8. (c) : Since, $(4, 0)$ is a point on the circle $x^2 + ax + y^2 = 0$.

$$\therefore 16 + 4a + 0 = 0 \Rightarrow 4a = -16 \Rightarrow a = -4$$

$$\therefore \text{Equation of circle is } x^2 - 4x + y^2 = 0$$

or $x^2 - 4x + 4 - 4 + y^2 = 0 \Rightarrow (x - 2)^2 + (y - 0)^2 = 4$

$$\therefore \text{Centre } (2, 0) \text{ and radius} = 2$$

9. (c) : We have, $(x + 3y^2) \frac{dy}{dx} = y$

$$\Rightarrow \frac{x + 3y^2}{y} = \frac{dx}{dy} \Rightarrow \frac{dx}{dy} - \frac{x}{y} = 3y$$

This is a linear differential equation.

$$\therefore \text{I.F.} = e^{-\int \frac{dy}{y}} = e^{-\log y} = e^{\log y^{-1}} = \frac{1}{y}$$

10. (d) : Every number between 100 and 1000 is a 3-digit number. We first have to count the permutations of 6 digits taken 3 at a time i.e., ${}^6P_3$. But, these permutations will include those also where 0 is at the 100's place. For example, 092, 042, ..., etc are such numbers which are actually 2-digit numbers. To get the number of such numbers, we fix 0 at the 100's place and rearrange the remaining 5 digits taking 2 at a time. This will give ${}^5P_2$ numbers.

So, the required numbers = ${}^6P_3 - {}^5P_2 = \frac{6!}{3!} - \frac{5!}{2!} = 100$


11. (a) : $n(A) = 5 \Rightarrow n(P(A)) = 2^5 = 32$.

12. (c) : Since the distance from the focus to the vertex is 5 cm, we have, $a = 5$. Also, the axis of the mirror lies along the positive x -axis, the equation of the parabolic section is

$$y^2 = 4(5)x = 20x$$

When $x = 45$, $y^2 = 900$

Therefore, $y = \pm 30$. Hence, $AB = 2y = 2 \times 30 = 60$ cm



13. (d) : Since, $\vec{a} = \hat{i} + \hat{j} + \hat{k}$, $\vec{b} = 2\hat{i} - 4\hat{k}$, $\vec{c} = \hat{i} + \lambda\hat{j} + 3\hat{k}$ are coplanar.

$$\therefore [\vec{a} \vec{b} \vec{c}] = 0 \Rightarrow \begin{vmatrix} 1 & 1 & 1 \\ 2 & 0 & -4 \\ 1 & \lambda & 3 \end{vmatrix} = 0$$

$$\Rightarrow 4\lambda - 1(6 + 4) + 2\lambda = 0 \Rightarrow \lambda = 5/3$$

14. (a) : In a cyclic quadrilateral, the sum of opposite angles is 180° .

$$\therefore A + C = 180^\circ \text{ and } B + D = 180^\circ$$

$$\Rightarrow C = 180^\circ - A \text{ and } D = 180^\circ - B \quad \dots(i)$$

$$\begin{aligned} \text{Now, } \cos A + \cos B + \cos C + \cos D \\ = \cos A + \cos B + \cos(180^\circ - A) + \cos(180^\circ - B) \\ \quad \quad \quad [\text{From (i)}] \\ = \cos A + \cos B - \cos A - \cos B = 0 \end{aligned}$$

15. (b) : Given, $\cot \alpha = \frac{1}{2} \Rightarrow \tan \alpha = 2$ and $\sec \beta = \frac{-5}{3}$

$$\text{Then, } \tan \beta = \sqrt{\sec^2 \beta - 1} \Rightarrow \tan \beta = \pm \frac{4}{3}$$

$$\text{But, } \tan \beta = \frac{-4}{3} \quad (\because \tan \beta \text{ is -ve in II quadrant})$$

$$\therefore \tan(\alpha + \beta) = \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \cdot \tan \beta} = \frac{2 + \left(-\frac{4}{3}\right)}{1 - (2)\left(-\frac{4}{3}\right)} = \frac{2}{11}$$

16. (d) : The r^{th} term of the series $t_r = r(2r + 1)^2 = 4r^3 + 4r^2 + r$

$$\begin{aligned} \sum_{r=1}^{20} t_r &= 4 \sum_{r=1}^{20} r^3 + 4 \sum_{r=1}^{20} r^2 + \sum_{r=1}^{20} r \\ &= 4 \cdot \left(\frac{20 \times 21}{2}\right)^2 + \frac{4 \cdot 20 \cdot 21 \cdot 41}{6} + \frac{20 \cdot 21}{2} \\ &= 4(210)^2 + 40 \cdot 7 \cdot 41 + 210 = 188090 \end{aligned}$$

17. (d) : Given equation is $x^2 + 2x + 4 = 0$

Since α, β are roots of this equation.

$$\therefore \alpha + \beta = -2 \text{ and } \alpha\beta = 4$$

$$\begin{aligned} \text{Now, } \frac{1}{\alpha^3} + \frac{1}{\beta^3} &= \frac{\alpha^3 + \beta^3}{(\alpha\beta)^3} = \frac{(\alpha + \beta)(\alpha^2 + \beta^2 - \alpha\beta)}{(\alpha\beta)^3} \\ &= \frac{(-2)((\alpha + \beta)^2 - 3\alpha\beta)}{4 \times 4 \times 4} = \frac{-2(4 - 12)}{4 \times 4 \times 4} = \frac{(-2) \times (-8)}{4 \times 4 \times 4} = \frac{1}{4} \end{aligned}$$

18. (b) : $\cos A = \frac{b^2 + c^2 - a^2}{2bc}$

$$\Rightarrow b^2 - 2bc \cos A + (c^2 - a^2) = 0$$

As, b_1 and b_2 are two values of b .

$$\therefore b_1 + b_2 = 2c \cos A \text{ and } b_1 b_2 = c^2 - a^2$$

$$\Rightarrow 3b_1 = 2c \cos A \text{ and } 2b_1^2 = c^2 - a^2 \quad [\because b_2 = 2b_1 \text{ (given)}]$$

$$\therefore 2 \left(\frac{2c \cos A}{3} \right)^2 = c^2 - a^2 \Rightarrow 8c^2 (1 - \sin^2 A) = 9c^2 - 9a^2$$

$$\Rightarrow \sin^2 A = \frac{9a^2 - c^2}{8c^2} \Rightarrow \sin A = \sqrt{\frac{(9a^2 - c^2)}{8c^2}}$$

19. (c) : Given relation is $y = \frac{A}{x} + B$

Differentiating w.r.t. x , we get

$$y_1 = -\frac{A}{x^2} \Rightarrow x^2 y_1 = -A$$

Again, differentiating w.r.t. x , we get

$$x^2 y_2 + y_1 \cdot 2x = 0 \Rightarrow xy_2 + 2y_1 = 0$$

20. (a) : $T_{17} = T_{16+1} = {}^{50}C_{16}(2)^{50-16}a^{16} = {}^{50}C_{16}2^{34}a^{16}$

Similarly, $T_{18} = T_{17+1} = {}^{50}C_{17}2^{33}a^{17}$

Given that $T_{17} = T_{18}$

So, ${}^{50}C_{16}(2)^{34}a^{16} = {}^{50}C_{17}(2)^{33}a^{17}$

Therefore $\frac{{}^{50}C_{16} \cdot 2^{34}}{{}^{50}C_{17} \cdot 2^{33}} = \frac{a^{17}}{a^{16}}$

$$\Rightarrow a = \frac{{}^{50}C_{16} \times 2}{{}^{50}C_{17}} = \frac{50!}{16! \times 34!} \times \frac{17! \times 33!}{50!} \times 2 = 1$$

21. (a) : Let $I = \int_{-\pi/4}^{\pi/4} \frac{dx}{1 + \cos 2x} = \int_{-\pi/4}^{\pi/4} \frac{dx}{2 \cos^2 x}$

$$= \frac{1}{2} \int_{-\pi/4}^{\pi/4} \sec^2 x dx = \int_0^{\pi/4} \sec^2 x dx$$

$$\left[\text{Using property for even function } f(x), \int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx \right]$$

$$= [\tan x]_0^{\pi/4} = \left[\tan \frac{\pi}{4} - \tan 0 \right] = 1$$

22. (d) : Given, $x^m y^n = (x + y)^{m+n}$

Taking log on both sides, we get

$$m \log x + n \log y = (m + n) \log(x + y)$$

Differentiating w.r.t. x , we get

$$\begin{aligned} \frac{m}{x} + \frac{n}{y} \frac{dy}{dx} &= \frac{m+n}{x+y} \times \left(1 + \frac{dy}{dx} \right) \\ \Rightarrow \frac{m}{x} - \frac{m+n}{x+y} &= \left(\frac{m+n}{x+y} - \frac{n}{y} \right) \frac{dy}{dx} \\ \Rightarrow \frac{my - nx}{x(x+y)} &= \left(\frac{my - nx}{y(x+y)} \right) \frac{dy}{dx} \Rightarrow \frac{dy}{dx} = \frac{y}{x} \end{aligned}$$

23. (d) : We have $f(x) = x e^{1-x}$

$$f'(x) = -x e^{1-x} + e^{1-x} = (1-x)e^{1-x}$$

If $f'(x) < 0 \Rightarrow (1-x)e^{1-x} < 0$

$\Rightarrow 1-x < 0 (\because e^{1-x} > 0)$

$\Rightarrow x > 1 \Rightarrow x \in (1, \infty)$ so function is strictly decreasing.

24. (b) : Number of ways in which exactly 4 out of 7 cards go into envelopes of respective colour is equal to selecting 3 cards out of 7 which is equal to ${}^7C_3$ and then dearranging them i.e., none of the 3 cards go into its respective envelope.

$\therefore$ Required ways $= {}^7C_3 \times 3! \left[1 - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} \right]$

$= {}^7C_3 \times 3! \left[\frac{1}{2} - \frac{1}{6} \right] = {}^7C_3 \times 6 \times \frac{1}{3} = 2 \cdot {}^7C_3$

25. (b) : Applying $C_2 \rightarrow C_2 - C_1$, $C_3 \rightarrow C_3 - C_1$, we get

$$\Delta = \begin{vmatrix} a+1 & 1 & 3 \\ a+3 & 2 & 5 \\ a+7 & 3 & 7 \end{vmatrix}$$

Applying $R_2 \rightarrow R_2 - R_1$, $R_3 \rightarrow R_3 - R_1$, we get

$$\Delta = \begin{vmatrix} a+1 & 1 & 3 \\ 2 & 1 & 2 \\ 6 & 2 & 4 \end{vmatrix} = 0 - (8-12) + 3(4-6) = -2$$

26. (a) : Let the centre of the circle is $(h, 0)$, where $h > 0$. Now, circle passes through $(0, 1)$

$\therefore r^2 = (h-0)^2 + (0-1)^2$

$\Rightarrow (\sqrt{17})^2 = h^2 + 1$

$\Rightarrow h^2 = 16 \Rightarrow h = \pm 4$

$\Rightarrow h = 4$ ($\because h$ can't be negative)

$\therefore$ Equation of circle is $(x-4)^2 + (y-0)^2 = 17$

$\Rightarrow x^2 - 8x + y^2 + 16 - 17 = 0 \Rightarrow x^2 + y^2 - 8x - 1 = 0$

27. (b) : Let $g(x_1) = g(x_2)$

$\Rightarrow x_1^2 - 4x_1 - 5 = x_2^2 - 4x_2 - 5 \Rightarrow x_1^2 - x_2^2 = 4(x_1 - x_2)$

$\Rightarrow (x_1 - x_2)(x_1 + x_2 - 4) = 0$

Either $x_1 = x_2$ or $x_1 + x_2 = 4$

Either $x_1 = x_2$ or $x_1 = 4 - x_2$

$\therefore$ There are two values of x for which $g(x_1) = g(x_2)$.

$\therefore g(x)$ is not one-one $\forall x \in R$

28. (a) : Let e_1 and e_2 be the eccentricity of given hyperbola and ellipse respectively.

$\therefore e_1 = \sqrt{1 + \frac{b^2}{a^2}}$ and $e_2 = \sqrt{1 - \frac{1}{9}} = \sqrt{\frac{8}{9}}$

Given, $e_1 = \frac{1}{e_2} \Rightarrow 1 + \frac{b^2}{a^2} = \frac{9}{8} \Rightarrow \frac{a^2}{b^2} = \frac{8}{1}$

29. (d) : We have, $\sin \left[\frac{\pi}{3} - \sin^{-1} \left(\frac{-1}{2} \right) \right]$

$= \sin \left[\frac{\pi}{3} + \sin^{-1} \left(\frac{1}{2} \right) \right] = \sin \left[\frac{\pi}{3} + \frac{\pi}{6} \right] = \sin \left(\frac{\pi}{2} \right) = 1$

30. (a) : We have, $y = 4 + 3x - x^2$, a parabola with vertex at $\left(\frac{3}{2}, \frac{25}{4} \right)$

Putting $y = 0$, we get $x^2 - 3x - 4 = 0$

$\Rightarrow (x-4)(x+1) = 0 \Rightarrow x = -1$ or $x = 4$

$\therefore$ Required area $= \int_{-1}^4 (4 + 3x - x^2) dx$

$= \left[4x + \frac{3x^2}{2} - \frac{x^3}{3} \right]_{-1}^4 = \frac{125}{6}$ sq. units

31. (a) : Since, $(\vec{a} \times \vec{b})^2 + (\vec{a} \cdot \vec{b})^2 = 676$

$\Rightarrow (|\vec{a}| \cdot |\vec{b}| \sin \theta)^2 + (|\vec{a}| \cdot |\vec{b}| \cos \theta)^2 = 676$

$\Rightarrow |\vec{a}|^2 \cdot |\vec{b}|^2 (\sin^2 \theta + \cos^2 \theta) = 676$

$\Rightarrow |\vec{a}|^2 (2)^2 = 676 \Rightarrow |\vec{a}|^2 = 169 \Rightarrow |\vec{a}| = 13$

32. (b) : We have, $|A| = 6 + 1 = 7$

Also, $\text{adj } A = \begin{bmatrix} 2 & -1 \\ 1 & 3 \end{bmatrix}$

Now, $A^{-1} = \frac{1}{|A|} \text{adj } (A) = \frac{1}{7} \begin{bmatrix} 2 & -1 \\ 1 & 3 \end{bmatrix}$

$\therefore 14A^{-1} = \begin{bmatrix} 4 & -2 \\ 2 & 6 \end{bmatrix}$

33. (c) : $P(\text{atleast one of } A \text{ and } B)$

$= P(A \cup B) = P(A) + P(B) - P(A \cap B)$

$= P(A) + P(B) - P(A)P(B)$ [$\because A, B$ are independent]

$= P(A) + P(B) [1 - P(A)] = [1 - P(A')] + P(B)P(A')$

$= 1 - P(A') + P(B)P(A') = 1 - P(A') [1 - P(B)]$

$= 1 - P(A')P(B')$

34. (c) : $\frac{z_2}{z_1} = ik$, k is real, $z_2 = ikz_1$

$\Rightarrow \frac{6z_1 - 8z_2}{4z_2 + 3z_1} = \frac{6z_1 - 8ikz_1}{4ikz_1 + 3z_1} = \frac{6 - 8ik}{4ik + 3}$

$\therefore \left| \frac{6z_1 - 8z_2}{4z_2 + 3z_1} \right| = \left| \frac{6 - 8ik}{4ik + 3} \right| = \frac{\sqrt{36 + 64k^2}}{\sqrt{16k^2 + 9}} = \sqrt{4} = 2$

35. (a) : From the given relation

$\sin C = \frac{1 - \cos A \cos B}{\sin A \sin B} \leq 1$... (i)

$\Rightarrow 1 \leq \cos A \cos B + \sin A \sin B$

$\Rightarrow 1 \leq \cos (A - B)$ but $\cos (A - B) \leq 1$

$$\Rightarrow \cos(A - B) = 1 \Rightarrow A - B = 0 \Rightarrow A = B \quad \dots(ii)$$

$$\text{From (i) and (ii), we get, } \sin C = \frac{1 - \cos^2 A}{\sin^2 A} = \frac{\sin^2 A}{\sin^2 A} = 1$$

$$\therefore \angle C = 90^\circ$$

$$\Rightarrow A + B = 90^\circ \Rightarrow A = B = 45^\circ \quad [\text{Using (ii)}]$$

$$\therefore a : b : c = \sin A : \sin B : \sin C = \frac{1}{\sqrt{2}} : \frac{1}{\sqrt{2}} : 1 = 1 : 1 : \sqrt{2}$$

36. (a) : We have,

$$\int_0^\infty \frac{dx}{(x^2 + 4)(x^2 + 9)} = \frac{1}{5} \int_0^\infty \left[\frac{1}{x^2 + 4} - \frac{1}{x^2 + 9} \right] dx$$

$$= \frac{1}{5} \left[\left[\frac{1}{2} \tan^{-1} \frac{x}{2} \right]_0^\infty - \left[\frac{1}{3} \tan^{-1} \frac{x}{3} \right]_0^\infty \right]$$

$$= \frac{1}{5} \left[\frac{1}{2} \tan^{-1} \infty - \frac{1}{3} \tan^{-1} \infty \right] = \frac{1}{5} \left[\frac{\pi}{4} - \frac{\pi}{6} \right] = \frac{\pi}{60}$$

37. (b) : $\lim_{x \rightarrow 0} \frac{\pi^x - 1}{\sqrt{1+x} - 1} \quad \left(\frac{0}{0} \text{ form} \right)$

$$= \lim_{x \rightarrow 0} \frac{\pi^x \log_e \pi}{\frac{1}{2\sqrt{1+x}} - 0} \quad [\text{Using L'Hospital's rule}]$$

$$= 2 \log_e \pi = \log_e (\pi^2)$$

38. (c) : 4th term from the end is $9 - 4 + 2$, i.e., 7th term from the beginning.

$$\therefore T_7 = {}^9C_6 \left(\frac{x^3}{2} \right)^3 \left(\frac{-2}{x^2} \right)^6 = {}^9C_3 \frac{x^9}{8} \cdot \frac{64}{x^{12}}$$

$$= \frac{9 \times 8 \times 7}{3 \times 2 \times 1} \times \frac{8}{x^3} = \frac{672}{x^3}$$

39. (b) : Let $\frac{x+1}{-3} = \frac{y-3}{2} = \frac{z+2}{1} = \lambda$

$\therefore$ The coordinates of a point on the line are $(-3\lambda - 1, 2\lambda + 3, \lambda - 2)$.

Similarly, let $\frac{x}{1} = \frac{y-7}{-3} = \frac{z+7}{2} = \mu$

The coordinates of a point on this line are $(\mu, -3\mu + 7, 2\mu - 7)$.

Since, two lines intersect for same values of λ and μ ,

$$\therefore (-3\lambda - 1, 2\lambda + 3, \lambda - 2) = (\mu, -3\mu + 7, 2\mu - 7)$$

$$\Rightarrow -3\lambda - 1 = \mu, 2\lambda + 3 = -3\mu + 7, \lambda - 2 = 2\mu - 7$$

$$\therefore -3\lambda - \mu = 1 \quad \dots(i)$$

$$2\lambda + 3\mu = 4 \quad \dots(ii)$$

$$\lambda - 2\mu = -5 \quad \dots(iii)$$

Solving equation (i) and (ii), we get $\lambda = -1$ and $\mu = 2$ and 3rd equation holds for these values.

$\therefore$ The lines intersect at the point $(2, 1, -3)$.

40. (d) : Variance $= \sigma^2 \Rightarrow \frac{1}{n} \sum_{i=1}^n x_i^2 - \left(\frac{1}{n} \sum_{i=1}^n x_i \right)^2 = \sigma^2$

$$\Rightarrow 0 = \frac{1}{n} (400) - (5)^2 \Rightarrow \frac{400}{n} = 25 \Rightarrow n = 16$$

41. (c) : From given differential equation, we have

$$\frac{dy}{dx} = \frac{e^x + x^2}{e^y}$$

Using variable separable form, we have

$$e^y dy = (e^x + x^2) dx$$

On integrating, we get $e^y = e^x + x^3/3 + c$

42. (b) : Let the slope of line through the point $(3, -2)$ be m .

Now, slope of line $\sqrt{3}x + y = 1$ is $-\sqrt{3}$.

$$\therefore \tan 60^\circ = \left| \frac{-\sqrt{3} - m}{1 + (-\sqrt{3})m} \right| \Rightarrow \sqrt{3} = \left| \frac{\sqrt{3} + m}{1 - \sqrt{3}m} \right|$$

$$\Rightarrow \pm \sqrt{3} = \frac{\sqrt{3} + m}{1 - \sqrt{3}m} \Rightarrow m = \sqrt{3}, 0$$

$\therefore$ Equations of lines passing through $(3, -2)$ and

having slopes $\sqrt{3}, 0$ are $y + 2 = \sqrt{3}(x - 3)$ and $y + 2 = 0$ (rejected) $\Rightarrow y - \sqrt{3}x + 3\sqrt{3} + 2 = 0$

43. (d) : Given, $x^2 - \frac{3ax}{2} + 1 - a^2$ is positive.

$$\Rightarrow x^2 - \frac{3ax}{2} + 1 - a^2 > 0$$

$$\therefore 9a^2 - 4(2)(2 - 2a^2) < 0 \quad (\because \text{If } f(x) > 0 \therefore D < 0)$$

$$\Rightarrow 9a^2 - 16 + 16a^2 < 0 \Rightarrow 25a^2 - 16 < 0$$

$$\Rightarrow 25a^2 < 16 \Rightarrow a^2 < \frac{16}{25} \Rightarrow |a| < \frac{4}{5}$$

44. (b) : Since, $P(3)$ is true and $P(k)$ is true

$\Rightarrow P(k + 1)$ is true, which means, if $P(3)$ is true

$\Rightarrow P(4)$ is true $\Rightarrow P(5)$ is true and so on. So statement is true for all $n \geq 3, n \in \mathbb{N}$.

45. (b) : $(A \cap B) \cup (A - B) = (A \cap B) \cup (A \cap B')$
 $[\because (A - B) = (A \cap B')]$
 $= A \cap (B \cup B') \quad (\text{By distributive law})$
 $= A \cap U = A \quad (\because B \cup B' = U)$

46. (c) : Let $y = a \sin^3 t$ and $x = a \cos^3 t$

$$\therefore \frac{dy}{dt} = 3a \sin^2 t \cos t \text{ and } \frac{dx}{dt} = -3a \cos^2 t \sin t$$

$$\Rightarrow \frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{\sin t}{-\cos t} = -\tan t$$

$$\Rightarrow \frac{d^2 y}{dx^2} = -\sec^2 t \cdot \frac{dt}{dx} = \frac{-\sec^2 t}{-3a \cos^2 t \sin t} = \frac{1}{3a \cos^4 t \sin t}$$

$$\therefore \left(\frac{d^2 y}{dx^2} \right)_{t=\pi/4} = \left(\frac{1}{3a \cos^4 t \sin t} \right)_{t=\pi/4} = \frac{4\sqrt{2}}{3a}.$$

47. (a) : To check continuity at $x = 1$

$$\text{L.H.L.} = \lim_{x \rightarrow 1^-} f(x) = \lim_{h \rightarrow 0} (1 - (1-h)^2) = 1 - 1 = 0$$

$$\text{R.H.L.} = \lim_{x \rightarrow 1^+} f(x) = \lim_{h \rightarrow 0} (1 + (1+h)^2) = 1 + 1 = 2$$

$$\therefore \text{L.H.L.} \neq \text{R.H.L.} \neq f(1)$$

$\therefore f(x)$ is not continuous at $x = 1$

48. (b) : Let $I = \int 1 \cdot \sin^{-1} x dx$

$$= (\sin^{-1} x)x - \int \frac{1}{\sqrt{1-x^2}} \cdot x dx + C'$$

Putting $1 - x^2 = t^2 \Rightarrow -2x dx = 2t dt$, we get

$$I = x \sin^{-1} x - \int \frac{(-t dt)}{t} + C' = x \sin^{-1} x + \sqrt{1-x^2} + C$$

49. (b) : From given equations, we have

$$\frac{x^2}{4} + \frac{y^2}{9} = 1 \text{ and } \frac{x^2}{4} - \frac{y^2}{9} = 1$$

$$\therefore e_1^2 = \frac{9-4}{9} \text{ and } e_2^2 = \frac{4+9}{4}$$

$$\text{Now, } e_1^2 + e_2^2 = \left(\frac{5}{9} \right) + \left(\frac{13}{4} \right) = \frac{137}{36} > 3 \text{ but } < 4$$

$$\therefore 3 < e_1^2 + e_2^2 < 4$$

$$50. (d) : \text{Let } D = \begin{vmatrix} y+k & y & y \\ y & y+k & y \\ y & y & y+k \end{vmatrix}$$

$$= (y+k)((y+k)^2 - y^2) - y(y^2 + ky - y^2) + y(y^2 - y^2 - yk)$$

$$= k^3 + 3k^2y = k^2(k + 3y)$$

$$51. (c) \quad 52. (d) \quad 53. (a)$$

$$54. (c) \quad 55. (a) \quad 56. (a)$$

$$57. (b) \quad 58. (c) \quad 59. (d)$$

$$60. (d) \quad 61. (a) \quad 62. (a)$$

$$63. (c) \quad 64. (b) \quad 65. (c)$$

$$66. (a, b, c) \quad 67. (a, b, c, d) \quad 68. (c, d)$$

$$69. (a, c, d) \quad 70. (a, c) \quad 71. (a, b, c)$$

$$72. (a, c) \quad 73. (a, b) \quad 74. (a, c)$$

$$75. (a, c)$$



UNSCRAMBLE ME

Unscramble the words given in column I and match them with their explanations in column II.

Column I

1. COCAPIRREL
2. TILEPERNEC
3. VINTNAIRA
4. OILBUEQ
5. ERESONISGR
6. VINALTER
7. RIVELTIAMULBA
8. ANARICOCEV
9. UNTIQUOET
10. SUTESREP

Column II

- (a) A statistical technique that relates a dependent variable to one or more independent variable.
- (b) In probability theory and statistics it is used to measure the joint variability of two random variables.
- (c) A straight line that is neither horizontal nor vertical.
- (d) The numbers resulting from the division of one number by another.
- (e) The multiplicative inverse of a value or a number.
- (f) A set that contains all the elements of another set.
- (g) A specific set of real numbers that lies between two conditional numbers of the set.
- (h) A function, quantity which remains unchanged when specified transformation is applied.
- (i) Two or more variables.
- (j) It is used in statistics indicating the value below which a given percentage of observations in a group of observations fall.

Readers can send their responses at editor@mtg.in or post us with complete address by 10th of every month. Winners' names will be published in next issue.

Are You Ready for JEE?



Top Weightage Topics for JEE Series 1

In this article we will discuss about some of the most important topics of mathematics in respect to JEE Main and Advanced. It is based on the analysis of previous years' questions of JEE Main and Advanced.

Sets

A well defined collection of distinct objects.

Representation of Sets


Set builder form or rule method:

$A = \{x : x \text{ is a prime number}\}$

Roster or tabular form :

$A = \{1, 2, 3, 4, 5\}$.

Intersection of two sets



$$A \cap B = \{x : x \in A \text{ and } x \in B\}$$

Order/Cardinality

If $B = \{\text{Red, Blue, White}\}$, then total number of element of B, i.e., $n(B) = 3$

Types of Sets

Empty or null set : $A = \phi$ or $\{\}$

Singleton set : $A = \{a\}$

Finite set : $A = \{1, 2, 3, 4, 5\}$

Infinite sets : $A = \{1, 2, 3, 4, \dots\}$

Equal sets : $A = \{1, 2, 3, 4\}$

$B = \{3, 1, 4, 2\} \Rightarrow A = B$


Equivalent set : $A = \{1, 2, 3\}$ and $B = \{2, 5, 6\} \Rightarrow n(A) = n(B)$

Power set : $A = \{1, 2, 3\}$

$P(A) = \{\{1, 2, 3\}, \{1, 2\}, \{1, 3\}, \{2, 3\}, \{1\}, \{2\}, \{3\}, \phi\}$


Disjoint sets : $A \cap B = \phi$

Union of two sets



$$A \cup B = \{x : x \in A \text{ or } x \in B\}$$

Symmetric difference of A and B




$$A \Delta B = (A - B) \cup (B - A)$$
$$x \in A \cup B \text{ but } x \notin A \cap B$$

Demorgan's law

$$(A \cup B)' = A' \cap B'$$

$$(A \cap B)' = A' \cup B'$$

Compliment of a Set



$$A' \text{ or } A^C = U - A$$

Relations

If A and B are two non empty sets, then relation R from A to B is a subset of $A \times B$.

Reflexive : A relation R on a set A is reflexive if $(a, a) \in R \forall a \in A$.

Symmetric : A relation R on a set A is symmetric if $(a_1, a_2) \in R \Rightarrow (a_2, a_1) \in R \forall a_1, a_2 \in A$.

Types of Relations

Transitive : A relation R on a set A is transitive if $(a_1, a_2) \in R$ and $(a_2, a_3) \in R \Rightarrow (a_1, a_3) \in R \forall a_1, a_2, a_3 \in A$.

Equivalence : A relation R is an equivalence relation if R is reflexive, symmetric and transitive.




Functions

A relation 'f' from a set A to a set B is said to be a function if every element of set A has one and only one image in set B.

Identity Function

$$y = f(x) = x \forall x \in R,$$


$$D_f = R = R_f$$



Logarithmic Function

$$y = f(x) = \log_a x, a > 0, a \neq 1 \text{ and } x > 0$$

$$D_f = (0, \infty), R_f = R.$$




Constant Function

$$y = f(x) = c,$$

$\forall x \in R$, where fixed number c is called a constant.

$$D_f = R$$

$$R_f = \{c\}$$




Modulus (Absolute Value) Function

$$y = f(x) = |x|, \text{ where}$$

$$|x| = \begin{cases} x, & \text{if } x \geq 0 \\ -x, & \text{if } x \leq 0 \end{cases}$$

$$D_f = R$$

$$R_f = [0, \infty)$$




Greatest Integer Function

$$y = f(x) = [x],$$

$$D_f = R$$

$$R_f = Z \text{ (Set of all integers)}$$




Signum Function

$$y = f(x) = \text{sgn}(x),$$

$$\text{where } f(x) = \begin{cases} 1, & \text{if } x > 0 \\ 0, & \text{if } x = 0 \\ -1, & \text{if } x < 0 \end{cases}$$

$$D_f = R$$

$$R_f = \{-1, 0, 1\}$$




Exponential Function

$$y = f(x) = a^x, a > 0, a \neq 1,$$

$$D_f = R,$$

$$R_f = (0, \infty)$$




Inverse of a Function

Let $f: A \rightarrow B$ is a objective function,

then define $g: B \rightarrow A$ such that

$$f(x) = y \Rightarrow g(y) = x$$



Even and Odd Function

Even function : $f(-x) = f(x)$

Odd function : $f(-x) = -f(x)$



Arithmetic Progression (A.P.)

A sequence in which each term is obtained by adding a fixed number to the preceding term except the first term.

General form of an A.P.

$a, a + d, a + 2d, \dots$,
where a = first term and d = common difference

Sum of n terms of an A.P.

$S_n = \frac{n}{2} [2a + (n-1)d]$ or $S_n = \frac{n}{2} [a + l]$,
where l = last term

General term (or n^{th} term) of an A.P.

$$a_n = n^{\text{th}} \text{ term} = a + (n-1)d$$

n^{th} term from last of an A.P.

$$a'_n = l - (n-1)d,$$

where n = number of terms

Arithmetic mean A.M.

- For two numbers a and b , A.M. is $\frac{a+b}{2}$
- $A_r = a + r \left(\frac{b-a}{n+1} \right), \forall r = 1, 2, \dots, n$

where $A_1, A_2, \dots, A_n$ are n arithmetic means inserted between two numbers a and b .



Geometric Progression (G.P.)

A sequence in which each term is obtained by multiplying the previous term by a non zero constant.

General form of a G.P.

$$a, ar, ar^2, \dots,$$

where a = first term and r = common ratio

General term (or n^{th} term) of a G.P.

$$a_n = ar^{n-1}$$

n^{th} term from the last of a G.P.

$$a'_n = \frac{l}{r^{n-1}}$$

Sum of ' n ' term of a G.P.

$$S_n = \frac{a(1-r^n)}{1-r} \quad \text{or} \quad S_n = \frac{a(r^n-1)}{r-1}, r \neq 1$$

or $S_n = na, r = 1$

Sum of infinite terms of a G.P.

$$S_\infty = \frac{a}{1-r}; |r| < 1$$

Geometric mean (G.M.)

- For two numbers a and b , G.M. is $\sqrt{ab}$

$$G_k = a \left(\frac{b}{a} \right)^{\frac{k}{n+1}}, \forall r = 1, 2, \dots, n$$

where $G_1, G_2, \dots, G_n$ are n geometric means inserted between two numbers a and b .

Relation between A.M. and G.M.

- If $a_1, a_2, a_3, \dots, a_n$ are ' n ' positive numbers, then $\text{A.M.} \geq \text{G.M.}$

i.e., $\frac{a_1 + a_2 + \dots + a_n}{n} \geq (a_1 a_2 \dots a_n)^{\frac{1}{n}}$; Equality holds if $a_1 = a_2 = \dots = a_n$



Permutation

Arrangement of number of things taken some or all of them at a time.

Notation

Number of ways of arranging ' n ' distinct things taken ' r ' at a time

$$\text{i.e., } {}^n P_r = P(n, r) = \frac{n!}{(n-r)!}$$

Properties

- ${}^n P_n = n!$
- ${}^n P_1 = n$
- ${}^n P_0 = 1$
- ${}^n P_r = n \cdot {}^{n-1} P_{r-1}$
- $\frac{{}^n P_r}{{}^n P_{r-1}} = n - r + 1$
- ${}^{n-1} P_r + r {}^{n-1} P_{r-1} = {}^n P_r$

Circular Permutation

- Number of circular permutations of n distinct object: $(n-1)!$
- Number of circular permutation of n distinct objects taken r at a time: ${}^n C_r (r-1)!$

Restricted Permutations

Number of permutations of n objects taken r at a time, when a particular object is :

- Always included (or never excluded)
= $r \cdot ({}^{n-1} P_{r-1})$
- Always excluded (or never included) = ${}^{n-1} P_r$



Combination

Different groups or selection which can be made by taking some or all the objects.

Notation

Number of selection of ' n ' distinct things taken ' r ' at a time, $0 \leq r \leq n$

$$\text{i.e., } {}^n C_r = C(n, r) = \frac{n!}{(r)! (n-r)!}$$

Properties

- ${}^n C_r = {}^n C_{n-r}$
- If ${}^n C_x = {}^n C_y$
 $\Rightarrow x = y$ or $x + y = n$
- ${}^n C_r + {}^n C_{r-1} = {}^{n+1} C_r$

Relation between Permutation and Combination

- In permutation, we deal with arrangements of objects whereas in combination, we have to deal with selection of objects.
- Order of objects is not important in combination but has importance in permutation.



Logarithm

Logarithm is the power to which a number must be raised in order to get some other number.

Properties of Logarithm

- $\log_b 1 = 0$
- $\log_b b = 1$
- $\log_b b^x = x$
- $b^{\log_b x} = x$
- $\log_b a = \log_c a \log_b c$
- $\log_b a = \frac{1}{\log_a b}$
- $\log_b a = \frac{\log_c a}{\log_c b}$
- $\log_b (xy) = \log_b x + \log_b y$
- $\log_b a^\alpha = \alpha \log_b a$
- $\log_\beta a = \frac{1}{\beta} \log_b a$
- $\log_\beta a^\alpha = \frac{\alpha}{\beta} \log_b a$
- $a^{\log_c b} = b^{\log_c a}$ ($a, b, c > 0$ and $c \neq 1$)
- $a^{\sqrt{\log_a b}} = b^{\sqrt{\log_b a}}$
- $\log_2 \log_2 \sqrt{\sqrt{\sqrt{\dots \sqrt{2}}}} = -n$, where n is the number of square roots.



Binomial Theorem

The algebraic expansion of powers of a binomial.

Binomial theorem for any positive integer n .

$$(x + y)^n = {}^nC_0 x^n y^0 + {}^nC_1 x^{n-1} y^1 + {}^nC_2 x^{n-2} y^2 + \dots + {}^nC_n x^0 y^n$$

$$\text{i.e., } (x + y)^n = \sum_{r=0}^n {}^nC_r x^{n-r} y^r,$$

where ${}^nC_0, {}^nC_1, {}^nC_2, \dots, {}^nC_n$ are known as binomial coefficients.

- $(x - y)^n = \sum_{r=0}^n {}^nC_r (-1)^r x^{n-r} y^r$
- $(1 + x)^n = \sum_{r=0}^n {}^nC_r x^r$
- $(1 - x)^n = \sum_{r=0}^n (-1)^r {}^nC_r x^r$

Properties of Binomial Coefficients

- $C_0 + C_1 + C_2 + \dots + C_n = 2^n$
- $C_0 + C_2 + C_4 + \dots = C_1 + C_3 + C_5 + \dots = 2^{n-1}$
- $C_0 - C_1 + C_2 - \dots + (-1)^n C_n = 0$
- $C_0^2 + C_1^2 + C_2^2 + \dots + C_n^2 = 2^n C_n$
- $C_0 C_r + C_1 C_{r+1} + C_2 C_{r+2} + \dots + C_{n-r} C_n = 2^n C_{n-r}$
- $\sum_{r=0}^n r {}^nC_r = n 2^{n-1}$
- $\sum_{r=0}^n r(r-1) {}^nC_r = n(n-1) 2^{n-2}$
- $C_0 + \frac{C_1}{2} + \frac{C_2}{3} + \dots + \frac{C_n}{n+1} = \frac{2^{n+1} - 1}{n+1}$
- $C_0 - \frac{C_1}{2} + \frac{C_2}{3} - \frac{C_3}{4} + \dots + \frac{(-1)^n C_n}{n+1} = \frac{1}{n+1}$

Number of terms

- For index n , $(a + b)^n$ has $(n + 1)$ terms.
- $(a + b)^n + (a - b)^n = 2 \{ {}^nC_0 a^n + {}^nC_2 a^{n-2} b^2 + {}^nC_4 a^{n-4} b^4 + \dots \}$ and $(a + b)^n - (a - b)^n = 2 \{ {}^nC_1 a^{n-1} b^1 + {}^nC_3 a^{n-3} b^3 + {}^nC_5 a^{n-5} b^5 + \dots \}$
- If n is odd, then number of terms in both $\{(a + b)^n + (a - b)^n\}$ and $\{(a + b)^n - (a - b)^n\} = (n+1)/2$
- If n is even, then number of terms in $\{(a + b)^n + (a - b)^n\} = n/2 + 1$; $\{(a + b)^n - (a - b)^n\} = n/2$



Complex Number

Combination of both the real number and imaginary number.

Definition

A number of the form $a + ib$, where $a, b \in \mathbb{R}$ and $i = \sqrt{-1}$ is called a complex number and denoted by 'z'.

$$z = a + ib$$

Power of i


Integral powers of i

$$i^0 = 1, i^1 = i, i^2 = -1, i^3 = i^2(i) = -i, i^4 = (i^2)^2 = (-1)^2 = 1$$

In general, for any integer k , $i^{4k} = 1$, $i^{4k+1} = i$, $i^{4k+2} = -1$, $i^{4k+3} = -i$

Argand Plane

$z = a + ib$ is represented by a point $P(a, b)$



Square root of complex number

$$\sqrt{a + ib} = \begin{cases} \pm \left(\sqrt{\frac{|z| + a}{2}} + i \sqrt{\frac{|z| - a}{2}} \right), & \text{for } b > 0 \\ \pm \left(\sqrt{\frac{|z| + a}{2}} - i \sqrt{\frac{|z| - a}{2}} \right), & \text{for } b < 0 \end{cases}$$

Modulus, Conjugate & Argument

Modulus of $z = a + ib$ is


$$r = |z| = \sqrt{a^2 + b^2}$$

Conjugate of $z = a + ib$ is

$$\bar{z} = a - ib$$


Argument of $z = a + ib$ is

$$\theta = \tan^{-1} \left(\frac{\text{Im}(Z)}{\text{Re}(Z)} \right), \quad -\pi < \theta \leq \pi$$



Polar representation

$$Z = r[\cos(2n\pi + \theta) + i \sin(2n\pi + \theta)], \quad n \in \mathbb{I}$$



Cube roots of unity

$$\text{Let } x = (1)^{1/3} \Rightarrow x^3 - 1 = 0$$

$$\Rightarrow x = 1, \frac{-1 + i\sqrt{3}}{2}, \frac{-1 - i\sqrt{3}}{2} \text{ or } x = 1, \omega, \omega^2$$

$$\bullet \quad \omega^3 = 1$$

$$\bullet \quad 1 + \omega + \omega^2 = 0$$

$$\bullet \quad 1 + \omega^n + \omega^{2n} = \begin{cases} 0, & \text{if } n \text{ is a not multiple of } 3 \\ 3, & \text{if } n \text{ is a not multiple of } 3 \end{cases}$$

Inequalities in complex numbers

If z_1 and z_2 are two complex numbers such that $|z_1| \leq 1$, $|z_2| \leq 1$, then

- $|z_1 - z_2|^2 \leq (|z_1| - |z_2|)^2 + (\text{Arg}(z_1) - \text{Arg}(z_2))^2$
- $|z_1 + z_2|^2 \geq (|z_1| + |z_2|)^2 - (\text{Arg}(z_1) - \text{Arg}(z_2))^2$
- $|z_1 \pm z_2| \leq |z_1| + |z_2|$
In general $|z_1 \pm z_2 \pm z_3 \pm \dots \pm z_n| \leq |z_1| + |z_2| + |z_3| + \dots + |z_n|$
- $|z_1 \pm z_2| \geq ||z_1| - |z_2||$
 $\Rightarrow ||z_1| - |z_2|| \leq |z_1 + z_2| \leq |z_1| + |z_2|$



Quadratic Equations

An equation of the form $ax^2 + bx + c, a \neq 0$ where a, b, c are real or complex number.

Common Roots of Quadratic Equation

Condition of one common root

Let $a_1x^2 + b_1x + c_1 = 0$ and $a_2x^2 + b_2x + c_2 = 0$ have a common root α .

$$\therefore \alpha = \frac{b_1c_2 - b_2c_1}{a_2c_1 - a_1c_2} = \frac{a_2c_1 - a_1c_2}{a_1b_2 - a_2b_1}, \alpha \neq 0$$

Condition for both the common roots


$$\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$$

Quadratic Expression and its Graph

Case-I : For $a > 0$

and $D > 0$


For $\alpha < x < \beta$,
 $y < 0$



Case-III : For $a > 0$

and $D < 0$


In this case, imaginary roots appears &
 $y > 0$ for $x \in \mathbb{R}$.



Case-II : For $a > 0$

and $D = 0$


$y = a(x - \alpha)^2$
and $y \geq 0$, for $x \in \mathbb{R}$.



Case-IV : For $a < 0$

and $D > 0$

$y > 0$, if $\alpha < x < \beta$
 $y < 0$, if $x < \alpha$
or $x > \beta$



Matrices

Any rectangular arrangement of numbers.

Types of Matrices

Orthogonal matrix : $A = [a_{ij}]_{m \times n}$

if $AA' = I_n = A'A$, then A is orthogonal matrix.

Idempotent matrix : $A = [a_{ij}]_{n \times n}$

If $A^2 = A$, then A is idempotent matrix.

Involutory matrix : $A = [a_{ij}]_{n \times n}$

If $A^2 = I$, then A is involutory matrix.

Nilpotent matrix : $A = [a_{ij}]$

If a positive integer k such that $A^k = 0$, then A is Nilpotent matrix.

Unitary matrix : $A = [a_{ij}]$

If $A' A = I$, then A is unitary matrix.

Rank of matrix

Rank of matrix = Maximum number of linearly independent columns (or rows of a matrix)

Trace of matrix

If $A = [a_{ij}]_{n \times n}$, then, $tr(A) = \sum_{i=1}^n a_{ii}$

- $tr(A) + tr(B) = tr(A + B)$
- $tr(kA) = k tr(A)$

Order of matrix

A matrix having m rows and n columns is called matrix of order $m \times n$.



Determinants

A compact form showing a set of numbers arranged in same number of rows and columns.

Minors and Cofactors

- For any matrix $A = [a_{ij}]_{n \times n}$, if we leave the row and the column of the element a_{ij} , then the value of determinant thus obtained is called the minor of a_{ij} and it is denoted by M_{ij} .
- The minor M_{ij} multiplied by $(-1)^{i+j}$ is called the cofactor of the element a_{ij} and denoted by A_{ij} . $\therefore A_{ij} = (-1)^{i+j} M_{ij}$

Singular and Non-Singular Matrices

A square matrix A of order n is said to be

- Singular if $|A| = 0$
- Non-singular if $|A| \neq 0$

Properties

- $(A^{-1})^{-1} = A$
- $(A')^{-1} = (A^{-1})'$
- $(AB)^{-1} = B^{-1} A^{-1}$
- $(ABC)^{-1} = C^{-1} B^{-1} A^{-1}$

Inverse of a Matrix

For any square matrix A , inverse of A is defined as

$$A^{-1} = \frac{1}{|A|} (\text{adj } A), |A| \neq 0$$

Area of a triangle

Let ABC be a triangle with vertices $A(x_1, y_1)$, $B(x_2, y_2)$ and $C(x_3, y_3)$, then area of ΔABC is

$$\Delta = \frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix}$$

Adjoint of a Matrix

Let $B = [A_{ij}]$ be the matrix of cofactors of matrix $A = [a_{ij}]$. Then the transpose of B is called the adjoint of matrix A .

Properties of adj (A)

If A is non-singular matrix of order n , then

- $A(\text{adj } A) = (\text{adj } A)A = |A| I_n$
- $|A \text{ adj } A| = |A|^n$
- $\text{adj } (AB) = (\text{adj } B) \cdot (\text{adj } A)$
- $|\text{adj } A| = |A|^{n-1}$
- $\text{adj } (\text{adj } A) = |A|^{n-2} A$
- $|\text{adj } (\text{adj } A)| = |A|^{(n-1)^2}$

Solution of System of Linear Equations

Let $AX = B$ be the given system of equations :

- If $|A| \neq 0$, the system is consistent and has a unique solution.
- If $|A| = 0$ and $(\text{adj } A)B \neq O$, then the system is inconsistent and hence it has no solution.
- If $|A| = 0$ and $(\text{adj } A)B = O$, then the system may be either consistent or inconsistent according as the system has either infinitely many solutions or no solution.

Cramer's rule

- If $a_1x + b_1y = c_1$ and $a_2x + b_2y = c_2$ then,
 $x = \frac{D_1}{D}, y = \frac{D_2}{D}$
 where $D = \begin{vmatrix} a_1 & b_1 \\ a_2 & b_2 \end{vmatrix}, D_1 = \begin{vmatrix} c_1 & b_1 \\ c_2 & b_2 \end{vmatrix}$
- If $a_1x + b_1y + c_1z = d_1, a_2x + b_2y + c_2z = d_2$ and $a_3x + b_3y + c_3z = d_3$
 then, $x = \frac{D_1}{D}, y = \frac{D_2}{D}, z = \frac{D_3}{D}$,
 where $D \neq 0$




INDIAN ECONOMY

Union Budget 2023-24

- The Finance Minister of India introduced the Union Budget (for 2023-24) before the Parliament on February 1, 2023. Highlights of Budget 2023-24 are:
 - Changes in the new income tax regime (in rebate limit and in tax slabs).
 - A 33% increase in capital investment outlay has been proposed, raising it to ₹ 10 lakh crore (the biggest in the past decade).
 - Capital outlay for the railways increased to the highest ever – ₹ 2.40 lakh crore.

Priorities of Budget 2023-24



Nidhi Aapke Nikat 2.0 Programme

- Employees' Provident Fund Organisation (EPFO) has launched a massive District Outreach Program in all

the districts of the country through a revamped Nidhi Aapke Nikat program. The aim of this program is to reach all the districts of the country on the same day i.e. 27th of every month. For that, the EPFO has planned to organize camps in 685 districts of the country.

Complete Transition to 'Trade-Plus-One' (T+1) Settlement Cycle

- India becomes the second country in the world after China to start 'Trade-plus-one' (T+1) settlement cycle in top listed securities.
- T+1 (trade-plus-one) means that trade-related settlements will be cleared within a day of the actual transactions. Earlier, trades on the Indian exchanges were settled in two working days after the transaction took place (T+2).

Financial Stability Report (FSR)

- Recently, the RBI released its 26th FSR which reflects the collective assessment of the Sub-Committee of the Financial Stability and Development Council (FSDC) on risks to financial stability, as also the resilience of the financial system. It is released twice every year.

RBI's e₹

- The RBI launched Central Bank Digital Currency (CBDC) or e-Rupee (e₹) in 2022 for retail markets in a pilot mode. It is in the form of a digital token that represents legal tender. Reliance Retail becomes the first retail to accept payments via e-rupee at stores. Reliance retail has



partnered with ICICI Bank and Kotak Mahindra Bank to enable acceptance of RBI-issued Digital Rupee.

Global Wage Report 2022-23

- The Global Wages Report by the International Labour Organisation (ILO) presents an analysis of how concurrent crises – COVID-19 pandemic followed

by cost-of-living crisis – have impacted wages and purchasing power across countries and regions.

- Global wages declined in 2022 for the first time since the 2008 global financial crisis due to increasing living expenses. This is threatening to worsen inequality and trigger social unrest.

Test Yourself!

- Which of the following banks are categorised as Domestically Systemically Important Banks (D-SIBs) or banks “too big to fail”?
(i) State Bank of India (ii) Punjab National Bank
(iii) HDFC Bank (iv) ICICI Bank
Select the correct option.
(a) (i) and (ii) only (b) (ii) and (iii) only
(c) (i), (iii) and (iv) only (d) (ii), (iii) and (iv) only
- Which among the following organisation releases Global Wage Report 2022-23?
(a) Bank for International Settlements
(b) World Bank
(c) International Monetary Fund
(d) International Labour Organisation
- Recently, New T+1 settlement cycle has come into effect. Select the incorrect statement about T+1 Settlement.
(a) It means that trade-related settlements must be done within a day, or 24 hours, of the completion of a transaction.
(b) Until 2001, stock markets had a weekly settlement system.
(c) After China and UK, India will become the third country in the world to start the ‘trade-plus-one’ (T+1) settlement cycle in top listed securities.
(d) None of the above
- ‘Nidhi Apke Nikat’ is an outreach program conducted by which institution?
(a) Department of Legal Affairs
(b) Employees’ Provident Fund Organisation
(c) Department of Border Management
(d) Khadi and Village Industries Commission
- Recently, the Reserve Bank of India (RBI) released its 26th Financial Stability Report (FSR). It reflects the collective assessment of
(a) Macro-Economic Framework Statement
(b) Sub-Committee of the Financial Stability and Development Council
(c) Annual Financial Statement
(d) Board for Regulation and Supervision of Payment and Settlement Systems
- Consider the following statements about the budget allocations of fiscal year 2023-24 across the ministries and choose which is the correct answer?
(a) Ministry of Defence has the largest share in budget allocations
(b) Ministry of Railways has largest share in budget allocations
(c) Ministry of Agriculture and Farmer's Welfare has received highest allocation amongst the ministries
(d) None of the above
- RBI has launched a virtual form of currency under the name Digital Rupee. Who can invest in digital money?
(a) Investors
(b) Secondary market transactions in government securities
(c) Wholesale and retail customers
(d) All of the above
- Which of the following statements is incorrect about Indian economy?
(a) The capital expenditure of Union government has increased approximately four times since Fiscal year 2016
(b) The non-performing assets of banks have been falling for last 2 years
(c) There has been constant increase in forex reserves.
(d) All of the above
- According to Union Budget 2023-24, which among the following are included in seven priorities of government?
(i) Reaching the last mile (ii) Green growth
(iii) Reducing inequality (iv) Unleashing potential
(a) Only (i), (ii) and (iii) (b) Only (ii) and (iv)
(c) Only (i), (ii) and (iv) (d) All of the above
- According to the Union Budget, 2023-24, what is the rebate limit of Personal Income Tax in the new tax regime?
(a) 8 lakh (b) 6 lakh
(c) 9 lakh (d) 7 lakh

Answer Key

- (c)
- (d)
- (c)
- (b)
- (b)
- (a)
- (d)
- (c)
- (c)
- (d)

UCD

Unique Career in Demand

Explore the available Unique Career Options!



B.Sc. in Actuarial Science

Bachelor of science in Actuarial Science course is 3-year undergraduate course, where students learn about insurance risk calculation, insurance premiums, application of economic and mathematical analysis for making any decision in financial planning, insurance investment and management among all the others. This is the best course to learn core mathematical skills.

Higher Studies options after B.Sc in Actuarial Science

- Master of Science in Actuarial Science
- Post Graduate Diploma in Actuarial Science
- PG Diploma in Insurance and Risk Management
- PG Diploma in Insurance Science
- PG Diploma in management of Insurance and Financial Services
- B.Sc. Actuarial Science graduates can also opt for MBA course as it will give them in-depth knowledge about the organizational work structure.

Job Perspectives

Actuaries mostly work in Insurance and Reinsurance companies. They also work as consultants for companies, banking, financial consulting firms, actuarial consulting firms, the government sector, pension funds, investment advisories and healthcare units.

According to the Bureau of labour statistics, the number of actuaries employed is expected to grow 21% from 2021 to 2031. For this reason, many universities offering educational degrees and courses on Actuarial science.

Eligibility Criteria

- The admission for the Actuarial Science course will be based on the merit scored in the Actuarial Common Entrance Exam (ACET) held by the Institute of Actuaries of India (IAI).
- The student must complete an Intermediate degree or 12th final year examination.
- The minimum percentage required may vary but candidates must have scored at least 70% marks in aggregate in the compulsory subjects.

JEEWORKCUTS

Single Option Correct Type

- $f: \left(0, \frac{\pi}{2}\right) \rightarrow A$, $f(x) = \log_e(\sin x^{\sin x} + 1)$, then the minimum value of $f(x)$ is

(a) $\log_e 2$ (b) $\log_e \left(\left(\frac{1}{e}\right)^{1/e} + 1\right)$
 (c) $\log_e((e)^2 + 1)$ (d) 2
- If $x^5 - x^3 + x = a$, then minimum value of x^6 is
 (a) a (b) $2a - 1$ (c) $a + 2$ (d) $a - 3$
- If z be a complex number satisfying, $|z|^2 + 2(z + \bar{z}) + 3i(z - \bar{z}) + 4 = 0$ then complex number $z + 3 + 2i$ will lie on
 (a) circle with centre $1 - 5i$ and radius 4
 (b) circle with centre $1 + 5i$ and radius 4
 (c) circle with centre $1 + 5i$ and radius 3
 (d) circle with centre $1 - 5i$ and radius 3
- If $f(x - y) = f(x)g(y) - f(y)g(x) \forall x, y \in R$, where $f(x)$ is not identically zero, then
 (a) If $f'(0^+)$ exists then $f'(0^-)$ also exists but not equal
 (b) If $f'(0^+)$ exists then $f'(0^-)$ does not exist
 (c) If $f'(0^+)$ exists then it is equal to $f'(0^-)$
 (d) none of these
- A quadratic polynomial maps from $[-2, 3]$ onto $[0, 3]$ and touches x -axis at $x = 3$, then the polynomial is
 (a) $\frac{3}{16}(x^2 - 6x + 16)$ (b) $\frac{3}{25}(x^2 - 6x + 9)$
 (c) $\frac{3}{25}(x^2 - 6x + 16)$ (d) None of these
- Let $f(x) = \int_0^x (t^2 + 2t + 2) dt$, where x belongs to the set of real numbers satisfying the inequation $\log_2(1 + \sqrt{6x - x^2 - 8}) \geq 0$. If $f(x) \in [\alpha, \beta]$, then maximum value of $|\alpha - \beta|$ is
 (a) $103/6$ (b) $104/3$
 (c) $106/3$ (d) None of these
- The set of values of a for which inequation $(a - 1)x^2 - (a + 1)x + a - 1 \geq 0$ is true for all $x \geq 2$ is
 (a) $\left(1, \frac{7}{3}\right]$ (b) $(-\infty, 1)$
 (c) $\left[\frac{7}{3}, \infty\right)$ (d) None of these
- If $x_1, x_2, x_3, \dots, x_n$ are the roots of $x^n + ax + b = 0$, then the value of $(x_1 - x_2)(x_1 - x_3)(x_1 - x_4) \dots (x_1 - x_n)$ is
 (a) n_1^n (b) $nx_1^{n-1} + a$
 (c) nx_1 (d) $nx_1 + a^n$
- The value of $\lim_{n \rightarrow \infty} \left(\frac{n!}{(mn)^n}\right)^{1/n}$ is
 (a) em (b) e/m
 (c) $1/em$ (d) None of these
- If one of the roots of the equation $\begin{vmatrix} 7 & 6 & x^2 - 13 \\ 2 & x^2 - 13 & 2 \\ x^2 - 13 & 3 & 7 \end{vmatrix} = 0$ is $x = 2$, then sum of all other five roots is
 (a) $2\sqrt{15}$ (b) -2
 (c) $\sqrt{20} + \sqrt{15} - 2$ (d) None of these
- A unit vector is orthogonal to $5\hat{i} + 2\hat{j} + 6\hat{k}$ and is coplanar to $2\hat{i} + \hat{j} + \hat{k}$ and $\hat{i} - \hat{j} + \hat{k}$, then the vector is
 (a) $\frac{3\hat{j} - \hat{k}}{\sqrt{10}}$ (b) $\frac{2\hat{i} + 5\hat{j}}{\sqrt{29}}$

$$(c) \frac{6\hat{i} - 5\hat{k}}{\sqrt{61}}$$

$$(d) \frac{2\hat{i} + 2\hat{j} - \hat{k}}{3}$$

12. The range of the function $y = \sqrt{2\{x\} - \{x\}^2 - \frac{3}{4}}$ is (where $\{ \cdot \}$ denotes fractional part)

$$(a) \left[-\frac{1}{4}, \frac{1}{4} \right]$$

$$(b) \left[0, \frac{1}{2} \right)$$

$$(c) \left[0, \frac{1}{4} \right]$$

$$(d) \left[\frac{1}{4}, \frac{1}{2} \right]$$

13. $\int \frac{dx}{\sqrt{\cos^3 x \cos(x - \alpha)}}$ is equal to

$$(a) 2\sqrt{\tan x \sin \alpha + \cot \alpha + c}$$

$$(b) 2\operatorname{cosec} \alpha \sqrt{\tan x \sin \alpha + \cos \alpha + c}$$

$$(c) 2\sqrt{\sin \alpha + \tan x \cos \alpha + c}$$

$$(d) 2\operatorname{cosec} \alpha \sqrt{\tan x \cos \alpha + \sin \alpha + c}$$

14. Inside a big circle exactly n small circles each of radius r can be drawn in such a way that each small circle touches the big circle and two small circles. If $n \geq 3$, then the radius of the bigger circle is

$$(a) r \operatorname{cosec} \left(\frac{\pi}{n} \right)$$

$$(b) r \left\{ 1 + \operatorname{cosec} \left(\frac{\pi}{n} \right) \right\}$$

$$(c) r \left\{ 1 + \operatorname{cosec} \left(\frac{2\pi}{n} \right) \right\}$$

$$(d) r \left\{ 1 + \operatorname{cosec} \left(\frac{\pi}{2n} \right) \right\}$$

15. If m_1 and m_2 be the slopes of two perpendicular chords of equal length passing through origin of circle $(x - 1)^2 + (y + 2)^2 = 5$, then the value of $m_1^2 + m_2^2$ is equal to

$$(a) 80/9$$

$$(b) 82/9$$

$$(c) 83/9$$

$$(d) \text{None of these}$$

One or More Than One Option(s) Correct Type

16. Let $f: [a, b] \rightarrow [1, \infty)$ be a continuous function and let $g: \mathbb{R} \rightarrow \mathbb{R}$ be defined as

$$g(x) = \begin{cases} 0 & , \text{ if } x < a \\ \int_a^x f(t) dt, & \text{ if } a \leq x \leq b \\ \int_a^b f(t) dt, & \text{ if } x > b \end{cases}$$

Then

$$(a) g(x) \text{ is continuous but not differentiable at } a$$

$$(b) g(x) \text{ is differentiable on } \mathbb{R}$$

$$(c) g(x) \text{ is continuous but not differentiable at } b$$

$$(d) g(x) \text{ is continuous and differentiable at either } a \text{ or } b \text{ but not both}$$

17. Let M and N be two 3×3 matrices such that $MN = NM$. Further, if $M \neq N^2$ and $M^2 = N^4$, then

$$(a) \text{determinant of } (M^2 + MN^2) \text{ is } 0$$

$$(b) \text{there is a } 3 \times 3 \text{ non-zero matrix } U \text{ such that } (M^2 + MN^2)U \text{ is the zero matrix}$$

$$(c) \text{determinant of } (M^2 + MN^2) \geq 1$$

$$(d) \text{for a } 3 \times 3 \text{ matrix } U, \text{ if } (M^2 + MN^2)U \text{ equals the zero matrix then } U \text{ is the zero matrix.}$$

18. An ellipse intersects the hyperbola $2x^2 - 2y^2 = 1$ orthogonally. The eccentricity of the ellipse is reciprocal of that of the hyperbola. If the axes of the ellipse are along the coordinate axes, then

$$(a) \text{Equation of ellipse is } x^2 + 2y^2 = 2$$

$$(b) \text{The foci of ellipse are } (\pm 1, 0)$$

$$(c) \text{Equation of ellipse is } x^2 + 2y^2 = 4$$

$$(d) \text{The foci of ellipse are } (\pm\sqrt{2}, 0)$$

19. If $I_n = \int_{-\pi}^{\pi} \frac{\sin nx}{(1 + \pi^x) \sin x} dx$, $n = 0, 1, 2, \dots$, then

$$(a) I_n = I_{n+2}$$

$$(b) \sum_{m=1}^{10} I_{2m+1} = 10\pi$$

$$(c) \sum_{m=1}^{10} I_{2m} = 0$$

$$(d) I_n = I_{n+1}$$

20. Let $\vec{x}$, $\vec{y}$ and $\vec{z}$ be three vectors each of magnitude $\sqrt{2}$ and the angle between each pair of them is $\pi/3$. If $\vec{a}$ is a non-zero vector perpendicular to $\vec{x}$ and $\vec{y} \times \vec{z}$ and $\vec{b}$ is non-zero vector perpendicular to $\vec{y}$ and $\vec{z} \times \vec{x}$, then

$$(a) \vec{b} = (\vec{b} \cdot \vec{z})(\vec{z} - \vec{x}) \quad (b) \vec{a} = (\vec{a} \cdot \vec{y})(\vec{y} - \vec{z})$$

$$(c) \vec{a} \cdot \vec{b} = -(\vec{a} \cdot \vec{y})(\vec{b} \cdot \vec{z}) \quad (d) \vec{a} = (\vec{a} \cdot \vec{y})(\vec{z} - \vec{y})$$

21. Let z_1 and z_2 be two distinct complex numbers and let $z = (1 - t)z_1 + tz_2$ for some real number t with $0 < t < 1$. If $\operatorname{Arg}(\omega)$ denotes the principal argument of a non-zero complex number ω , then

$$(a) |z - z_1| + |z - z_2| = |z_1 - z_2|$$

$$(b) \operatorname{Arg}(z - z_1) = \operatorname{Arg}(z - z_2)$$

$$(c) \begin{vmatrix} z - z_1 & \bar{z} - \bar{z}_1 \\ z_2 - z_1 & \bar{z}_2 - \bar{z}_1 \end{vmatrix} = 0$$

$$(d) \operatorname{Arg}(z - z_1) = \operatorname{Arg}(z_2 - z_1)$$

22. If $\sin y = x \cos(a + y)$, then dy/dx is

$$(a) \frac{\cos^2(a + y)}{\cos a}$$

$$(b) \frac{\cos(a + y)}{x \sin(a + y) + \cos y}$$

$$(c) \frac{\cos a}{3x^2 - 2x \cos a - 1}$$

$$(d) \text{none of these}$$

23. The $\det \Delta = \begin{vmatrix} d^2 + r & de & df \\ de & e^2 + r & ef \\ df & ef & f^2 + r \end{vmatrix}$ is divisible by

- (a) r (b) $(d + e^2 + f^2 + r)$
(c) $(d^2 + e^2 + f^2 + r)$ (d) $(d^2 + e + f^2 + r^2)$

24. If both the roots of equation $ax^2 + x + c - a = 0$ are imaginary and $c > -1$, then

- (a) $3a > 2 + 4c$ (b) $a > 0$
(c) $c < a$ (d) None of these

25. The solution of the equation

$$3^{\sin 2x + 2 \cos^2 x} + 3^{1 - \sin 2x + 2 \sin^2 x} = 28 \text{ is}$$

- (a) $(2n+1)\frac{\pi}{2}, n \in I$ (b) $n\pi - \frac{\pi}{4}, n \in I$
(c) $2n\pi$ (d) none of these

Numerical Answer Type

26. Let m and n be two positive integers greater than 1.

If $\lim_{\alpha \rightarrow 0} \left(\frac{e^{\cos(\alpha^n)} - e}{\alpha^m} \right) = -\left(\frac{e}{2}\right)$, then the value of $\frac{m}{n}$ is _____.

27. Let X be the set consisting of the first 2018 terms of the arithmetic progression 1, 6, 11, ... , and Y be the set consisting of the first 2018 terms of the arithmetic progression 9, 16, 23, ... Then, the number of elements in the set $X \cup Y$ is _____.

28. A farmer F_1 has a land in the shape of a triangle with vertices at $P(0, 0)$, $Q(1, 1)$ and $R(2, 0)$. From this land, a neighbouring farmer F_2 takes away the region which lies between the side PQ and a curve of the form $y = x^n$ ($n > 1$). If the area of the region taken away by the farmer F_2 is exactly 30% of the area of ΔPQR , then the value of n is _____.

29. A debate club consists of 6 girls and 4 boys. A team of 4 members is to be selected from this club including the selection of a captain (from among these 4 members) for the team. If the team has to include at most one boy, then the number of ways of selecting the team is _____.

30. In ΔABC if BC is unity,

$$\sin \frac{A}{2} = x_1, \sin \frac{B}{2} = x_2, \cos \frac{A}{2} = x_3 \text{ and } \cos \frac{B}{2} = x_4$$

with $\left(\frac{x_1}{x_2}\right)^{2007} - \left(\frac{x_3}{x_4}\right)^{2006} = 0$, then the length of AC is _____.

SOLUTIONS

1. (b): Let $h(x) = \log(x^x + 1)$, for $0 < x < 1$. Then, $h'(x) > 0$ if $x > \frac{1}{e}$. Thus, for $\frac{1}{e} < x < 1$, function will be increasing. Also, $h'(x) < 0$, if $x < \frac{1}{e}$. So, at $x = \frac{1}{e}$, there is minimum. $f_{\min} = \log_e \left(\left(\frac{1}{e}\right)^{1/e} + 1 \right)$.

2. (b): $x(x^4 - x^2 + 1) = \frac{x(x^6 + 1)}{x^2 + 1}$. Suppose $x > 0$, then $a > 0$ and $x^6 + 1 = \frac{a(x^2 + 1)}{x} = a \left(x + \frac{1}{x} \right) \geq 2a$, $x^6 > 2a - 1$.

If $x \leq 0 \Rightarrow a \leq 0$, $2a - 1 < 0 \leq x^6 \Rightarrow x^6 > 2a - 1$.

3. (c): Given equation is $|z|^2 + z(2 + 3i) + \bar{z}(2 - 3i) + 4 = 0$, which represents a circle with centre $-(2 - 3i)$, radius $= \sqrt{a\bar{a} - b} = 3$

Let $\omega = z + 3 + 2i = z + 2 - 3i + 1 + 5i$,

$$|\omega - 1 - 5i| = |z + 2 - 3i| = 3$$

so, ω lies on circle whose centre is $1 + 5i$, radius $= 3$.

4. (c): $f(x - y) = f(x)g(y) - f(y)g(x)$... (i)

Put $x = y$ in (i), $f(0) = 0$,

Put $y = 0$ in (i), $f(x) = f(x)g(0) - f(0)g(x) \Rightarrow g(0) = 1$

$$\text{Now, } f'(0^+) = \lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{f(0)g(-h) - f(-h)g(0)}{h} = \lim_{h \rightarrow 0} \frac{f(-h)}{-h} = f'(0^-)$$

i.e. $f'(0^+) = f'(0^-)$.

5. (b)

$$6. (b): \log_2(1 + \sqrt{6x - x^2 - 8}) \geq 0$$

$$\Rightarrow 1 + \sqrt{6x - x^2 - 8} \geq 1$$

$$\Rightarrow 6x - x^2 - 8 \geq 0 \Rightarrow x^2 - 6x + 8 \leq 0$$

$$\Rightarrow (x - 2)(x - 4) \leq 0 \Rightarrow 2 \leq x \leq 4$$

Now $f'(x) = x^2 + 2x + 2 > 0 \forall x \in R$

$\Rightarrow f(x)$ is strictly increasing in $[2, 4]$

$$f(x) = \frac{x^3}{3} + x^2 + 2x, \alpha = f(2) = \frac{8}{3} + 4 + 4 = \frac{32}{3}$$


$$\beta = f(4) = \frac{64}{3} + 16 + 8 = \frac{136}{3}, |\alpha - \beta|_{\max} = \frac{104}{3}$$

7. (c)

8. (b): Given $x^n + ax + b = 0 = (x - x_1)(x - x_2)(x - x_3) \dots (x - x_n)$

$$\Rightarrow (x - x_2)(x - x_3) \dots (x - x_n) = \frac{x^n + ax + b}{x - x_1}$$

How can history help to succeed in JEE!



₹625/-



₹625/-



₹625/-

Wouldn't you agree that previous years' test papers provide great insights into the pattern and structure of future tests. Studies corroborate this, and have shown that successful JEE aspirants begin by familiarising themselves with problems that have appeared in past JEEs, as early as 2 years in advance.

Which is why the MTG team created 45 Years Chapterwise Solutions. The most comprehensive 'real' question bank out there, complete with detailed solutions by experts. An invaluable aid in your quest for success in JEE. Visit www.mtg.in to order online. Or simply scan the QR code to check for current offers.



Scan now with your
smartphone or tablet

Application to read
QR codes required

Note: 45 Years Chapterwise Solutions are also available for each subject separately.

Available at all leading book shops throughout India. To buy online visit www.mtg.in.

For more information or for help in placing your order, call 0124-6601200 or e-mail info@mtg.in

$$\therefore (x_1 - x_2)(x_1 - x_3)(x_1 - x_4) \dots (x_1 - x_n) \\ = \lim_{x \rightarrow x_1} \left(\frac{x^n + ax + b}{x - x_1} \right) = nx_1^{n-1} + a.$$

9. (c)

10. (b): Let $x^2 - 13 = t$, then $t^3 - 67t + 126 = 0$
 $\Rightarrow t = -9, 2, 7 \Rightarrow x = \pm 2, \pm \sqrt{20}, \pm \sqrt{15}.$

11. (a): A vector orthogonal to $\vec{c}$ and coplanar with $\vec{a}$ and $\vec{b}$ is $\vec{c} \times (\vec{a} \times \vec{b})$

Here, $\vec{c} = (5, 2, 6), \vec{a} = (2, 1, 1), \vec{b} = (1, -1, 1)$

Then $\vec{c} \times (\vec{a} \times \vec{b}) = (0, 27, -9)$

$\therefore$ Unit vector is $\frac{3\hat{j} - \hat{k}}{\sqrt{10}}.$

12. (b): Set $\{x\} = t$ so that $0 \leq t < 1, y = \sqrt{-t^2 + 2t - \frac{3}{4}}$

The domain of function is $\frac{1}{2} \leq t < 1.$

$$y\left(\frac{1}{2}\right) = \sqrt{-\frac{1}{4} + 1 - \frac{3}{4}} = 0; y(1) = \sqrt{-1 + 2 - \frac{3}{4}} = \sqrt{\frac{1}{4}} = \frac{1}{2}$$

But $\frac{1}{2}$ is not attained, so the range is $\left[0, \frac{1}{2}\right).$

$$13. (b): \text{Let } I = \int \frac{dx}{\sqrt{\cos^3 x (\cos x \cos \alpha + \sin x \sin \alpha)}} \\ = \int \frac{\sec^2 x dx}{\sqrt{\cos \alpha + \tan x \sin \alpha}}, \text{ Put } \tan x \sin \alpha + \cos \alpha = t^2$$

$$\Rightarrow \sec^2 x dx = \frac{2t dt}{\sin \alpha} \therefore I = \int \frac{2t dt}{\sin \alpha \cdot t} = 2 \operatorname{cosec} \alpha \cdot t$$

$$= 2 \operatorname{cosec} \alpha \sqrt{\tan x \sin \alpha + \cos \alpha} + c$$

14. (b): We have, $\sin\left(\frac{\pi}{n}\right) = \frac{r}{R-r} \Rightarrow R-r = r \operatorname{cosec}\left(\frac{\pi}{n}\right)$

$$\therefore R = r \left\{ 1 + \operatorname{cosec}\left(\frac{\pi}{n}\right) \right\}$$

15. (b): Let slope of chord, $OA = m, \tan 45^\circ = \left| \frac{m+2}{1-2m} \right|$

$$\Rightarrow |m+2| = |1-2m| \Rightarrow m+2 = \pm (1-2m) \Rightarrow m=3 \text{ or } -\frac{1}{3}$$

$$\Rightarrow m_1^2 + m_2^2 = (m_1 + m_2)^2 - 2m_1 m_2 = \frac{64}{9} + 2 = \frac{82}{9}$$

16. (a, c): $g(a^-) = 0$

$$\text{Also, } g(a^+) = \lim_{h \rightarrow 0} \int_a^{a+h} f(t) dt = 0$$

Thus g is continuous at $x = a$. Similarly, g is continuous at $x = b$.

$$g'(a^-) = \lim_{h \rightarrow 0} \frac{g(a-h) - g(a)}{-h} \text{ (by definition)}$$

$$= \lim_{h \rightarrow 0} \frac{0 - \int_a^{a-h} f(t) dt}{-h} = \lim_{h \rightarrow 0} \frac{0}{-h} = 0 \\ g'(a^+) = \lim_{h \rightarrow 0} \frac{g(a+h) - g(a)}{h} = \lim_{h \rightarrow 0} \frac{\int_a^{a+h} f(t) dt - 0}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\int_a^{a+h} f(t) dt}{h} = \lim_{h \rightarrow 0} \frac{f(a+h)}{1} = f(a) \geq 1$$

So, $g'(a^-) \neq g'(a^+)$

Hence g is not differentiable at $x = a$.

Similarly, g is not differentiable at $x = b$.

17. (a, b): $M^2 = N^4 \Rightarrow (M - N^2)(M + N^2) = 0$
as M and N commute.

$M - N^2 \neq 0$ so $\det(M + N^2) = 0$

Recall that if $AB = O$ and $A \neq 0$, then $\det B = 0$

$$\text{Now, } \det(M^2 + MN^2) = \det(M(M + N^2)) \\ = (\det M)(\det(M + N^2)) \\ = (\det M) \cdot 0 = 0$$

Recall that if $\det A = 0$, then $\exists X \neq 0$ such that $AX = 0$

Let $M^2 + MN^2$ play the role of A and X that of U , we get $(M^2 + MN^2)U = O$ for some 3×3 non-zero matrix U .

18. (a, b): The hyperbola is $\frac{x^2}{1/2} - \frac{y^2}{1/2} = 1 \dots (i)$

Eccentricity of the hyperbola = $\sqrt{2}$

$\therefore$ Eccentricity of the ellipse = $\frac{1}{\sqrt{2}}$

Let the equation of ellipse be $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 (a > b)$
Then $a^2 e^2 = a^2 - b^2$

$$\Rightarrow a^2 \cdot \frac{1}{2} = a^2 - b^2 \Rightarrow b^2 = \frac{a^2}{2} \therefore a^2 = 2b^2$$

The equation is $\frac{x^2}{2b^2} + \frac{y^2}{b^2} = 1$

$$x^2 + 2y^2 = 2b^2 \dots (ii)$$

Let (x_0, y_0) be a point of intersection of hyperbola (i) and ellipse (ii),

$$\text{Then for hyperbola } \left[\frac{dy}{dx} \right]_{(x_0, y_0)} = \frac{x_0}{y_0}$$

$$\text{and for ellipse } \left[\frac{dy}{dx} \right]_{(x_0, y_0)} = -\frac{x_0}{2y_0}$$

As the intersection is orthogonal, we have

$$-\frac{x_0^2}{2y_0^2} = -1 \Rightarrow x_0^2 = 2y_0^2$$

From (i) and (ii), $2x_0^2 - 2y_0^2 = 1$ and $x_0^2 + 2y_0^2 = 2b^2$

Multiplying the 1st equation by $2b^2$ and subtracting we get $4b^2x_0^2 - 4y_0^2b^2 = x_0^2 + 2y_0^2$

$$\Rightarrow x_0^2(4b^2 - 1) = 2y_0^2(1 + 2b^2)$$

Also $x_0^2 = 2y_0^2$ which gives

$$4b^2 - 1 = 1 + 2b^2 \Rightarrow 2b^2 = 2$$

$$\therefore b^2 = 1$$

Thus the equation to ellipse is $x^2 + 2y^2 = 2$

$\therefore$ Foci of the ellipse are $(\pm\sqrt{a^2 - b^2}, 0)$ i.e. $(\pm 1, 0)$

$$19. (a, b, c) : I_n = \int_{-\pi}^{\pi} \frac{\sin nx}{(1 + \pi^x) \sin x} dx \quad \dots(i)$$

$$\Rightarrow I_n = \int_{-\pi}^{\pi} \frac{\sin(-nx)}{(1 + \pi^{-x})(-\sin x)} dx \quad \dots(ii)$$

Adding (i) and (ii), we get

$$2I_n = \int_{-\pi}^{\pi} \left(\frac{1}{1 + \pi^x} + \frac{1}{1 + \pi^{-x}} \right) \frac{\sin nx}{\sin x} dx = \int_{-\pi}^{\pi} \frac{\sin nx}{\sin x} dx$$

$$\Rightarrow I_n = \frac{1}{2} \int_{-\pi}^{\pi} \frac{\sin nx}{\sin x} dx$$

$$\text{Now, } I_{n+2} - I_n = \frac{1}{2} \int_{-\pi}^{\pi} \frac{\sin(n+2)x - \sin nx}{\sin x} dx$$

$$= \frac{1}{2} \int_{-\pi}^{\pi} \frac{2 \sin x \cos(n+1)x}{\sin x} dx$$

$$= \int_{-\pi}^{\pi} \cos(n+1)x dx = \left(\frac{\sin(n+1)x}{(n+1)} \right)_{-\pi}^{\pi} = 0$$

$$\therefore I_{n+2} = I_n$$

Thus $I_0 = I_2 = I_4 = \dots = I_{10}$

$$I_1 = I_3 = I_5 = \dots = I_{11}$$

$$I_0 = \frac{1}{2} \int_{-\pi}^{\pi} 0 dx = 0$$

$$\therefore I_1 = \frac{1}{2} \int_{-\pi}^{\pi} \frac{\sin x}{\sin x} dx = \frac{1}{2} \cdot 2\pi = \pi$$

$$\sum_{m=1}^{10} I_{2m} = 0 \text{ and } \sum_{m=1}^{10} I_{2m+1} = 10\pi$$

20. (a, b, c) : $\vec{a} = p(\vec{x} \times (\vec{y} \times \vec{z}))$, where $p \in \mathbb{R} - \{0\}$

$$= p\{(\vec{x} \cdot \vec{z})\vec{y} - (\vec{x} \cdot \vec{y})\vec{z}\} = p\{\vec{y} - \vec{z}\}$$

$$\left(\because \vec{x} \cdot \vec{y} = |\vec{x}| |\vec{y}| \cos \frac{\pi}{3} = \sqrt{2} \cdot \sqrt{2} \left(\frac{1}{2} \right) = 1 = \vec{y} \cdot \vec{z} = \vec{z} \cdot \vec{x} \right)$$

$$\text{Now, } \vec{a} \cdot \vec{y} = p\{\vec{y} \cdot \vec{y} - \vec{y} \cdot \vec{z}\} = p\{2 - 1\} = p$$

$$\therefore p = \vec{a} \cdot \vec{y} \quad (\because \vec{y} \cdot \vec{y} = \vec{y}^2 = (\sqrt{2})^2 = 2)$$

$$\text{Thus, } \vec{a} = (\vec{a} \cdot \vec{y})(\vec{y} - \vec{z}) \quad \dots (i)$$

$$\text{Similarly, } \vec{b} = q(\vec{y} \times (\vec{z} \times \vec{x})) = q\{(\vec{y} \cdot \vec{x})\vec{z} - (\vec{y} \cdot \vec{z})\vec{x}\} = q(\vec{z} - \vec{x})$$

$$\text{Now, } \vec{b} \cdot \vec{z} = q(\vec{z} \cdot \vec{z} - \vec{z} \cdot \vec{x}) = q(\vec{z}^2 - \vec{z} \cdot \vec{x}) = q$$


$$\text{Thus } \vec{b} = (\vec{b} \cdot \vec{z})(\vec{z} - \vec{x}) \quad \dots (ii)$$

$$\text{Now, } \vec{a} \cdot \vec{b} = (\vec{a} \cdot \vec{y})(\vec{b} \cdot \vec{z})\{(\vec{y} - \vec{z}) \cdot (\vec{z} - \vec{x})\} \quad (\text{From (i) \& (ii)})$$

$$= (\vec{a} \cdot \vec{y})(\vec{b} \cdot \vec{z})(\vec{y} \cdot \vec{z} - \vec{y} \cdot \vec{x} - \vec{z} \cdot \vec{z} + \vec{z} \cdot \vec{x})$$

$$= (\vec{a} \cdot \vec{y})(\vec{b} \cdot \vec{z})(1 - 1 - 2 + 1) = -(\vec{a} \cdot \vec{y})(\vec{b} \cdot \vec{z})$$

21. (a, c, d) : The given statement implies that z is on the line segment joining A and B.



As $z = (1 - t)z_1 + tz_2$ indicate.

$$\therefore AP + PB = AB \Rightarrow |z - z_1| + |z - z_2| = |z_1 - z_2|$$

$$\text{Now, } z_1, z, z_2 \text{ are collinear, so } \text{Arg} \left(\frac{z - z_1}{z_2 - z_1} \right) = 0$$

$$\text{giving } \text{Arg}(z - z_1) = \text{Arg}(z_2 - z_1)$$

$$\text{Note that } \text{Arg}(z - z_1) - \text{Arg}(z - z_2) = \pi$$

Form IV

1. Place of Publication	: New Delhi
2. Periodicity of its Publication	: Monthly
3. Printer's Name	: HT Media Ltd.
3a. Publisher's Name	: MTG Learning Media Pvt. Ltd.
Nationality	: Indian
Address	: 406, Taj Apartment, New Delhi - 110029
4. Editor's Name	: Anil Ahlawat
Nationality	: Indian
Address	: 19, National Media Centre, Gurugram, Haryana - 122002
5. Name and address of individuals who own the newspapers and partners or shareholders holding more than one percent of the total capital	: Mahabir Singh Ahlawat 64, National Media Centre, Nathupur, Gurugram : Krishna Devi 64, National Media Centre, Nathupur, Gurugram : Anil Ahlawat & Sons 19, National Media Centre, Nathupur, Gurugram : Anil Ahlawat 19, National Media Centre, Nathupur, Gurugram

I, Mahabir Singh, authorised signatory for MTG Learning Media Pvt. Ltd. hereby declare that particulars given above are true to the best of my knowledge and belief.

For MTG Learning Media Pvt. Ltd.
Mahabir Singh
Director

Also $\frac{z-z_1}{\bar{z}-\bar{z}_1} = \frac{z_2-z_1}{\bar{z}_2-\bar{z}_1}$ = complex slope of the line which

can be rearranged as

$$(z-z_1)(\bar{z}_2-\bar{z}_1) - (z_2-z_1)(\bar{z}-\bar{z}_1) = 0$$

$$\text{i.e., } \begin{vmatrix} z-z_1 & \bar{z}-\bar{z}_1 \\ z_2-z_1 & \bar{z}_2-\bar{z}_1 \end{vmatrix} = 0$$

$$22. (a, b) : \sin y = x \cos(a+y) \Rightarrow \frac{\sin y}{\cos(a+y)} = x \quad \dots(i)$$

$$\frac{dx}{dy} = \frac{\cos y \cos(a+y) + \sin y \sin(a+y)}{\cos^2(a+y)} = \frac{\cos(a+y)}{\cos^2(a+y)}$$

$$\Rightarrow \frac{dy}{dx} = \frac{\cos^2(a+y)}{\cos a} \quad \dots(ii)$$

$$= \frac{\cos^2(a+y)}{\cos y \cos(a+y) + \sin y \sin(a+y)} = \frac{\cos(a+y)}{\cos y + x \sin(a+y)} \quad [\text{Using (i)}]$$

23. (a, c) : Multiply R_1, R_2, R_3 by d, e, f and take out d, e, f common from C_1, C_2, C_3 , then

$$\Delta = \begin{vmatrix} d^2+r & d^2 & d^2 \\ e^2 & e^2+r & e^2 \\ f^2 & f^2 & f^2+r \end{vmatrix}$$

Operate $R_1 \rightarrow R_1 + R_2 + R_3$ and take out $d^2 + e^2 + f^2 + r$ common from R_1 , then

$$\Delta = \begin{vmatrix} 1 & 1 & 1 \\ e^2 & e^2+r & e^2 \\ f^2 & f^2 & f^2+r \end{vmatrix} (d^2 + e^2 + f^2 + r) \\ = r^2(d^2 + e^2 + f^2 + r).$$

24. (b) : Here $f(1) = a + 1 + c - a$

As $c > -1 \Rightarrow f(1) > 0$

25. (a, b) : We have,

$$3^{\sin 2x + 2 \cos^2 x} + 3^{1 - \sin 2x + 2 \sin^2 x} = 28, \quad y + \frac{27}{y} = 28,$$

$$\Rightarrow y^2 - 28y + 27 = 0 \Rightarrow y = 1$$

$$(\because y \neq 27, \text{ since } \max. (\sin 2x + \cos 2x) = \sqrt{2})$$

$$\therefore \sin 2x + 2 \cos^2 x = 0 \Rightarrow 2 \cos x (\sin x + \cos x) = 0$$

$$\Rightarrow x = (2n+1)\frac{\pi}{2} \text{ or } x = n\pi - \frac{\pi}{4}$$

$$26. (2) : \lim_{\alpha \rightarrow 0} \left(\frac{e^{\cos(\alpha^n)} - e}{\alpha^m} \right) = \lim_{\alpha \rightarrow 0} \frac{e(e^{\cos(\alpha^n)-1} - 1)}{\alpha^m}$$

$$= \lim_{\alpha \rightarrow 0} \frac{e(\alpha^n)^2}{\alpha^m} \left[\frac{e^{\cos(\alpha^n)-1} - 1}{\cos(\alpha^n) - 1} \right] \left[\frac{\cos(\alpha^n) - 1}{(\alpha^n)^2} \right] = \frac{-1}{2} e \alpha^{2n-m}$$

As limit is given to be $-e/2$, we have

$$2n - m = 0. \therefore m/n = 2.$$

27. (3748) : Here $X = \{1, 6, 11, \dots, 10086\}$

and $Y = \{9, 16, 23, \dots, 14128\}$

The intersection of X and Y is an A.P. with 16 as first term and 35 as common difference.


The series becomes 16, 51, 86, ...

Now, k^{th} term = $16 + (k-1)35 \leq 10086$

$$\text{i.e. } k \leq \frac{10105}{35} \therefore k \leq 288 \text{ (as } k \text{ is to be an integer)}$$

$$\text{Hence, } n(X \cup Y) = n(X) + n(Y) - n(X \cap Y) \\ = 2018 + 2018 - 288 = 3748$$

28. (4) :



$$\int_0^1 (x - x^n) dx = \frac{3}{10} \cdot \left(\frac{1}{2} \times 2 \times 1 \right)$$

$$\Rightarrow \left[\frac{x^2}{2} - \frac{x^{n+1}}{n+1} \right]_0^1 = \frac{3}{10} \Rightarrow \frac{1}{2} - \frac{1}{n+1} = \frac{3}{10}$$

$$\Rightarrow \frac{1}{2} - \frac{3}{10} = \frac{1}{n+1} \Rightarrow \frac{2}{10} = \frac{1}{n+1}$$

$$\Rightarrow n+1 = 5 \therefore n = 4$$

29. (380) : The team can have either 0 boy or 1 boy.

$$\text{So the number of selection is } ({}^6C_4 \cdot {}^4C_0 + {}^6C_3 \cdot {}^4C_1) \cdot {}^4C_1 \\ = (15 + 20 \cdot 4) \cdot 4 = 95 \cdot 4 = 380$$

30. (1) : In given ΔABC both $A/2$ and $B/2$ lie strictly between $(0, \pi/2)$ and $\sin x$ is always increasing in $(0, \pi/2)$ whereas $\cos x$ is always decreasing throughout $(0, \pi/2)$.

$$\text{So if } \frac{A}{2} > \frac{B}{2} \Rightarrow \sin \frac{A}{2} > \sin \frac{B}{2} \Rightarrow x_1 > x_2$$

$$\text{and } \frac{1}{x_3} > \frac{1}{x_4}. \text{ Now, } x_1^{2007} x_4^{2006} = x_2^{2007} x_3^{2006} \text{ is not valid.}$$

Similarly for $A/2 < B/2$

$$\sin \frac{A}{2} < \sin \frac{B}{2} \Rightarrow x_1 < x_2 \text{ and } \frac{1}{x_3} < \frac{1}{x_4}. \text{ Again equality is not possible.}$$

Therefore only possible case is when $A/2 = B/2$

$$\Rightarrow x_1 = x_2 \text{ and } \frac{1}{x_3} = \frac{1}{x_4}.$$

Hence in this case ΔABC is isosceles with $\angle ABC = \angle CAB$





QUANTITATIVE APTITUDE

For Various Competitive Exams

- $(1+2)(1+2^2)(1+2^4)(1+2^8)(1+2^{16})(1+2^{32})(1+2^{64})(1+2^{128}) = ?$
 (a) $2^{512} - 1$ (b) $2^{1024} - 1$
 (c) $2^{255} - 1$ (d) $2^{256} - 1$
- A college has raised 75% of the amount it needs for a new building by receiving an average donation of ₹ 600 from the people already solicited. The people already solicited represent 60% of the people the college will ask for donations. If the college is to raise exactly 25% of the amount needed for the new building, what should be the average donation from the remaining people to be solicited?
 (a) ₹ 300 (b) ₹ 250
 (c) ₹ 400 (d) ₹ 500
- Mr. 'X' invested certain amounts in two different schemes 'A' & 'B' for two years. Scheme 'A' offers simple interest at 12 % p.a. and Scheme 'B' offers compound interest at 10 % per annum. Interest accrued on the amount invested in Scheme 'A' in 2 years was ₹ 3,600 and the total amount invested was ₹ 35,000. What was interest accrued on the amount invested in Scheme 'B' ?
 (a) ₹ 4,800 (b) ₹ 4,200
 (c) ₹ 4,000 (d) Cannot be determined
- Given a number 12345678901234567890..... upto 500 digits. Find the smallest number n that should be added to the number such that the sum is exactly divisible by 11.
 (a) 6 (b) 5 (c) 8 (d) 1
- A certain manufacture sells a product to the distributor at 10% profit. Then distributor sells it to the dealer and the dealer sells to the retailer at a mark up of 10% and 20% respectively. The retailer marks up his cost by 20% and then offers a 10% discount to the customers. If the customer had bought it from distributor directly, then how much reduction in price would he get with respect to buying it from the retailer?
 (a) 29.6% (b) 20% (c) 33.33% (d) 22.8%
- Mr. X invested a certain amount in Debit and Equity funds in the ratio of 4 : 5 respectively. At the end of one year, he earned a total dividend of 30% on his investment. After one year he reinvested the amount including dividend in the ratio of 6 : 7 in Debit and Equity Funds. If the amount reinvested in Equity Funds was ₹ 94,500, what was the original amount invested in Equity Funds?
 (a) ₹ 75,000 (b) ₹ 81,000
 (c) ₹ 60,000 (d) ₹ 65,000
- Professor John noticed something peculiar while entering the quiz marks of his five students into a spreadsheet. The after each score was entered, Professor John entered the marks in a random order and noticed that after each mark was entered, the average was always an integer. In ascending order, the marks of the students were 71, 76, 80, 82 and 91. What were the fourth and fifth marks that Professor John entered?
 (a) 71 and 82 (b) 71 and 76
 (c) 71 and 80 (d) 76 and 80
- The diluted wine contains only 6 litres of wine and the rest is water. A new mixture whose concentration is 50% is to be formed by replacing with wine. How many litres of mixture be replaced with pure wine if there are initially 18 litres of water in the mixture?
 (a) 5 litres (b) 6 litres
 (c) 8 litres (d) 12 litres
- If 15 men or 24 women or 36 boys can do a piece of work in 12 days, working 8 hours a day, how many men must be associated with 12 women and 6 boys to do another piece of work $2\frac{1}{4}$ times as great in 30 days working 6 hours a day?
 (a) 4 (b) 8 (c) 6 (d) 10
- Navjivan Express from Ahmedabad to Chennai leaves Ahmedabad at 6:30 a.m. and travels at 50 kmph towards Baroda situated 100 km away. At 7:00 a.m.

Howrah-Ahmedabad Express leaves Baroda towards Ahmedabad and travels at 40 kmph. At 7:30 a.m., Mr. Shah, the traffic controller at Baroda realizes that both the trains are running on the same track. How much time does he have to avert a head-on collision between the two trains?

- (a) 20 minutes (b) 30 minutes
(c) 45 minutes (d) None of these


11. The 288th term of the series $a, b, b, c, c, c, d, d, d, d, e, e, e, e, e, f, f, f, f, f, f, \dots$ is

- (a) u (b) v (c) w (d) x

12. Rahim sells two types of coffees viz. DC and VC. He sells DC at ₹ 18 per kg and incurs a loss of 10% whereas on selling the VC at ₹ 30 per kg, he gains 20%. In what proportion should the DC and VC be mixed such that he can gain a profit of 25% by selling the mixture at ₹ 27.5 per kg?

- (a) 5 : 3 (b) 3 : 2 (c) 2 : 3 (d) 7 : 4

13. Three identical balls fit snugly into a cylindrical can. The radius of each of the spheres equals the radius of the can and the balls just touch the bottom and the top of the can. What fraction of the volume of the can is taken up by the balls?



- (a) $3/5$ (b) $4/3$ (c) $3/2$ (d) $2/3$

14. How many four letters words can be formed using the letters of 'SERIES'?

- (a) 24 (b) 42 (c) 84 (d) 102

15. From amongst the available 15 cricket players, of whom 5 are bowlers, a team of 11 is to be selected. What is the probability that the selected team will have at least 3 bowlers?

- (a) $7/13$ (b) $5/13$ (c) $12/13$ (d) $9/13$

16. A water tank has three taps A, B and C. A fills 8 glasses in 48 minutes, B fills 16 glasses in 2 hour, and C fills 4 glasses in 40 minutes. If all the taps are opened together, a full tank is emptied in 4 hours. If a glass contains 500 ml water, what is the capacity of the tank?

- (a) 32 litres (b) 48 litres (c) 56 litres (d) 64 litres

17. The speed of the boat in still water is 16 kmph and speed of the current is 2 kmph. It takes a total of 6.5 hours to row upstream from point A to point B and downstream from point B to point C. If the distance from point A to point B is two-third the distance between point B and C, what is the total distance travelled by the boat (both upstream and downstream)?

- (a) 112 km (b) 98 km (c) 124 km (d) 105 km

18. Left pan of a faulty balance weighs 100 gram more than its right pan. A shopkeeper keeps the weight measure in the left pan while buying goods but keeps it in the right pan while selling his goods. He uses only 1 kg weight measure. If he sells his goods at the listed cost price, what is his gain?

- (a) $\frac{200}{9}\%$ (b) $\frac{100}{11}\%$ (c) $\frac{100}{9}\%$ (d) $\frac{200}{11}\%$

19. ₹ 1500 were invested for 5 years in scheme A which offers simple interest at the rate of 14% per annum. The amount received after 5 years and some additional money, is then invested in scheme B, for 2 years, which offers compound interest (compounded annually) at the rate of 20% per annum. If the compound interest received from scheme B after 2 years is ₹ 1408, what was the additional money invested in scheme B apart from the amount received from scheme A?

- (a) ₹ 450 (b) ₹ 650 (c) ₹ 500 (d) ₹ 280

20. In 2013, the arithmetic mean of the annual incomes of Ram and Shyam was ₹ 3800. The arithmetic mean of the annual incomes of Shyam and Pratap was ₹ 4800, and the arithmetic mean of the annual incomes of Pratap and Ram was ₹ 5800. What is the arithmetic mean of the incomes of the three?

- (a) ₹ 4800 (b) ₹ 5600 (c) ₹ 4200 (d) ₹ 5000

SOLUTIONS

1. (d): $(2 + 1) = 3 = 2^2 - 1$

$$(2 + 1)(2^2 + 1) = 3 \times 5 = 15 = 2^4 - 1$$

$$(2 + 1)(2^2 + 1)(2^4 + 1) = 3 \times 5 \times 17 = 2^8 - 1$$

$$\text{So, } (2 + 1)(2^2 + 1)(2^4 + 1) \dots (2^{128} + 1) = 2^{256} - 1$$

2. (a): Let x be the total number of people the college will ask for donations.

$$\therefore \text{People already solicited} = 0.6x$$

$$\text{Amount raised from the people solicited}$$

$$= 600 \times 0.6x = 360x$$

Now, $360x$ constitutes 75% of the amount.

$$\text{Hence, remaining } 25\% = 120x$$

$$\therefore \text{Average donation from remaining people}$$

$$= \frac{120x}{0.4x} = ₹ 300$$

3. (b): Scheme A : $R = 12\%$, $t = 2$ year, $I = 3600$

$$3600 = \frac{A \times 12 \times 2}{100}$$

$$A = 15,000$$

$$\text{Amount invested in } B = 35000 - 15000 = ₹ 20000$$

$$\text{Scheme B : } R = 10\%, P = 20,000$$

$$CI = (1.1)^2 \times 20000 - 20000 = 20000 (1.21 - 1)$$

$$= 20000 \times \frac{21}{100} = 4200$$

$$CI = 4,200$$

So, Interest accrued on scheme B is ₹ 4200.

4. (c) : If A is the sum of all the alternate digits of the number starting from the units place and B is the sum of all the alternate digits of the number starting from tens place, then remainder by 11 is given by the number $A - B$.

For one group of ten digits of the number, $A - B = -5$

For the 50 groups, it is -250 .

So, when this number is divided by 11, the remainder will be -8 or $+3$.

Therefore 8 has to be added to the number, so that sum is divisible by 11.

5. (d) : Let manufacturer C.P = ₹ 100

Then, his S.P would be $= 1.1 \times 100 = ₹ 110$. It is distributor's C.P.

Then, distributor S.P = $1.1 \times 110 = ₹ 121$.

Dealer S.P = $1.2 \times 121 = ₹ 145.2$.

Retailer S.P = $1.2 \times 145.2 = ₹ 174.24$.

After discount $= 0.9 \times 174.24 = ₹ 156.816$.

If he had bought from distributor he would save $= 35.816$

$$\text{Percentage reduction} = \frac{35.816}{156.816} \times 100 = 22.8\%$$

6. (a) : Amount reinvested in Debit and Equity is in the ratio 6 : 7.

Let the amount invested in Equity be $7x$ and amount invested in Debit will be $6x$.

$$\text{Now, } 7x = ₹ 94500 \Rightarrow x = 13500$$

$$\text{Amount reinvested in Debit} = 6 \times 13500 = ₹ 81000$$

$$\text{Total amount reinvested} = 94500 + 81000 = ₹ 175500.$$

As this is the original amount + 30% dividend on the original amount.

Let original amount be ₹ y

$$\Rightarrow y + \frac{30}{100} \times y = ₹ 175500 \Rightarrow 1.3y = 175500$$

$$\Rightarrow y = 135000$$

Hence, total amount invested is ₹ 135000.

$$\therefore \text{Amount invested in Equity is } \frac{5}{9} \times 135000 = ₹ 75000$$

Hence amount invested in Equity is ₹ 75000.

7. (c) : The marks of the 5 student and the remainder that they leave when divided by 3 are tabulated below.

Marks	71	76	80	82	91
Remainder	2	1	2	1	1

After the first 3 marks are entered, the total has to be a multiple of 3. The remainders can only be 1, 1, 1. (2, 1, 1 or 2, 2, 1 would not produce a multiple of 3).

So, the fourth and fifth marks to be entered were 71 and 80.

8. (c) : Wine Water


$$6 L \quad 18 L$$

$$1 : 3$$

$$25\% \quad 75\% \quad (\text{Original ratio})$$

$$50\% \quad 50\% \quad (\text{Required ratio})$$

PUZZLE CORNER



MATHDOKU

Introducing MATHDOKU, a mixture of ken-ken, sudoku and Mathematics. In this puzzle 6×6 grid is given, your objective is to fill the digits 1-6 so that each appear exactly once in each row and each column.

Notice that most boxes are part of a cluster. In the upper-left corner of each multibox cluster is a value that is combined using a specified operation on its numbers. For example, if that value is 3 for a two-box cluster and operation is multiply, you know that only 1 and 3 can go in there. But it is your job to determine which number goes where! A few cluster may have just one box and that is the number that fills that box.

3-		1-	4-	2/	
6*	2-			2/	
		2-		7+	1-
7+		3+	6		
11+			2-		1-
3/		15+			

Readers can send their responses at editor@mtg.in or post us with complete address. Winners' name with their valuable feedback will be published in next issue.

In this case, the percentage of water is being reduced when the mixture is being replaced with wine.

So, the ratio of left quantity to the initial quantity is 2 : 3.

$$\text{Therefore, } \frac{2}{3} = \left[1 - \frac{k}{24}\right] \Rightarrow \frac{k}{24} = \frac{1}{3} \Rightarrow k = 8 \text{ litres}$$

$$\mathbf{9. (b):} 15m = 24w = 36b \quad \dots(1)$$

$$\text{Total work} = 15 \times 12 \times 8 = 1440 \text{ men hours}$$

$$\text{New work} = \frac{9}{4} \times 1440 = 3240 \text{ men hours.}$$

$$\text{Now, } x \text{ men} + 12 \text{ women} + 6 \text{ boys}$$

$$= x \text{ men} + \frac{15}{2} \text{ men} + \frac{15}{6} \text{ men} \quad [\text{Using (1)}]$$

$$= \frac{6x + 60}{6} \text{ men are working on 3240 units.}$$

$$\text{Number of men} = \frac{3240}{6 \times 30} \Rightarrow \frac{6x + 60}{6} = 18$$

$$\Rightarrow 6x = 108 - 60 \Rightarrow x = 8 \text{ men}$$


10. (a):



From 6:30 to 7:00 *i.e.* for half hour, the train from Ahmedabad will move for half hour and will cover a

$$\text{distance of } 50 \times \frac{1}{2} \text{ i.e. } 25 \text{ km.}$$

Now, past 7:00 a.m. both the trains will move.



Now, both the trains are moving in the opposite direction.

So, they will collide after

$$= \frac{75}{50 + 40} = \frac{75}{90} = \frac{5}{6} \text{ hr or 50 minutes}$$

So, they will collide 50 minutes after 7:00 a.m. *i.e.* at 7:50 a.m.

Mr. Shah came to know at 7:30 that both the trains are on the same track, so he has 20 minutes to avoid the collision between the two trains.

11. (d): The number of terms of the series forms the sum of first n natural numbers *i.e.* $\frac{n(n+1)}{2}$.

Thus the first 23 letters will account for the first

$$= \frac{23 \times 24}{2} = 276 \text{ terms of the series.}$$

The 288th term will be the 24th letter which is x .

12. (b): The S.P. of DC = ₹ 18 per kg

The S.P. of VC = ₹ 30 per kg

The C.P. of DC = ₹ 20 per kg

The C.P. of VC = ₹ 25 per kg

The S.P. of mixture = ₹ 27.5 per kg

The C.P. of mixture = ₹ 22 per kg

$$22 = 20x + 25(1-x) \Rightarrow x = \frac{3}{5} \Rightarrow 1-x = \frac{2}{5}$$

Therefore, the ratio of DC is to VC is 3 : 2.

13. (d): Assume that the radii of each of the spheres and

the can are 1, then the volume of each ball is $\frac{4}{3}\pi(1)^3 = \frac{4}{3}\pi$ and the total volume of 3 balls is $3\left(\frac{4}{3}\pi\right) = 4\pi$. Since the

height of the can is 6 (the diameter of each sphere is 2), the volume of the can is $\pi(1)^2(6) = 6\pi$

So, the balls take up $\frac{4\pi}{6\pi} = \frac{2}{3}$ of the can.

14. (d): The given word SERIES, this word contains 2S, 2E and rest are distinct.

Case 1: 4 Letters are distinct

Using S, E, R, I

$$\text{Number of ways} = 4! = 24$$

Case 2: 2 letters are same and rest are distinct

Using SS or EE and using two letter out of remaining R and I.

$$\text{Ways} = {}^2C_1 \times {}^3C_2 \times \frac{4!}{2!} = 72 \text{ ways}$$

Case 3: SS and EE

$$\text{Number of ways} = \frac{4!}{2!2!} = 6 \text{ ways}$$

$$\text{Total number of ways} = 24 + 72 + 6 = 102$$

15. (c): Total players = 15

Total number of selected players = 11

$$\therefore \text{Total possible outcomes} = {}^{15}C_{11} = 1365$$

Favourable number of cases

$$= n(3 \text{ bowlers}) + n(4 \text{ bowlers}) + E(5 \text{ bowlers})$$

($\because$ Total number of bowlers = 5)

$$= ({}^5C_3 \times {}^{10}C_8) + ({}^5C_4 \times {}^{10}C_7) + ({}^5C_5 \times {}^{10}C_6)$$

$$= (10 \times 45) + (5 \times 120) + (1 \times 210) = 1260$$

$$\therefore P(\text{at least 3 bowlers}) = \frac{1260}{1365} = \frac{12}{13}$$

16. (b) : A = 8 glasses in 48 minutes

B = 16 glasses in 2 hours

C = 4 glasses in 40 minutes

In 4 hours, $A = \frac{4 \times 60}{48} \times 8 = 40$ glasses


$B = \frac{4 \times 60}{120} \times 16 = 32$ glasses

$C = \frac{4 \times 60}{40} \times 4 = 24$ glasses

Total water = 40 + 32 + 24 = 96 glasses.

Capacity = 96 × 0.5 L = 48 litres

17. (d) :



Let distance between B and C be x , $AB = \frac{2}{3}x$

$\therefore AC = \frac{1}{3}x$

Speed of boat upstream = 16 – 2 = 14 km/h

Speed of boat downstream = 16 + 2 = 18 km/h

$$\frac{2x}{3 \times 14} + \frac{x}{18} = \frac{13}{2}; \frac{1}{2} \left[\frac{2x}{3 \times 7} + \frac{x}{9} \right] = \frac{13}{2}; \frac{18x + 21x}{3 \times 7 \times 9} = 13$$

$39x = 3 \times 7 \times 9 \times 13$; $x = 63$ km

$\therefore$ Total distance travelled

$$= \frac{2}{3}x + x = \frac{5}{3}x = \frac{5}{3} \times 63 = 105 \text{ km.}$$

18. (a) : Let the price of 1 g be ₹ 1.

The shopkeeper buys 1100 g at the rate of ₹ 1000.

He sells 900 g at the cost price of ₹ 1000.

So, CP for 900 g would be $= \frac{1000}{1100} \times 900 = ₹ 818.18$

So, gain % $= \frac{1000 - 818.18}{818.18} \times 100 = 22.22\% = \frac{200}{9}\%$

19. (b) : Compound interest received after 2 year = ₹ 1408

Let the principal invested in scheme B is x at rate = 20%

Now, $1408 = x[(1.2)^2 - 1]$

$$1408 = x[1.44 - 1]; x = \frac{1408}{0.44} = 3200$$

Amount invested in scheme B is ₹ 3200.

Now, interest received in scheme A = $1500 \times \frac{5}{100} \times 14$
 $= 70 \times 15 = ₹ 1050$

Amount received from scheme A = 1500 + 1050 = ₹ 2550

Now, additional amount that is added is

$$3200 - 2550 = ₹ 650$$

20. (a) : Let a , b , and c be the annual incomes of Ram, Shyam, and Pratap, respectively.

Now, we have given that

The arithmetic mean of the annual incomes of Ram and

Shyam was ₹ 3800. Hence, $\left(\frac{a+b}{2} \right) = 3800$

Multiplying by 2 yields $a + b = 2 \times 3800 = 7600$.

The arithmetic mean of the annual incomes of Shyam and Pratap was ₹ 4800.

Hence, $\left(\frac{b+c}{2} \right) = 4800$

Multiplying by 2 yields $b + c = 2 \times 4800 = 9600$.

The arithmetic mean of the annual incomes of Pratap and Ram was ₹ 5800.

Hence, $\left(\frac{c+a}{2} \right) = 5800$

Multiplying by 2 yields $c + a = 2 \times 5800 = 11,600$

Summing these three equations yields:

$$(a+b) + (b+c) + (c+a) = 7600 + 9600 + 11600$$

$$2a + 2b + 2c = 28,800; a + b + c = 14,400$$

The average of the incomes of the three

$$= \left(\frac{a+b+c}{3} \right) = \frac{14400}{3} = ₹ 4800$$



PUZZLE CORNER

ANSWER • FEBRUARY 2023

$2 \div$ 2	1	$3 \div$ 6	$2 \div$ 3	$8 \div$ 4	$2 -$ 5
$1 -$ 5	4	2	6	1	3
$3 -$ 4	6 6	$40 \times$ 5	2	3	$9 \div$ 1
1	$2 -$ 5	3	4	2	6
$54 \times$ 6	3	$3 -$ 4	1	$3 -$ 5	2
3	$1 -$ 2	1	$15 \div$ 5	6	4

Brain Teaser

LOGICAL REASONING

for Various Competitive Exams

Direction Q.(1 and 2) : What will come in place of '?' in the given series?

1. 9, 31, 73, 141, ?
(a) 164 (b) 280 (c) 239 (d) 241
2. A2E, C5H, E12K, G27N, ?
(a) I58Q (b) I77R (c) J58R (d) J57Q

Direction Q.(3 and 4) : The two words or numbers on the right of (:) are related in the same way as the words on the left of (:). Identify what will come in place of blank.


3. GRIN : RING :: : NECKLACE
(a) NEECKLAC (b) LACENECK
(c) ENECKLAC (d) NECKLAEC
4. 216 : 81 :: 512 :
(a) 100 (b) 64 (c) 48 (d) 256
5. In a code language, 'FUN' is coded as 'HWP', then 'TCFKQ' is code of
(a) REALER (b) RIDER
(c) RADIO (d) ROMAN
6. If LUXOR is coded as 30, then GUILDS will be coded as
(a) 40 (b) 36 (c) 38 (d) 24

Direction Q.(7 and 8) : Three of the following four are alike in a certain way and so form a group. Which is the one that does not belong to that group?

7.
(a) Bagel (b) Bun (c) Partly (d) Damper
8.
(a) Cuckoo (b) Crow (c) Bat (d) Parrot

Direction Q.(9 and 10) : In each of the following figures, find the number which replace the sign of '?'

9.



- (a) -173 (b) 173 (c) 713 (d) 143

- 10.
-
- (a) 5 (b) 2 (c) 7 (d) 9

Direction Q.(11 and 12) : Study the following information carefully and answer the questions given below:

P is son of Q. Q is mother of R. R is wife of T. T is father of V. V is brother of W. Y is mother of T.

11. Which of the following is true based upon the relationships given above?
(a) W is granddaughter of Y
(b) R is sister of P
(c) V is son of Q
(d) V is brother-in-law of Y
12. Which of the following is/are required to establish that W is the daughter of R?
(a) R has only three children, one son and two daughters
(b) Q has only one grandson
(c) Y has only two children, a son and a daughter
(d) Either (a) or (b)

13. Bimla left her house to go to a temple. She walked 10 m straight to north and then took a left to walk 3 m and then a right to walk 5 m and then a left to walk 2 m and then a right to walk 5 m and then again a right to walk 30 m to reach the temple. How far is the temple and in which direction from Bimla's house?

- (a) 25 m North (b) $15\sqrt{2}$ m North-East
(c) $5\sqrt{41}$ m North-East (d) $15\sqrt{2}$ m North

Direction Q.(14 and 15) : In these questions, relationship between different elements is shown in the statements. The statements are followed by two conclusions.

Mark answer

- (a) if only conclusion (I) is true.
(b) if only conclusion (II) is true.
(c) if neither conclusion (I) nor (II) is true.
(d) if both conclusions (I) and (II) are true.

14. Statements : $G \leq K = O \geq U > P < S$

Conclusions :

I. $G \leq U$ II. $S > O$

15. Statements: $O = L \leq T > S$; $L > I$; $T \leq Z$

Conclusions :

I. $Z > I$ II. $S < I$

16. Vinay remembers that his brother Anupam's birthday is after 20th of August but before 28th of August, while Rita remembers that Anupam's birthday is before 22nd of August but after 12th of August. On which date does Anupam's birthday fall?

- (a) 20th August (b) 21th August
(c) 22nd August (d) Cannot be determined

Direction Q.(17 and 19) : Study the following information carefully and answer the questions given below:

Each of the six friends, A, B, C, D, E and F scored different marks in an examination. C scored more than only A and E. D scored less than only B. E did not score the least. The one who scored the third highest marks scored 81 marks. E scored 62 marks.

17. Which of the following could possibly be C's score?

- (a) 70 (b) 94 (c) 86 (d) 61

18. Which of the following is true with respect to the given information?

- (a) D's score was definitely less than 60
(b) F scored the maximum marks
(c) Only two people scored more than C
(d) There is a possibility that B scored 89 marks

19. The person who scored the maximum, scored 13 marks more than F's marks. Which of the following can be D's score?

- (a) 94 (b) 60 (c) 88 (d) 78

20. Seventeen people are standing in a straight line facing south. What is Bhavna's position from the left end of the line?

- I. Sandeep is standing second to the left of Sheetal. Only five people stand between Sheetal and the one who is standing at the extreme right end of the line. Four people stand between Sandeep and Bhavna.
II. Anita is standing fourth to the left of Sheetal. Less than three people are standing between Bhavna and Anita.
(a) if the data in statement (I) alone are sufficient to answer the question, while the data in statement

(II) alone are not sufficient to answer the question.

- (b) if the data in statement (II) alone are sufficient to answer the question, while the data in statement (I) alone are not sufficient to answer the question.
(c) if the data either in statement (I) alone or in statement (II) alone are sufficient to answer the question.
(d) if the data given in both the statements (I) and (II) together are not sufficient to answer the question.

SOLUTIONS


- 1. (d) :** $(2)^3 + (1)^2 \rightarrow (3)^3 + (2)^2 \rightarrow (4)^3 + (3)^2 \rightarrow (5)^3 + (4)^2 \rightarrow (6)^3 + (5)^2$
 $9 \rightarrow 31 \rightarrow 73 \rightarrow 141 \rightarrow 241$

SAMURAI SUDOKU



Samurai Sudoku puzzle consists of five overlapping sudoku grids. The standard sudoku rules apply to each 9×9 grid. Place digits from 1 to 9 in each empty cell. Every row, every column and every 3×3 box should contain one of each digit.

The puzzle has a unique answer.



Readers can send their responses at editor@mtg.in or post us with complete address. Winners' name with their valuable feedback will be published in next issue.

2. (a):

A	2	E
↓ +2	↓ × 2+1	↓ +3
C	5	H
↓ +2	↓ × 2+2	↓ +3
E	12	K
↓ +2	↓ × 2+3	↓ +3
G	27	N
↓ +2	↓ × 2+4	↓ +3
I	58	Q

3. (c): The relationship between GRIN and RING is same as 4123 and 1234. Last letter becomes the first letter and rest remain as it is.

4. (d): The relationship is $x^3 : (x/2)^4$.

5. (c): F $\xrightarrow{+2}$ H, U $\xrightarrow{+2}$ W, N $\xrightarrow{+2}$ P
Similarly, T $\xrightarrow{-2}$ R, C $\xrightarrow{-2}$ A, F $\xrightarrow{-2}$ D
K $\xrightarrow{-2}$ I, Q $\xrightarrow{-2}$ O

6. (d): As,

$$\frac{L+U+X+O+R}{3} = \frac{12+21+24+15+18}{3} = \frac{90}{3} = 30$$

In the same way, $\frac{G+U+I+L+D+S}{3}$

$$= \frac{7+21+9+12+4+19}{3} = \frac{72}{3} = 24$$

7. (c): Partly is not a type of bread.

8. (c): Except Bat, all others are birds. Bat is a mammal.


9. (a): $10 + (-3)^1 = 7$; $7 + (-3)^2 = 7 + 9 = 16$
 $16 + (-3)^3 = 16 - 27 = -11$
 $-11 + (-3)^4 = -11 + 81 = 70$
 $70 + (-3)^5 = 70 - 243 = -173$

10. (d): $29 + 46 = 75$

Now, $7 + 5 = 12$ and $1 + 2 = 3$; $18 + 59 = 77$


Now, $7 + 7 = 14$ and $1 + 4 = 5$

So, $158 + 283 = 441$; $4 + 4 + 1 = 9$



11. (b): Clearly, R is sister of P.

12. (d): Either (a) or (b)



Thus, $AB^2 = AC^2 + BC^2 = (20)^2 + (25)^2$
 $\Rightarrow AB^2 = 1025 \Rightarrow AB = 5\sqrt{41}$ m

According to the figure, direction is North-East.

14. (c): **Statements:** $G \leq K = O \geq U > P < S$
Conclusions:

I. $G \leq U$: Not true II. $S > O$: Not true
Hence, neither conclusion (I) nor (II) is true.

15. (a): **Conclusions:**

I. $Z > I$: True II. $S < I$: Not true
Hence, only conclusion (I) is true.

16. (b): According to Vinay, his brother Anupam's birthday may be on 21st, 22nd, 23rd, 24th, 25th, 26th, or 27th August.

According to Rita, Anupam's birthday may be on 13th, 14th, 15th, 16th, 17th, 18th, 19th, 20th or 21st August.

Hence, the common date is 21st August.

(17 - 19): B, D, F > C > A, E

B > D > F > C > A, E; B > D > F > C > E > A

Score of F = 81; Score of E = 62

17. (a): C scored more than 62 and less than 81.
 $62 < 70 < 81$


18. (d): D's score was more than 81 but less than B score. B scored the maximum marks.

If B scored 89, then there is a possibility that D's score can be 88, 87, 86, 85, 84, 83 and 82.

19. (c): Score of F = 81; Score of B = $81 + 13 = 94$

Therefore, score of D = more than 81 and less than 94
 $81 < 88 < 94$

20. (a): From statement (I),



We can conclude that Bhavna's position is 4th from the left end. From statement (II), nothing can be concluded.

beat the TIME TRAP



Duration : 60 minutes

SECTION-I

Single Option Correct Type

- The domain of $f(x) = \sqrt{\left(\frac{1}{\sqrt{x}} - \sqrt{x+1}\right)}$ is
 (a) $x > -1$ (b) $(-1, \infty) \setminus \{0\}$
 (c) $\left(0, \frac{\sqrt{5}-1}{2}\right]$ (d) $\left[\frac{1-\sqrt{5}}{2}, 0\right]$
- The solution of system of equations $x + (\sin \alpha)y = 1$ and $(\sin \alpha)x + 4y = 2$ satisfy $x \geq \frac{4}{5}$ & $y \leq \frac{1}{2}$, then
 (a) $\alpha \in \left[\frac{\pi}{4}, \frac{\pi}{3}\right]$ (b) $\alpha \in \left[0, \frac{\pi}{6}\right]$
 (c) $\alpha \in \left[\frac{\pi}{6}, \frac{\pi}{2}\right]$ (d) None of these
- If $f(x) = \left(2 - \frac{x}{a}\right)^{\tan\left(\frac{\pi x}{2a}\right)}$, $x \neq a$, the value which should be assigned to f at $x = a$ so that it is continuous every where is
 (a) $2/\pi$ (b) $e^{-2/\pi}$ (c) 2 (d) $e^{2/\pi}$
- A parallelogram is constructed on the vectors $\vec{a} = 3\vec{\alpha} - \vec{\beta}$, $\vec{b} = \vec{\alpha} + 3\vec{\beta}$. If $|\vec{\alpha}| = |\vec{\beta}| = 2$ and angle between $\vec{\alpha}$ and $\vec{\beta}$ is $\pi/3$, then length of a diagonal of the parallelogram is
 (a) $4\sqrt{5}$ units (b) $3\sqrt{2}$ units
 (c) $4\sqrt{3}$ units (d) $5\sqrt{3}$ units
- If $1, \alpha_1, \alpha_2, \dots, \alpha_{2008}$ are $(2009)^{\text{th}}$ roots of unity, then the value of $\sum_{r=1}^{2008} r(\alpha_r + \alpha_{2009-r})$ equals
 (a) 2009 (b) 2008 (c) 0 (d) -2009
- In the next world cup of cricket there will be 12 teams, divided equally in two groups. Teams of each group will play a match against each other. From each group 3 top teams will qualify for the next round. In this

round each team will play against others once. Four top teams of this round will qualify for the semifinal round, when each team will play against the others once. Two top teams of this round will go to the final round, where they will play the best of three matches. The minimum number of matches in the next world cup will be
 (a) 54 (b) 53 (c) 52 (d) 51

- If X denote the number of times heads occur in n tosses of a fair coin. If $P(X = 4)$, $P(X = 5)$ and $P(X = 6)$ are in A.P.; then value of n is
 (a) 7, 14 (b) 10
 (c) 12 (d) none of these
- The angle between the pair of tangents drawn from the point $(2, 4)$ to the circle $x^2 + y^2 = 4$ is
 (a) $\tan^{-1}(3/8)$ (b) $\tan^{-1}(4/3)$
 (c) 90° (d) None of these
- If $x \in (-\infty, -2)$ and $y^3 - 3y + x = 0$, then
 (a) y is not a monotonic function
 (b) y is an increasing function of x
 (c) y is not a function of x
 (d) y is a decreasing function of x
- The value of the integral $\int \frac{\sin^3 x + \sin^5 x}{\cos^2 x + \cos^4 x} dx$ is
 (a) $-\cos x + 2(\cos x)^{-1} + 6 \tan^{-1}(\cos x) + C$
 (b) $\cos x - 2(\cos x)^{-1} + 4 \tan^{-1}(\cos x) + C$
 (c) $\cos x - 2(\cos x)^{-1} + \tan^{-1}(\cos x) + C$
 (d) $\sec x + \tan x \sin x - 6 \tan^{-1}(\cos x) + C$

SECTION-II

Numerical Answer Type

- If $\tan \alpha$, $\tan \beta$ and $\tan \gamma$; $\alpha, \beta, \gamma \neq \frac{(2n-1)\pi}{2}$, $n \in \mathbb{N}$ be the slopes of three line segments OA , OB and OC , respectively, where O is origin. If circumcentre of $\triangle ABC$ coincides with origin and its orthocentre lies on y -axis, then the value of $\left(\frac{\cos 3\alpha + \cos 3\beta + \cos 3\gamma}{\cos \alpha \cos \beta \cos \gamma}\right)^2$ is equal to _____.

12. If the greatest and least values of the function $f(x) = x^3 - 6x^2 + 9x + 1$ on $[0, 2]$ are λ and μ , then the value of $\lambda^4 + \mu^8$ must be _____.

13. If p and r are the roots of the equation $x^2 - 8ax + 2a = 0$ and q and s are the roots of the equation $x^2 + 12bx + 6b = 0$, such that $\frac{1}{p}, \frac{1}{q}, \frac{1}{r}, \frac{1}{s}$ are

in A.P., (where a and b are two non zero real numbers) then $a^{-1} - b^{-1}$ is equal to _____.

14. If $f: \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function such that $f(x) + f(x+1) = 2$, for all $x \in \mathbb{R}$. If $I_1 = \int_0^8 f(x)dx$ and $I_2 = \int_{-1}^3 f(x)dx$, then the value of $I_1 + 2I_2$ is equal to _____.

15. A tangent is drawn to the curve $y = f(x)$ at $P(x, y)$ cuts the x and y axes at A and B respectively such that $AP : PB = 1 : 3$. If $f(1) = 1$, then the curve passes through $\left(k, \frac{1}{8}\right)$, where $k =$ _____.

SOLUTIONS

1. (c) : Here $f(x) = \sqrt{\left(\frac{1}{\sqrt{x}} - \sqrt{x+1}\right)}$

$f(x)$ will be defined, if

$$\frac{1}{\sqrt{x}} - \sqrt{x+1} \geq 0, x \neq 0, x+1 \geq 0 \text{ and } x > 0$$

$$\Rightarrow 1 - \sqrt{x^2 + x} \geq 0 \Rightarrow \sqrt{x^2 + x} \leq 1 \Rightarrow x^2 + x - 1 \leq 0$$

$$\Rightarrow \left(x + \frac{1-\sqrt{5}}{2}\right) \left(x + \frac{1+\sqrt{5}}{2}\right) \leq 0$$

$$\Rightarrow \frac{-1-\sqrt{5}}{2} \leq x \leq \frac{\sqrt{5}-1}{2} \quad \dots(i)$$

$$\text{Also, } x+1 \geq 0 \Rightarrow x \geq -1 \text{ and } x > 0 \quad \dots(ii)$$

$$\text{From (i) and (ii), } x \in \left(0, \frac{\sqrt{5}-1}{2}\right)$$

2. (b)

3. (d) : For $f(x)$ to be continuous, we must have

$$f(a) = \lim_{x \rightarrow a} \left(2 - \frac{x}{a}\right)^{\tan\left(\frac{\pi x}{2a}\right)}$$

$$\text{Now, } \log f(x) = \tan\left(\frac{\pi x}{2a}\right) \log\left(2 - \frac{x}{a}\right)$$

$$\Rightarrow \lim_{x \rightarrow a} \log f(x) = \frac{\log\left(2 - \frac{x}{a}\right)}{\cot\left(\frac{\pi x}{2a}\right)} \left(\frac{0}{0} \text{ form}\right)$$

$$= \lim_{x \rightarrow a} \frac{-1/a}{-\frac{\pi}{2a} \operatorname{cosec}^2\left(\frac{\pi x}{2a}\right)} \quad (\text{Using } L' \text{ Hospital's Rule})$$

$$= \frac{2}{\pi} \lim_{x \rightarrow a} \frac{1}{2 - \frac{x}{a}} \times \lim_{x \rightarrow a} \sin^2\left(\frac{\pi x}{2a}\right) = \frac{2}{\pi}$$

$$\text{Hence, } \lim_{x \rightarrow a} f(x) = e^{2/\pi} \Rightarrow f(a) = e^{2/\pi}$$

4. (c) : Let $AB = \vec{a} = 3\vec{\alpha} - \vec{\beta}$, $BC = \vec{b} = \vec{\alpha} + 3\vec{\beta}$

Diagonal, $AC = AB + BC = \vec{a} + \vec{b}$

$$\Rightarrow |AC| = |\vec{a} + \vec{b}| \Rightarrow |AC| = |4\vec{\alpha} + 2\vec{\beta}|$$

$$\Rightarrow |AC|^2 = 16|\vec{\alpha}|^2 + 4|\vec{\beta}|^2 + 16\vec{\alpha} \cdot \vec{\beta}$$

$$\Rightarrow |AC|^2 = 64 + 16 + 16|\vec{\alpha}||\vec{\beta}|\cos\frac{\pi}{3}$$

$$= 80 + 16 \times 4 \times \frac{1}{2} = 112 \Rightarrow |AC| = 4\sqrt{7} \text{ units}$$

Other diagonal is $|BD| = |\vec{a} - \vec{b}|$

$$\Rightarrow |BD|^2 = |2\vec{\alpha} - 4\vec{\beta}|^2 = 4|\vec{\alpha}|^2 + 16|\vec{\beta}|^2 - 16|\vec{\alpha}||\vec{\beta}|\cos\frac{\pi}{3}$$

$$\Rightarrow |BD|^2 = 16 + 64 - 64\left(\frac{1}{2}\right) \Rightarrow |BD| = \sqrt{48} = 4\sqrt{3} \text{ units}$$

5. (d)

6. (b) : The number of matches in first round

$$= {}^6C_2 + {}^6C_2 = 30$$

The number of matches in next round = ${}^6C_2 = 15$

and the number of matches in the semifinal round

$$= {}^4C_2 = 6$$

Number of matches in the final round = 2

Hence, required number of matches

$$= 30 + 15 + 6 + 2 = 53.$$

7. (a) : Clearly, X is a binomial variate with parameters

n and $p = 1/2$ such that $P(X = r)$

$$= {}^nC_r \left(\frac{1}{2}\right)^r \left(\frac{1}{2}\right)^{n-r} = {}^nC_r \left(\frac{1}{2}\right)^n$$

Now, $P(X = 4)$, $P(X = 5)$ and $P(X = 6)$ are in A.P.

$$\Rightarrow 2P(X = 5) = P(X = 4) + P(X = 6)$$

$$\Rightarrow 2 \cdot {}^nC_5 \left(\frac{1}{2}\right)^n = {}^nC_4 \left(\frac{1}{2}\right)^n + {}^nC_6 \left(\frac{1}{2}\right)^n$$

$$\Rightarrow 2 \cdot {}^nC_5 = {}^nC_4 + {}^nC_6$$

$$\Rightarrow 2 \frac{n!}{(n-5)!5!} = \frac{n!}{(n-4)!4!} + \frac{n!}{(n-6)!6!}$$

$$\Rightarrow \frac{2}{5(n-5)} = \frac{1}{(n-4)(n-5)} + \frac{1}{6 \times 5}$$


$$\Rightarrow n^2 - 21n + 98 = 0 \Rightarrow (n - 7)(n - 14) = 0$$

$$\Rightarrow n = 7 \text{ or } 14.$$

8. (b): Equation of pair of tangents to a circle is $T^2 = SS_1$, where $T = xx_1 + yy_1 - 4$

$$S_1 = x_1^2 + y_1^2 - 4$$

$$S = x^2 + y^2 - 4$$



$$\therefore (2x + 4y - 4)^2 = (x^2 + y^2 - 4)(4 + 16 - 4)$$

$$\Rightarrow 4(x + 2y - 2)^2 = 16(x^2 + y^2 - 4)$$

$$\Rightarrow x^2 + 4y^2 + 4xy - 8y - 4x + 4 = 4x^2 + 4y^2 - 16$$

$$\Rightarrow 3x^2 - 4xy + 8y + 4x - 20 = 0$$

$$= ax^2 + 2hxy + by^2 + 2gx + 2fy + c$$

$$\therefore \tan \theta = \frac{2\sqrt{h^2 - ab}}{a + b} = \frac{2 \times 2}{3} \Rightarrow \theta = \tan^{-1}(4/3)$$

9. (d): $y^3 - 3y + x = 0$

$$\Rightarrow x = -y^3 + 3y < -2 \quad (\because x \in (-\infty, -2))$$

$$\Rightarrow y^3 - 3y - 2 > 0 \Rightarrow (y + 1)^2(y - 2) > 0 \therefore y > 2$$

$$\text{Now, } x = -y^3 + 3y$$

$$\therefore \frac{dx}{dy} = -3y^2 + 3 = 3(1 - y^2)$$

$$\Rightarrow \frac{dy}{dx} = \frac{1}{3(1 - y^2)} < 0 \quad (\text{as } y > 2)$$

$\Rightarrow y$ is a decreasing function of x .

10. (a)

11. (144): Let $A(r \cos \alpha, r \sin \alpha)$, $B(r \cos \beta, r \sin \beta)$ and $C(r \cos \gamma, r \sin \gamma)$.

Since, circumcentre and orthocentre both lies on y -axis.

$\therefore$ Centroid also lies on y -axis.

$$\Rightarrow \cos \alpha + \cos \beta + \cos \gamma = 0$$

$$\therefore \cos^3 \alpha + \cos^3 \beta + \cos^3 \gamma = 3 \cos \alpha \cdot \cos \beta \cdot \cos \gamma \dots (i)$$

$$\text{Now, } \cos 3\alpha + \cos 3\beta + \cos 3\gamma = 4(\cos^3 \alpha + \cos^3 \beta + \cos^3 \gamma)$$

$$\Rightarrow \frac{\cos 3\alpha + \cos 3\beta + \cos 3\gamma}{\cos^3 \alpha + \cos^3 \beta + \cos^3 \gamma} = 4 \dots (ii)$$

Using (i) and (ii), we get

$$\left(\frac{\cos 3\alpha + \cos 3\beta + \cos 3\gamma}{\cos \alpha \cdot \cos \beta \cdot \cos \gamma} \right)^2 = 12^2 = 144$$

12. (626): Given, $f(x) = x^3 - 6x^2 + 9x + 1$

$$\therefore f'(x) = 3x^2 - 12x + 9 = 3(x - 1)(x - 3)$$

For maxima or minima put $f'(x) = 0$

$$\Rightarrow x = 1, 3 \text{ but } 3 \notin [0, 2]$$

$\therefore$ Only one critical point in $[0, 2]$ is $x = 1$.

Now, greatest value, $\lambda = \max. \{f(0), f(1), f(2)\} = \max. \{1, 5, 3\} = 5$

and least value, $\mu = \min. \{f(0), f(1), f(2)\} = \min. \{1, 5, 3\} = 1$

$$\therefore \lambda^4 + \mu^8 = 5^4 + 1^8 = 625 + 1 = 626$$

13. (38): p and r are the roots of the equation, $x^2 - 8ax + 2a = 0$

$$\therefore p + r = 8a, pr = 2a \Rightarrow 1/p + 1/r = 4 \dots (i)$$

Also, q and s are the roots of $x^2 + 12bx + 6b = 0$

$$\therefore q + s = -12b, qs = 6b \Rightarrow 1/q + 1/s = -2 \dots (ii)$$

Given that $\frac{1}{p}, \frac{1}{q}, \frac{1}{r}, \frac{1}{s}$ are in A.P.

$$\therefore \frac{1}{q} - \frac{1}{p} = \frac{1}{r} - \frac{1}{q} \Rightarrow \frac{2}{q} = \frac{1}{r} + \frac{1}{p} = 4 \quad [\text{Using (i)}]$$

$$\Rightarrow q = \frac{1}{2} \text{ and } s = \frac{-1}{4}$$

$$\text{Clearly, } \frac{1}{r} - \frac{1}{q} = \frac{1}{s} - \frac{1}{r} \Rightarrow \frac{2}{r} = \frac{1}{q} + \frac{1}{s}$$

$$\Rightarrow \frac{2}{r} = -2 \Rightarrow r = -1 \text{ and } p = \frac{1}{5} \quad [\text{Using (ii)}]$$

$$\therefore \frac{1}{a} - \frac{1}{b} = \frac{2}{pr} - \frac{6}{qs} = 38$$

14. (16): We have, $f(x) + f(x + 1) = 2 \dots (i)$

Replacing x with $x + 1$, we get

$$f(x + 1) + f(x + 2) = 2 \dots (ii)$$

From (i) and (ii), we have $f(x) = f(x + 2)$

$\Rightarrow f(x)$ is periodic with period 2.

$$\text{Now, } I_1 = \int_0^8 f(x) dx = 4 \int_0^2 f(x) dx$$

$$\text{and } I_2 = \int_{-1}^3 f(x) dx = 2 \int_0^2 f(x) dx \Rightarrow I_1 = 2I_2$$

$$\therefore I_1 = 4 \left[\int_0^1 f(x) dx + \int_1^2 f(x) dx \right]$$

$$= 4 \left[\int_0^1 f(x) dx + \int_0^1 f(x + 1) dx \right]$$

$$= 4 \left[\int_0^1 f(x) dx + \int_0^1 (2 - f(x)) dx \right] \quad [\text{Using (i)}]$$

$$= 4 \left[\int_0^1 f(x) dx + \int_0^1 2 dx - \int_0^1 f(x) dx \right] = 8$$

$$\Rightarrow I_1 = 8 \Rightarrow I_2 = 4 \therefore I_1 + 2I_2 = 8 + 8 = 16$$

15. (2): $\frac{3x}{h} + \frac{y}{k} = 1$ is the

equation of the tangent whose slope is $-3k/h$.

$$\frac{dy}{dx} = -\frac{3y}{x} \Rightarrow x \frac{dy}{dx} + 3y = 0$$


$$\Rightarrow \frac{dy}{y} = -3 \frac{dx}{x} \Rightarrow \log y = -3 \log x + \log c$$

$$\Rightarrow y = cx^{-3} \Rightarrow y = \frac{c}{x^3}, \text{ which passes through } (1, 1)$$

$$\Rightarrow c = 1$$

$\therefore$ Equation of the curve is $y = 1/x^3$.

$$\therefore \text{ It passes through } \left(k, \frac{1}{8}\right) \therefore \frac{1}{8} = \frac{1}{k^3} \Rightarrow k = 2$$



YOU ASK WE ANSWER

Do you have a question that you just can't get answered?

Use the vast expertise of our MTG team to get to the bottom of the question. From the serious to the silly, the controversial to the trivial, the team will tackle the questions, easy and tough. The best questions and their solutions will be printed in this column each month.


1. If a circular sector of perimeter 60 meter with maximum area is to be constructed, then what will be the radius of the circular arc in meter. (Anushka, Karnataka)

Ans. Let OAB be the circular sector.

Perimeter of $OAB = 60$ m

$$\therefore \frac{\theta}{360} \times 2\pi r + 2r = 60$$

$$\Rightarrow \frac{\theta}{360} = \frac{60 - 2r}{2\pi r} \quad \dots(i)$$



We need maximum area, and we know that area of a

$$\text{sector } (A) = \frac{\theta}{360} \times \pi r^2$$

$$\text{i.e., } A = \frac{60 - 2r}{2\pi r} \times \pi r^2$$

$$\Rightarrow A = 30r - r^2$$

Differentiating w.r.t 'r', $\frac{dA}{dr} = 30 - 2r$

For critical point, we have $\frac{dA}{dr} = 0 \Rightarrow 30 - 2r = 0$

$$\Rightarrow r = \frac{30}{2} = 15 \text{ m. Now, } \frac{d^2A}{dr^2} = -2 < 0$$

$\therefore r = 15$ m gives the maximum area.

2. If $\vec{a} = -\hat{i} - \hat{k}$, $\vec{b} = -\hat{i} + \hat{j}$ and $\vec{c} = \hat{i} + 2\hat{j} + 3\hat{k}$ be three given vectors and $\vec{r}$ is a vector such that $\vec{r} \times \vec{b} = \vec{c} \times \vec{b}$ and $\vec{r} \times \vec{a} = 0$, then show that $\vec{r} \cdot \vec{b} = 9$.

(Rakesh, Assam)

Ans. We have, $\vec{a} = -\hat{i} - \hat{k}$, $\vec{b} = -\hat{i} + \hat{j}$, $\vec{c} = \hat{i} + 2\hat{j} + 3\hat{k}$.

Let $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$

Now given that, $\vec{r} \times \vec{b} = \vec{c} \times \vec{b}$

$$\text{So, } \vec{c} \times \vec{b} = -3\hat{i} - 3\hat{j} + 3\hat{k} \quad \dots(i)$$

$$\text{and } \vec{r} \times \vec{b} = -z\hat{i} - z\hat{j} + (x + y)\hat{k} \quad \dots(ii)$$

On comparing (i) and (ii), we get $z = 3$

$$\text{and } x + y = 3 \quad \dots(iii)$$

$$\text{Now, } \vec{r} \cdot \vec{a} = 0 \Rightarrow -x - z = 0 \Rightarrow x = -z \Rightarrow x = -3$$

$$\text{From (iii), } y = 3 - x = 3 - (-3) = 6$$

$$\Rightarrow \vec{r} = -3\hat{i} + 6\hat{j} + 3\hat{k}$$

$$\text{Now, } \vec{r} \cdot \vec{b} = (-3)(-1) + 6 \cdot 1 = 3 + 6 = 9$$

3. In a triangle PQR , the co-ordinates of the points P and Q are $(-2, 4)$ and $(4, -2)$ respectively and equation of the perpendicular bisector of PR is $2x - y + 2 = 0$, then find the centre of the circumcircle of the ΔPQR .

(Vibhor Singh, A.P.)

Ans. Let A is mid point of PQ .

So, coordinates of point A

$$= \left(\frac{-2 + 4}{2}, \frac{4 - 2}{2} \right) \text{ i.e., } (1, 1)$$

$$\text{Slope of } PQ = \frac{4 + 2}{-2 - 4} = -1$$

$$\Rightarrow \text{Slope of perpendicular bisector of } PQ = 1$$

So, equation of perpendicular bisector of PQ is


$$(y - 1) = 1(x - 1) \Rightarrow x - y = 0 \quad \dots(i)$$

Also, equation of perpendicular bisector of PR is

$$2x - y + 2 = 0 \quad \dots(ii)$$

Solving (i) and (ii), we get $x = -2$ and $y = -2$

$\Rightarrow$ Centre of circumcircle of ΔPQR is $(-2, -2)$.





ONLINE TEST SERIES

Practice Part Syllabus/ Full Syllabus
24 Mock Tests for


JEE Main



Now on your android smart phones
with the same login of web portal.

Log on to test.pcmbtoday.com

MONTHLY TEST DRIVE



This specially designed column enables students to self analyse their extent of understanding of specified chapters. Give yourself four marks for correct answer and deduct one mark for wrong answer. Self check table given at the end will help you to check your readiness.

Total Marks : 80

Series 8 : Straight Lines

Time Taken : 60 Min.

Only One Option Correct Type

- The equation of a line passing through $(a \cos^3 \theta, a \sin^3 \theta)$ and perpendicular to the line $x \sec \theta + y \operatorname{cosec} \theta = a$, is
 (a) $x \cos \theta - y \sin \theta = a \cos \theta$
 (b) $x \cos \theta - y \sin \theta = a \cos 2\theta$
 (c) $x \cos \theta - y \sin \theta = -a \cos 2\theta$
 (d) None of these
- If the points (1, 2) and (3, 4) were to be on the same side of the line $3x - 5y + a = 0$, then
 (a) $7 < a < 11$ (b) $a = 7$
 (c) $a = 1$ (d) $a < 7$ or $a > 11$
- A vertex of an equilateral triangle is (2, 3) and the opposite side is $x + y = 2$. The equations of other sides are
 (a) $(2 \pm \sqrt{3})x - y = 1 \pm 2\sqrt{3}$
 (b) $(2 \pm \sqrt{3})x - y = 1 \mp 2\sqrt{3}$
 (c) $(1 \pm 2\sqrt{3})x - y = 2 \pm 2\sqrt{3}$
 (d) $(2 \pm \sqrt{3})x + y = 1 \pm 2\sqrt{3}$
- Find the transformed equation of the straight line $2x - 3y + 5 = 0$, when the origin is shifted to the point (3, -1) after translation of axes.
 (a) $2x - 3y + 14 = 0$ (b) $2x - 5y + 14 = 0$
 (c) $2x - 3y + 15 = 0$ (d) $x - 4y + 14 = 0$
- The orthocentre of the triangle formed by the points (0, 0), (4, 0) and (3, 4) is
 (a) (2, 0) (b) (3/2, 2)
 (c) (3/4, 3) (d) (3, 3/4)
- The product of the perpendiculars drawn from the point (1, 2) to the pair of lines $x^2 + 4xy + y^2 = 0$ is

- $\frac{9}{4}$
- $\frac{3}{4}$
- $\frac{9}{16}$
- None of these

One or More Than One Option(s) Correct Type

- Equations of the diagonals of a rectangle are $y + 8x - 17 = 0$ and $y - 8x + 7 = 0$. If the area of the rectangle is 8 sq. units, then the equation of the sides of the rectangle is/are
 (a) $x = 1$ (b) $x + y = 1$
 (c) $y = 9$ (d) $x - 2y = 3$
- If (a, a) is not outside of lines $|x + y| = 2$, then
 (a) $|a| \in [0, 1]$ (b) $|a| \in [0, 2]$
 (c) $[|a|] \in \{0, 1, 2\}$ (d) $[|a|] = \{0, 1\}$
- Which of the following line is parallel to an angle bisector of pair of lines $x^2 - y^2 - 4y - 4 = 0$?
 (a) $x + 1 = 0$ (b) $y - 2 = 0$
 (c) $x + y + 2 = 0$ (d) $x - 1 = 0$



EXAM ALERT 2023

Exam	Date
JEE Main Session 2	6 th , 8 th , 10 th , 11 th and 12 th April
WB JEE	30 th April
CUET	21 st to 31 st May
MHT CET PCM	9 th to 13 th May
BITSAT	Session - 1 : 22 nd to 26 th May
	Session - 2 : 18 th to 22 nd June
JEE Advanced	4 th June

10. If the line $y = mx$ is one of the bisector of the lines $x^2 + 4xy - y^2 = 0$, then the value of m is equal to

(a) $\frac{-1+\sqrt{5}}{2}$ (b) $\frac{1+\sqrt{5}}{2}$
(c) $\frac{-1-\sqrt{5}}{2}$ (d) $\frac{1-\sqrt{5}}{2}$

11. All points inside the triangle formed by the points $(1, 3)$, $(5, 0)$ and $(-1, 2)$ satisfy

(a) $3x + 2y \geq 0$ (b) $2x + y - 13 \geq 0$
(c) $2x - 3y - 12 \leq 0$ (d) $-2x + y \geq 0$

12. If the point $P(x, y)$ is equidistant from the points $A(a + b, b - a)$ and $B(a - b, a + b)$, then

(a) $ax = by$ (b) $bx = ay$
(c) $x^2 - y^2 = 2(ax + by)$ (d) P can be (a, b) .

13. ABC is an equilateral triangle whose centroid is origin and base BC is along the line $11x + 60y = 122$, then

- (a) area of the triangle is numerically equal to the perimeter.
(b) area of triangle is numerically double the perimeter.
(c) area of triangle is numerically three times the perimeter.
(d) area of triangle is $12\sqrt{3}$ sq units.

Comprehension Type

Paragraph for Q. No. 14 to 15

A straight line L with negative slope passes through $(1, 4)$ and meets the coordinate axes at A and B .

14. What will be the value of the sum of the intercepts made by the line L on the coordinates axes if x intercept is 3.
(a) 12 (b) 9 (c) 8 (d) 6
15. If the area of the triangle OAB is 8 units then the equation of the line L is
(a) $8x + y = 16$ (b) $16x + y = 32$
(c) $4x + y = 8$ (d) $8x + y = 8$

Matrix Match Type

16. Match the following :

Column-I		Column-II	
(P)	The coordinates of the points P and Q on the line $x + 5y = 13$ which are at a distance of 2 units from the line $12x - 5y + 26 = 0$ are	(1)	$(3, 1), (-7, 11)$
(Q)	The coordinates of the point on the line $x + y = 4$, which are at a unit distance from the line $4x + 3y - 10 = 0$ are	(2)	$\left(-\frac{1}{3}, \frac{11}{3}\right), \left(\frac{4}{3}, \frac{7}{3}\right)$
(R)	The coordinates of the point on the line joining $A(-2, 5)$ and $B(3, 1)$ such that $AP = PQ = QB$ are	(3)	$\left(1, \frac{12}{5}\right), \left(-3, \frac{16}{5}\right)$

	P	Q	R
(a)	1	2	3
(b)	2	3	1
(c)	3	1	2
(d)	3	2	1

Numerical Answer Type

17. If pair of lines $x^2 - 4pxy - y^2 = 0$ and $x^2 + 6qxy - y^2 = 0$ is such that each pair bisects the angle between the other pair, then pq equals _____.
18. If the straight lines joining the origin to the points of intersection of $2x + 3y = k$ and $3x^2 - xy + 3y^2 + 2x - 3y - 4 = 0$ are at right angle, then $6k - \frac{52}{k}$ is _____.
19. The area enclosed by the graphs of $|x + y| = 2$ and $|x| = 1$ (in sq. units) is _____.
20. If the equation $12x^2 - 10xy + 2y^2 + 11x - 5y + \lambda = 0$ represents a pair of straight lines, then the value of $221\lambda^2 + 361\lambda + 582$ is equal to _____.



Keys are published in this issue. Search now! ☺

SELF CHECK

No. of questions attempted
No. of questions correct
Marks scored in percentage

Check your score! If your score is

> 90%	EXCELLENT WORK !	You are well prepared to take the challenge of final exam.
90-75%	GOOD WORK !	You can score good in the final exam.
74-60%	SATISFACTORY !	You need to score more next time.
< 60%	NOT SATISFACTORY!	Revise thoroughly and strengthen your concepts.





Happy Pi DAY!



March 14 marks **Pi Day**, an annual celebration of the mathematical symbol **Pi**. **Pi Day** was first organised by **Larry Shaw** in 1988. The earliest known use of the Greek letter π came in 1706. The purpose of **Pi Day** is simply to celebrate the important and mysterious number.

What is Pi?

Pi is the mathematical constant whose value is the ratio of a circle's circumference to its diameter. The first calculation of π was done by Archimedes of Syracuse.



Albert Einstein's Birthday
March 14, 1879

In addition to it being **Pi Day**, it is also a significant date in the lives of two renowned scientists! On this day we commemorate the birth of Albert Einstein and the passing of Stephen Hawking. Both were aged 76 when they died. Is this Cosmic or Coincidence?



Stephen Hawking's Death
Anniversary
March 14, 2018



CBSE warm-up!

Exam on
11th March
2023

CLASS-XII

Practice questions for CBSE Exams as per the latest pattern
and reduced syllabus by CBSE for the academic session 2022-23.

Practice Paper 2022-23

Time Allowed : 3 hours
Maximum Marks : 80

General Instructions :

- This Question paper contains - five sections A, B, C, D and E. Each section is compulsory. However, there are internal choices in some questions.
- Section A has 18 MCQ's and 02 Assertion-Reason based questions of 1 mark each.
- Section B has 5 Very Short Answer (VSA)-type questions of 2 marks each.
- Section C has 6 Short Answer (SA)-type questions of 3 marks each.
- Section D has 4 Long Answer (LA)-type questions of 5 marks each.
- Section E has 3 source based/case based/passage based/integrated units of assessment (4 marks each) with sub parts.

SECTION - A (MULTIPLE CHOICE QUESTIONS)

Each question carries 1 mark.

- If $A = \begin{bmatrix} k & 8 \\ 4 & 2k \end{bmatrix}$ is a singular matrix, then the value of k is
(a) 4 (b) -4 (c) ± 4 (d) 0

- If $A = \begin{bmatrix} 2 & 5 \\ -11 & 7 \end{bmatrix}$, then $(\text{adj } A)'$ is equal to

- $\begin{bmatrix} -2 & -5 \\ 11 & -7 \end{bmatrix}$
- $\begin{bmatrix} 7 & 5 \\ 11 & 2 \end{bmatrix}$
- $\begin{bmatrix} 7 & 11 \\ -5 & 2 \end{bmatrix}$
- $\begin{bmatrix} 7 & -5 \\ 11 & 2 \end{bmatrix}$

- $\vec{a} \cdot (\vec{b} + \vec{c})$ is equal to

- $\vec{a} \cdot \vec{b} + \vec{a} \cdot \vec{c}$
- $\vec{a} + \vec{b} \cdot \vec{c}$
- $\vec{a} \cdot \vec{b} + \vec{c}$
- $\vec{a} \cdot \vec{b} + \vec{a} \cdot \vec{b} \cdot \vec{c}$

- If $y = 5\cos x - 3\sin x$, then $\frac{d^2 y}{dx^2}$ is equal to

- y
- y
- 25y
- 9y


- Evaluate : $\int_0^{\pi/2} x \cos x \, dx$

- $\frac{\pi}{6} - 1$
- $\frac{\pi}{3}$
- $\frac{\pi}{2} - 1$
- $\frac{\pi}{3} - 2$

- Solve the differential equation $\frac{dy}{dx} = \left(\frac{y}{x}\right)^{1/3}$.

- $y^{1/3} - x^{-1/3} = c$
- $y^{2/3} - x^{2/3} = c$
- $y^{-1/3} - x^{-1/3} = c$
- $y^{2/3} - x^{-2/3} = c$

- Maximise $Z = 3x + 2y$, if the shaded region is the feasible region.



- (a) Max. $Z = 10$ (b) Max. $Z = 15$
 (c) Max. $Z = 18$ (d) none of these
8. Find the value of $(\vec{u} \cdot \hat{i})^2 + (\vec{u} \cdot \hat{j})^2 + (\vec{u} \cdot \hat{k})^2$.
 (a) $|\vec{u}|^2$ (b) $|\vec{u}|$ (c) 0 (d) 1
9. Evaluate : $\int |x| dx$
 (a) $I = \frac{|x|}{2} + C$ (b) $I = \frac{x|x|}{2} + C$
 (c) $I = \frac{x^2|x|}{2} + C$ (d) $I = \frac{x}{2} + C$
10. If $A = \begin{bmatrix} \alpha & 2 \\ 2 & \alpha \end{bmatrix}$ and $|A^3| = 27$, then $\alpha =$
 (a) ± 2 (b) $\pm \sqrt{5}$ (c) ± 1 (d) $\pm \sqrt{7}$
11. Corner points of the feasible region for an LPP are $(0, 2)$, $(3, 0)$, $(6, 0)$, $(6, 8)$ and $(0, 5)$. Let $F = 4x + 6y$ be the objective function. The minimum value of F occurs at
 (a) $(0, 2)$ only (b) $(3, 0)$ only
 (c) the mid-point of the line segment joining the points $(0, 2)$ and $(3, 0)$ only
 (d) any point on the line segment joining the points $(0, 2)$ and $(3, 0)$
12. If $e \begin{bmatrix} e^x & e^y \\ e^y & e^x \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$, then the values of x and y are respectively
 (a) $-1, -1$ (b) $1, 1$ (c) $0, 1$ (d) $1, 0$
13. Let A and B be independent events with $P(A) = 1/4$ and $P(A \cup B) = 2P(B) - P(A)$. Find $P(B)$.
 (a) $1/5$ (b) $2/5$ (c) $3/5$ (d) $4/5$
14. If the matrix $A = \begin{bmatrix} 1 & -2 & 3 \\ 1 & 2 & 1 \\ x & 2 & -3 \end{bmatrix}$ is singular, then find x .
 (a) 1 (b) -2 (c) -1 (d) 0
15. The general solution of the differential equation $\frac{dy}{dx} = 1 - x + y - xy$ is
 (a) $\log|1+y| = x - \frac{x^2}{2} + C$
 (b) $\log|1+y| = x^2 - \frac{x}{2} + C$
 (c) $\log|1+y| = x^2 - \frac{x}{3} + C$

(d) $\log|1+y| = \frac{x^3}{3} - x + C$

16. Find the order of the differential equation whose general solution is given by $y = (A + B) \cos(x + C) + De^x$.
 (a) 4 (b) 5 (c) 3 (d) 2
17. If the vector $\vec{a} = 3\hat{i} - 2\hat{j} + 6\hat{k}$, then the magnitude of $\vec{a}$ is
 (a) 6 (b) 7 (c) 9 (d) 5
18. The direction cosines of the line $\frac{4-x}{2} = \frac{y}{6} = \frac{1-z}{3}$ are
 (a) $\frac{2}{7}, \frac{3}{7}, \frac{6}{7}$ (b) $\frac{1}{7}, \frac{2}{7}, \frac{3}{7}$
 (c) $\frac{2}{7}, \frac{-6}{7}, \frac{3}{7}$ (d) $\frac{1}{7}, \frac{-6}{7}, \frac{-2}{7}$

ASSERTION-REASON BASED QUESTIONS

In the following questions, a statement of Assertion (A) is followed by a statement of Reason (R). Choose the correct answer out of the following choices.

- (a) Both A and R are true and R is the correct explanation of A.
 (b) Both A and R are true but R is not the correct explanation of A.
 (c) A is true but R is false.
 (d) A is false but R is true.
19. Assertion (A): The domain of the function $\cos^{-1}(2x - 1)$ is $[0, 1]$
 Reason (R): The domain of the function $\cos^{-1} x$ is $[0, \pi]$
20. Assertion (A): The two lines with direction cosines $\left(\frac{12}{13}, \frac{-3}{13}, \frac{-4}{13}\right), \left(\frac{4}{13}, \frac{12}{13}, \frac{3}{13}\right)$ are mutually perpendicular.
 Reason (R): The two lines with direction cosines (l_1, m_1, n_1) and (l_2, m_2, n_2) are mutually perpendicular iff $l_1 l_2 + m_1 m_2 + n_1 n_2 = 0$.

SECTION - B

This section comprises of very short answer type questions (VSA) of 2 marks each.

21. Find the value of $\left[\frac{\sin^{-1}\left(\frac{1}{\sqrt{2}}\right) - 3 \sin^{-1}\left(\frac{\sqrt{3}}{2}\right)}{\cos^{-1}\left(\frac{1}{\sqrt{2}}\right) - 3 \cos^{-1}\left(\frac{\sqrt{3}}{2}\right)} \right]$,

where $[\cdot]$ is greatest integer function.

OR

If $\vec{a} = 3\hat{i} - \hat{j}$ and $\vec{b} = 2\hat{i} + \hat{j} - 3\hat{k}$ then express $\vec{b}$ in the form $\vec{b} = \vec{b}_1 + \vec{b}_2$ where $\vec{b}_1 \parallel \vec{a}$ and $\vec{b}_2 \perp \vec{a}$.

34. Express the following matrix as the sum of a symmetric matrix and a skew-symmetric matrix

$$\begin{bmatrix} 3 & -2 & -4 \\ 3 & -2 & -5 \\ -1 & 1 & 2 \end{bmatrix}$$

35. Let $A = \mathbb{R} - \{2\}$ and $B = \mathbb{R} - \{1\}$. If $f: A \rightarrow B$ is a function defined by $f(x) = \frac{x-1}{x-2}$, then show that f is one-one and onto.

OR


Show that the relation R in the set of real numbers, defined as $R = \{(a, b) : a \leq b^2\}$ is neither reflexive, nor symmetric, nor transitive.

SECTION - E

This section comprises of 3 case-study/passage-based questions of 4 marks each with sub-parts. First two case study questions have three sub-parts (i), (ii), (iii) of marks 1, 1, 2 respectively. The third case study question has two sub-parts of 2 marks each.

36. **Case-Study 1 :** Read the following passage and answer the questions given below.

On a holiday, a father gave a puzzle from a newspaper to his son Ravi and his daughter Priya. The probability of solving this specific puzzle independently by Ravi and Priya are $\frac{1}{4}$ and $\frac{1}{5}$ respectively.



- Find the chance that both Ravi and Priya solved the puzzle.
- Find the probability that puzzle is solved by Ravi but not by Priya.
- Find the probability that puzzle is solved.

OR

Find the probability that exactly one of them solved the puzzle.

37. **Case-Study 2 :** Read the following passage and answer the questions given below.

The fuel cost per hour for running a train is proportional to the square of the speed it generates in km per hour. If the fuel costs



₹ 48 per hour at speed 16 km per hour and the fixed charges to run the train amount to ₹ 1200 per hour. Assume the speed of the train as v km/h.


- Given that the fuel cost per hour is k times the square of the speed the train generates in km/h, find the value of k .
- If the train has travelled a distance of 500 km, then find the function of total cost of running the train.
- Determine the most economical speed to run the train.

OR

Determine the fuel cost for the train to travel 500 km at the most economical speed.

38. **Case-Study 3 :** Read the following passage and answer the questions given below.


Rohan, a student of class XII, visited his uncle's flat with his father. He observed that the window of the house is in the form of a rectangle surmounted by a semicircular opening having perimeter 10 m as shown in the figure.



- Find the area (A) of the window.
- What is the maximum area of the window?



COMIC CAPSULE



SOLUTIONS

1. (c) : $\because A$ is singular matrix.

$$\therefore |A| = 0$$

$$\Rightarrow \begin{vmatrix} k & 8 \\ 4 & 2k \end{vmatrix} = 0 \Rightarrow 2k^2 - 32 = 0 \Rightarrow k^2 = 16 \Rightarrow k = \pm 4$$

2. (c) : We know that, $(\text{adj } A)' = \text{cofactor matrix of } A$

$$\text{Here, cofactor matrix of } A = \begin{bmatrix} 7 & 11 \\ -5 & 2 \end{bmatrix} = (\text{adj } A)'$$

3. (a) : $\vec{a} \cdot (\vec{b} + \vec{c}) = \vec{a} \cdot \vec{b} + \vec{a} \cdot \vec{c}$ (By distributive law)

4. (a) : We have, $y = 5 \cos x - 3 \sin x$

$$\Rightarrow \frac{dy}{dx} = -5 \sin x - 3 \cos x$$

$$\Rightarrow \frac{d^2 y}{dx^2} = -5 \cos x + 3 \sin x = -y$$

$$5. (c) : \int_0^{\pi/2} x \cos x \, dx = [x \sin x]_0^{\pi/2} - \int_0^{\pi/2} 1 \cdot \sin x \, dx$$

[Integrating by parts]

$$= \frac{\pi}{2} + [\cos x]_0^{\pi/2} = \frac{\pi}{2} - 1$$

$$6. (b) : \text{We have, } \frac{dy}{dx} = \left(\frac{y}{x}\right)^{1/3} \Rightarrow y^{-1/3} dy = x^{-1/3} dx$$

Integrating both sides, we get

$$\frac{3}{2} y^{2/3} = \frac{3}{2} x^{2/3} + k \Rightarrow y^{2/3} - x^{2/3} = \frac{2}{3} k = c \text{ (say)}$$

Hence, required solution is $y^{2/3} - x^{2/3} = c$

7. (c) : Let us find the value of Z at corner points of the feasible region.

$$\text{Clearly, } Z(O) = 0 ; Z(C) = 3 \times 5 = 15$$

$$Z(E) = 3 \times 4 + 2 \times 3 = 18 \text{ and } Z(B) = 2 \times 5 = 10$$

Thus, maximum value of Z is 18, at point $E(4, 3)$.

$$8. (a) : \text{Let } \vec{u} = x\hat{i} + y\hat{j} + z\hat{k} \Rightarrow (\vec{u} \cdot \hat{i})^2 = x^2$$

$$\text{Similarly, } (\vec{u} \cdot \hat{j})^2 = y^2 \text{ and } (\vec{u} \cdot \hat{k})^2 = z^2$$

$$\therefore (\vec{u} \cdot \hat{i})^2 + (\vec{u} \cdot \hat{j})^2 + (\vec{u} \cdot \hat{k})^2 = x^2 + y^2 + z^2 = |\vec{u}|^2$$

$$9. (b) : \text{Let } I = \int |x| \cdot 1 \, dx$$

$$= |x| \cdot x - \int \frac{|x|}{x} x \, dx + K = x|x| - \int |x| \, dx + K$$

$$\Rightarrow I = x|x| - I + K \Rightarrow 2I = x|x| + K$$

$$\Rightarrow I = \frac{x|x|}{2} + C \left[\text{where } \frac{K}{2} = C \right]$$

10. (d) : Given, $|A^3| = 27 \Rightarrow |A|^3 = 27$

$$\Rightarrow |A| = 3 \Rightarrow \begin{vmatrix} \alpha & 2 \\ 2 & \alpha \end{vmatrix} = 3$$

$$\Rightarrow \alpha^2 - 4 = 3 \Rightarrow \alpha^2 = 7 \Rightarrow \alpha = \pm\sqrt{7}$$

11. (d) : Construct the following table of values of objective function :

Corner Points	Value of $F = 4x + 6y$	
(0, 2)	$4 \times 0 + 6 \times 2 = 12$	} ← Minimum
(3, 0)	$4 \times 3 + 6 \times 0 = 12$	
(6, 0)	$4 \times 6 + 6 \times 0 = 24$	
(6, 8)	$4 \times 6 + 6 \times 8 = 72$	
(0, 5)	$4 \times 0 + 6 \times 5 = 30$	

Since the minimum value (F) = 12 occurs at two distinct corner points, it occurs at every point of the segment joining these two points.

$$12. (a) : e \begin{bmatrix} e^x & e^y \\ e^y & e^x \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \Rightarrow \begin{bmatrix} e^{x+1} & e^{y+1} \\ e^{y+1} & e^{x+1} \end{bmatrix} = \begin{bmatrix} e^0 & e^0 \\ e^0 & e^0 \end{bmatrix}$$

$$\Rightarrow e^{x+1} = e^0 \text{ and } e^{y+1} = e^0 \Rightarrow x = -1 \text{ and } y = -1$$

13. (b) : We have, $P(A) = 1/4$

$$\text{Now, } P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$= P(A) + P(B) - P(A)P(B) \quad (\because A, B \text{ are independent})$$

$$\Rightarrow 1/4 + P(B) - (1/4)P(B) = 2P(B) - 1/4 \quad (\text{Given})$$

$$\Rightarrow P(B) = 2/5$$

14. (c) : Given that matrix A is singular $\Rightarrow |A| = 0$

$$\Rightarrow \begin{vmatrix} 1 & -2 & 3 \\ 1 & 2 & 1 \\ x & 2 & -3 \end{vmatrix} = 0 \Rightarrow 1(-6-2) + 2(-3-x) + 3(2-2x) = 0$$

$$\Rightarrow -8 - 6 - 2x + 6 - 6x = 0 \Rightarrow -8x - 8 = 0 \Rightarrow x = -1$$

$$15. (a) : \text{We have } \frac{dy}{dx} = (1-x)(1+y) \Rightarrow \frac{dy}{1+y} = (1-x)dx$$

$$\Rightarrow \int \frac{dy}{1+y} = \int (1-x)dx \Rightarrow \log|1+y| = x - \frac{x^2}{2} + C$$

16. (c) : Given $y = (A + B) \cos(x + C) + De^x$

$$\text{or } y = k \cos(x + C) + De^x$$

Now order of a differential equation is same as the number of arbitrary unknowns present in the solution. Hence order of differential equation is 3.

17. (b) : Here, $\vec{a} = 3\hat{i} - 2\hat{j} + 6\hat{k} \therefore$ Its magnitude = $|\vec{a}|$

$$= \sqrt{3^2 + (-2)^2 + 6^2} = \sqrt{9 + 4 + 36} = \sqrt{49} = 7.$$

18. (c) : The given equation of line is

$$\frac{4-x}{2} = \frac{y}{6} = \frac{1-z}{3}$$

$$\Rightarrow \frac{x-4}{-2} = \frac{y}{6} = \frac{z-1}{-3} \Rightarrow \frac{x-4}{2} = \frac{y}{-6} = \frac{z-1}{3}$$

$$\text{Now, as } \sqrt{2^2 + (-6)^2 + 3^2} = 7$$

$$\therefore \text{ D.c's. of (i) are } \frac{2}{7}, \frac{-6}{7}, \frac{3}{7}.$$

19. (c) : We know, $0 \leq \cos^{-1}(2x-1) \leq \pi$

$$\Rightarrow -1 \leq 2x-1 \leq 1 \Rightarrow 0 \leq 2x \leq 2 \Rightarrow 0 \leq x \leq 1$$

$$\therefore \text{ Domain of } \cos^{-1}(2x-1) = [0, 1]$$

Also, reason is not true as domain of $\cos^{-1}x$ is $[-1, 1]$.

$\therefore$ A is true but Reason (R) is false.

20. (a) : For two lines, having direction cosines

$$\left(\frac{12}{13}, \frac{-3}{13}, \frac{-4}{13}\right) \text{ and } \left(\frac{4}{13}, \frac{12}{13}, \frac{3}{13}\right), \text{ we obtain}$$

$$l_1 l_2 + m_1 m_2 + n_1 n_2$$

$$= \frac{12}{13} \times \frac{4}{13} + \left(\frac{-3}{13}\right) \times \frac{12}{13} + \left(\frac{-4}{13}\right) \times \frac{3}{13} = 0$$

Therefore, the lines are perpendicular.

So, both A and R are true and R is the correct explanation of A.

$$21. \left[\frac{\sin^{-1}\left(\frac{1}{\sqrt{2}}\right) - 3\sin^{-1}\left(\frac{\sqrt{3}}{2}\right)}{\cos^{-1}\left(\frac{1}{\sqrt{2}}\right) - 3\cos^{-1}\left(\frac{\sqrt{3}}{2}\right)} \right] = \left[\frac{\frac{\pi}{4} - 3\left(\frac{\pi}{3}\right)}{\frac{\pi}{4} - 3\left(\frac{\pi}{6}\right)} \right]$$

$$= \left[\frac{\frac{\pi}{4} - \pi}{\frac{\pi}{4} - \frac{\pi}{2}} \right] = 3$$

$$\left[\therefore \text{ Principal value of } \sin^{-1}\left(\frac{1}{\sqrt{2}}\right) = \frac{\pi}{4}, \sin^{-1}\left(\frac{\sqrt{3}}{2}\right) = \frac{\pi}{3}, \right.$$

$$\left. \cos^{-1}\left(\frac{1}{\sqrt{2}}\right) = \frac{\pi}{4} \text{ and } \cos^{-1}\left(\frac{\sqrt{3}}{2}\right) = \frac{\pi}{6} \right]$$

$$22. \text{ Any point on the line } \frac{x+2}{3} = \frac{y+1}{2} = \frac{z-3}{2} = \lambda (\text{say}) \text{ is}$$

$$\text{of the form } Q(3\lambda - 2, 2\lambda - 1, 2\lambda + 3) \quad \dots(i)$$

Now, distance PQ, where P is (1, 3, 3), is 5.

$$\text{So, } (3\lambda - 2 - 1)^2 + (2\lambda - 1 - 3)^2 + (2\lambda + 3 - 3)^2 = 5^2$$

$$\Rightarrow 9\lambda^2 + 9 - 18\lambda + 4\lambda^2 + 16 - 16\lambda + 4\lambda^2 = 25$$

$$\Rightarrow 17\lambda^2 - 34\lambda = 0 \Rightarrow 17\lambda(\lambda - 2) = 0$$

$$\Rightarrow \lambda = 0 \text{ or } \lambda = 2$$

Putting values of λ in (i), the required points are (-2, -1, 3) and (4, 3, 7).

OR

We have, $\vec{a}$ and $\vec{b}$ are collinear.

$$\therefore \vec{a} = \lambda \vec{b} \Rightarrow -3\hat{i} + n\hat{j} + 4\hat{k} = \lambda(-2\hat{i} + 4\hat{j} + p\hat{k})$$

$$\Rightarrow \lambda = \frac{3}{2}$$

$$\text{Also, } n = 4\lambda \Rightarrow n = 4 \times \frac{3}{2} = 6$$

$$\text{And, } \lambda p = 4 \Rightarrow p = 4 \times \frac{2}{3} = \frac{8}{3}$$

23. We have, L.H.L. (at $x = 0$)

$$= \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0} \frac{\sqrt{1+kx} - \sqrt{1-kx}}{x}$$

$$= \lim_{x \rightarrow 0} \frac{1+kx-1+kx}{x(\sqrt{1+kx} + \sqrt{1-kx})}$$

$$= \lim_{x \rightarrow 0} \frac{2k}{\sqrt{1+kx} + \sqrt{1-kx}} = \frac{2k}{\sqrt{1} + \sqrt{1}} = \frac{2k}{2} = k$$

R.H.L. (at $x = 0$)

$$= \lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0} \frac{2x+1}{x-1} = -1 \text{ and } f(0) = -1$$

Since $f(x)$ is continuous at $x = 0$. $\therefore k = -1$.

24. We have,

$$f(x) = (x-1)^2 + (x-2)^2 + (x-3)^2 + (x-4)^2 + (x-5)^2$$

$$\Rightarrow f'(x) = 2[x-1+x-2+x-3+x-4+x-5]$$

$$= 2(5x-15)$$

$$f'(x) = 0 \Rightarrow 2(5x-15) = 0 \Rightarrow x = 3$$

Also, $f''(x) = 10 > 0$, for all x .

$\therefore f(x)$ is minimum for $x = 3$.

25. Direction ratios of AB are

$$(-1-0, 6-7, 6-10) \text{ or } (-1, -1, -4)$$

$$\text{Also, } \sqrt{(-1)^2 + (-1)^2 + (-4)^2} = 3\sqrt{2}$$

$$\therefore \text{ Direction cosines are } \left(-\frac{1}{3\sqrt{2}}, -\frac{1}{3\sqrt{2}}, -\frac{4}{3\sqrt{2}}\right)$$

$$\text{or } \left(\frac{1}{3\sqrt{2}}, \frac{1}{3\sqrt{2}}, \frac{4}{3\sqrt{2}}\right)$$

OR

The position vector of the point (12, n) is $12\hat{i} + n\hat{j}$.

$$\therefore \vec{a} = 12\hat{i} + n\hat{j} \Rightarrow |\vec{a}| = \sqrt{12^2 + n^2} = 13 \quad (\text{Given})$$

$$\Rightarrow 12^2 + n^2 = 169 \Rightarrow n^2 = 25 \Rightarrow n = \pm 5$$

26. Let E : 'a total of 8' and F : 'red die resulted in a number less than 4'

i.e., $E = \{(2, 6), (3, 5), (4, 4), (5, 3), (6, 2)\}$

and $F = \{(x, y) : x \in \{1, 2, 3, 4, 5, 6\}, y \in \{1, 2, 3\}\}$

i.e., $F = \{(1, 1), (1, 2), (1, 3), (2, 1), (2, 2), (2, 3), (3, 1), (3, 2), (3, 3), (4, 1), (4, 2), (4, 3), (5, 1), (5, 2), (5, 3), (6, 1), (6, 2), (6, 3)\}$

Hence, $E \cap F = \{(5, 3), (6, 2)\}$

$P(E) = 5/36$, $P(F) = 18/36$, $P(E \cap F) = 2/36$

$\therefore$ Required probability = $P(E|F)$

$$= \frac{P(E \cap F)}{P(F)} = \frac{2/36}{18/36} = \frac{2}{18} = \frac{1}{9}$$

27. Let $I = \int \frac{x^2}{(ax+b)^2} dx$

Put $ax + b = t \Rightarrow dx = \frac{1}{a} dt$

$$\therefore I = \frac{1}{a^3} \int \frac{(t-b)^2}{t^2} dt = \frac{1}{a^3} \int \left(1 + \frac{b^2}{t^2} - \frac{2b}{t} \right) dt$$

$$= \frac{1}{a^3} \left(t - \frac{b^2}{t} - 2b \log t \right) + C$$

$$= \frac{1}{a^3} \left(ax + b - \frac{b^2}{ax+b} - 2b \log |ax+b| \right) + C$$

OR

$$\text{Let } I = \int \frac{\sin 2x}{a^2 \sin^2 x + b^2 \cos^2 x} dx$$


$$\text{Let } a^2 \sin^2 x + b^2 \cos^2 x = t \Rightarrow (a^2 - b^2) \sin 2x dx = dt$$

$$\therefore I = \frac{1}{(a^2 - b^2)} \int \frac{1}{t} dt = \frac{1}{(a^2 - b^2)} \log |t| + C$$

$$\Rightarrow I = \frac{1}{(a^2 - b^2)} \log |a^2 \sin^2 x + b^2 \cos^2 x| + C$$

28. Let $l_1 : 2x + 4y = 8$, $l_2 : 3x + y = 6$, $l_3 : x + y = 4$; $x = 0, y = 0$

Solving l_1 and l_2 we get $B\left(\frac{8}{5}, \frac{6}{5}\right)$



Shaded portion $OABC$ is the feasible region, where coordinates of the corner points are $O(0, 0)$, $A(0, 2)$,

$B\left(\frac{8}{5}, \frac{6}{5}\right)$, $C(2, 0)$


The value of objective function at these points are :

Corner Points	Value of the objective function $z = 2x + 5y$
$O(0, 0)$	$2 \times 0 + 5 \times 0 = 0$
$A(0, 2)$	$2 \times 0 + 5 \times 2 = 10$ (Maximum)
$B\left(\frac{8}{5}, \frac{6}{5}\right)$	$2 \times \frac{8}{5} + 5 \times \frac{6}{5} = 9.2$
$C(2, 0)$	$2 \times 2 + 5 \times 0 = 4$

$\therefore$ The maximum value of z is 10, which is at $A(0, 2)$.

OR

Let $l_1 : 2x + 3y = 6$, $l_2 : 3x - 2y = 6$, $l_3 : y = 1$; $x = 0, y = 0$



Solving l_1 and l_3 , we get $D(1.5, 1)$

Solving l_1 and l_2 , we get $C\left(\frac{30}{13}, \frac{6}{13}\right)$

Shaded portion $OADCBO$ is the feasible region, where coordinates of the corner points are $O(0, 0)$, $A(0, 1)$,

$D(1.5, 1)$, $C\left(\frac{30}{13}, \frac{6}{13}\right)$, $B(2, 0)$.

The value of the objective function at these points are :

Corner Points	Value of the objective function $z = 8x + 9y$
$O(0, 0)$	$8 \times 0 + 9 \times 0 = 0$
$A(0, 1)$	$8 \times 0 + 9 \times 1 = 9$
$D(1.5, 1)$	$8 \times 1.5 + 9 \times 1 = 21$
$C\left(\frac{30}{13}, \frac{6}{13}\right)$	$8 \times \frac{30}{13} + 9 \times \frac{6}{13} = 22.6$ (Maximum)
$B(2, 0)$	$8 \times 2 + 9 \times 0 = 16$

The maximum value of z is 22.6, which is at $C\left(\frac{30}{13}, \frac{6}{13}\right)$

CBSE CHAPTERWISE TOPICWISE 10 SOLVED PAPERS (2022-2013)



CBSE CHAMPION Chapterwise-Topicwise Solved Papers Series contains topicwise questions and solutions asked over last decade in CBSE-Board examination.

Examination papers for Class 10 and 12 Boards are based on a certain pattern. To excel, studying right is therefore more important than studying hard, which is why we created this series.

Questions are supported with topicwise graphical analysis of previous years CBSE Board questions as well as comprehensive and lucid theory. Questions from Term I, Term II, Delhi, All India and Foreign papers are included.

Important feature of these books is that the solutions to all the questions have been given according to CBSE marking scheme with Preparatory Tools and Topper's Answers. Practice papers are also supplemented.

HIGHLIGHTS OF 2022-23 EDITION

- Based on Reduced Syllabus
- Topicwise Graphical Analysis
- Quick Recap with Brain Map
- Previous 10 years (2022-2013) questions of CBSE
- CBSE Cognitive Level Tagging
- Key Points, Answer Tips, Concept Applied, Shortcuts, Alternating Methods, Commonly Made Mistakes provided
- CBSE Topper's Answers
- Past 3 years CBSE Sample Question Papers
- Chapterwise Self-Assessment
- 5 Practice Papers
- CBSE Sample Paper (Issued on 16th September 2022)

LATEST
2022-23
PRACTICE PAPERS
INCLUDED



Available at all leading book shops throughout India.
For more information or for help in placing your order:
Call 0124-6601200 or email info@mtg.in
Visit www.mtg.in for latest offers and to buy online!

29. Given, $\frac{dy}{dx} = \frac{x^2 + y^2 + 1}{2xy}$
 $\Rightarrow 2xydy = (x^2 + y^2 + 1)dx$
 $\Rightarrow 2xydy - y^2dx = (x^2 + 1)dx$
 $\Rightarrow xd(y^2) - y^2dx = (x^2 + 1)dx$
 $\Rightarrow \frac{xd(y^2) - y^2dx}{x^2} = \left(1 + \frac{1}{x^2}\right)dx \Rightarrow d\left(\frac{y^2}{x}\right) = d\left(x - \frac{1}{x}\right)$

Integrating both sides, we get

$$\frac{y^2}{x} = x - \frac{1}{x} + C \Rightarrow y^2 = x^2 - 1 + Cx$$

Now, given that $y(1) = 1$

$$\therefore 1 = 1 - 1 + C \Rightarrow C = 1$$

Thus, curve becomes $y^2 = x^2 - 1 + x$

OR

We have, $y = cx + \frac{a}{c}$... (i)

Differentiating with respect to x , we get $\frac{dy}{dx} = c$

$$\therefore x \frac{dy}{dx} + \frac{a}{c} = xc + \frac{a}{c} \quad \left[\text{Putting } \frac{dy}{dx} = c \right]$$

$$\Rightarrow x \frac{dy}{dx} + \frac{a}{c} = y \quad [\text{Using (i)}]$$

Hence, $y = cx + \frac{a}{c}$ is a solution of the differential equation $x \frac{dy}{dx} + \frac{a}{dy/dx} = y$.

30. Let $I = \int \frac{dx}{\sqrt{1 - (x^2 + 2x)}} = \int \frac{dx}{\sqrt{2 - (x^2 + 2x + 1)}}$
 $= \int \frac{dx}{\sqrt{2 - (1 + x)^2}} = \int \frac{dx}{\sqrt{(\sqrt{2})^2 - (1 + x)^2}}$

Let $1 + x = z \Rightarrow dx = dz$

$$\therefore I = \int \frac{dz}{\sqrt{(\sqrt{2})^2 - z^2}} = \sin^{-1} \frac{z}{\sqrt{2}} + c = \sin^{-1} \left(\frac{1+x}{\sqrt{2}} \right) + c$$

31. Let $I = \int \frac{x^2 + 9}{x^4 + 81} dx \Rightarrow I = \int \frac{1 + (9/x^2)}{x^2 + \frac{81}{x^2}} dx$
 $\Rightarrow I = \int \frac{1 + (9/x^2)}{x^2 + \left(\frac{9}{x}\right)^2 - 18 + 18} dx = \int \frac{1 + (9/x^2)}{\left(x - \frac{9}{x}\right)^2 + 18} dx$

Let $x - \frac{9}{x} = t \Rightarrow \left(1 + \frac{9}{x^2}\right) dx = dt$

$$\therefore I = \int \frac{dt}{t^2 + 18} \Rightarrow I = \int \frac{dt}{t^2 + (3\sqrt{2})^2}$$

$$\Rightarrow I = \frac{1}{3\sqrt{2}} \tan^{-1} \left(\frac{t}{3\sqrt{2}} \right) + c$$

$$\Rightarrow I = \frac{1}{3\sqrt{2}} \tan^{-1} \left(\frac{x^2 - 9}{3\sqrt{2}x} \right) + c$$

32. Given, $y = |x + 1| + 1$

$$\therefore y = \begin{cases} x + 2, & \text{if } x \geq -1 \\ -x, & \text{if } x < -1 \end{cases}$$

We now draw the lines : $y = 0, x = 3, x = -3$ and

$$y = x + 2 \text{ if } x \geq -1 \quad \dots (i)$$

$$y = -x \text{ if } x < -1 \quad \dots (ii)$$


Lines (i) and (ii) intersect each other at $(-1, 1)$

$\therefore$ Required area

$$= \int_{-3}^{-1} (-x) dx + \int_{-1}^3 (x + 2) dx$$

$$= -\left[\frac{x^2}{2}\right]_{-3}^{-1} + \left[\frac{x^2}{2} + 2x\right]_{-1}^3$$

$$= -\frac{1}{2}(1 - 9) + \frac{1}{2}(9 - 1) + 2(3 + 1) = 4 + 4 + 8 = 16 \text{ sq. units.}$$



33. Given lines are

$$l_1 : \vec{r} = \hat{i} + 2\hat{j} - 4\hat{k} + \lambda(2\hat{i} + 3\hat{j} + 6\hat{k});$$

$$l_2 : \vec{r} = 3\hat{i} + 3\hat{j} - 5\hat{k} + \mu(4\hat{i} + 6\hat{j} + 12\hat{k})$$

$$\therefore \text{We have } \vec{a}_1 = \hat{i} + 2\hat{j} - 4\hat{k}, \vec{b}_1 = 2\hat{i} + 3\hat{j} + 6\hat{k}$$

$$\text{and } \vec{a}_2 = 3\hat{i} + 3\hat{j} - 5\hat{k}, \vec{b}_2 = 4\hat{i} + 6\hat{j} + 12\hat{k}$$

$$\text{So, } \vec{a}_2 - \vec{a}_1 = 2\hat{i} + \hat{j} - \hat{k}$$

$$\text{Also, } \vec{b}_2 = 4\hat{i} + 6\hat{j} + 12\hat{k} = 2\vec{b}_1 \Rightarrow \vec{b}_1 \parallel \vec{b}_2$$

Hence l_1 and l_2 are parallel lines.

Shortest distance between two parallel lines is,

$$d = \left| \frac{\vec{b} \times (\vec{a}_2 - \vec{a}_1)}{|\vec{b}|} \right| \Rightarrow d = \left| \frac{(2\hat{i} + 3\hat{j} + 6\hat{k}) \times (2\hat{i} + \hat{j} - \hat{k})}{\sqrt{2^2 + 3^2 + 6^2}} \right|$$

$$\Rightarrow d = \left| \frac{-9\hat{i} + 14\hat{j} - 4\hat{k}}{7} \right|$$

$$\Rightarrow d = \frac{\sqrt{(-9)^2 + 14^2 + (-4)^2}}{7} = \frac{\sqrt{293}}{7} \text{ units.}$$

OR

$$\text{Here, } \vec{a} = 3\hat{i} - \hat{j}, \vec{b} = 2\hat{i} + \hat{j} - 3\hat{k}$$

We have to express : $\vec{b} = \vec{b}_1 + \vec{b}_2$, where

$$\vec{b}_1 \parallel \vec{a} \text{ and } \vec{b}_2 \perp \vec{a}$$

Let $\vec{b}_1 = \lambda \vec{a} = \lambda(3\hat{i} - \hat{j})$ and $\vec{b}_2 = x\hat{i} + y\hat{j} + z\hat{k}$

Now $\vec{b}_2 \perp \vec{a} \Rightarrow \vec{b}_2 \cdot \vec{a} = 0$

$$\Rightarrow (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (3\hat{i} - \hat{j}) = 0 \Rightarrow 3x - y = 0 \quad \dots(i)$$

Now, $\vec{b} = \vec{b}_1 + \vec{b}_2$

$$\Rightarrow 2\hat{i} + \hat{j} - 3\hat{k} = \lambda(3\hat{i} - \hat{j}) + (x\hat{i} + y\hat{j} + z\hat{k})$$

On comparing, we get

$$\begin{cases} 2 = 3\lambda + x \\ 1 = -\lambda + y \end{cases} \Rightarrow x + 3y = 5$$

and $-3 = z \Rightarrow z = -3$

Solving (i) and (ii), we get $x = \frac{1}{2}, y = \frac{3}{2}$

$$\therefore 1 = -\lambda + y = -\lambda + \frac{3}{2} \Rightarrow \lambda = \frac{1}{2}$$

Hence, $\vec{b}_1 = \lambda(3\hat{i} - \hat{j}) = \frac{3}{2}\hat{i} - \frac{1}{2}\hat{j}$ and $\vec{b}_2 = \frac{1}{2}\hat{i} + \frac{3}{2}\hat{j} - 3\hat{k}$

$$34. \text{ Let } A = \begin{bmatrix} 3 & -2 & -4 \\ 3 & -2 & -5 \\ -1 & 1 & 2 \end{bmatrix} \Rightarrow A' = \begin{bmatrix} 3 & 3 & -1 \\ -2 & -2 & 1 \\ -4 & -5 & 2 \end{bmatrix}$$

$$\therefore A + A' = \begin{bmatrix} 3 & -2 & -4 \\ 3 & -2 & -5 \\ -1 & 1 & 2 \end{bmatrix} + \begin{bmatrix} 3 & 3 & -1 \\ -2 & -2 & 1 \\ -4 & -5 & 2 \end{bmatrix}$$

$$= \begin{bmatrix} 6 & 1 & -5 \\ 1 & -4 & -4 \\ -5 & -4 & 4 \end{bmatrix}$$

$$\Rightarrow \frac{1}{2}(A + A') = \begin{bmatrix} 3 & \frac{1}{2} & -\frac{5}{2} \\ \frac{1}{2} & -2 & -2 \\ -\frac{5}{2} & -2 & 2 \end{bmatrix}$$

which is clearly a symmetric matrix and

$$A - A' = \begin{bmatrix} 3 & -2 & -4 \\ 3 & -2 & -5 \\ -1 & 1 & 2 \end{bmatrix} - \begin{bmatrix} 3 & 3 & -1 \\ -2 & -2 & 1 \\ -4 & -5 & 2 \end{bmatrix} = \begin{bmatrix} 0 & -5 & -3 \\ 5 & 0 & -6 \\ 3 & 6 & 0 \end{bmatrix}$$

$$\Rightarrow \frac{1}{2}(A - A') = \begin{bmatrix} 0 & -\frac{5}{2} & -\frac{3}{2} \\ \frac{5}{2} & 0 & -3 \\ \frac{3}{2} & 3 & 0 \end{bmatrix}$$

which is clearly a skew symmetric matrix.

Since $A = \frac{1}{2}(A + A') + \frac{1}{2}(A - A')$

$$\therefore A = \begin{bmatrix} 3 & \frac{1}{2} & -\frac{5}{2} \\ \frac{1}{2} & -2 & -2 \\ -\frac{5}{2} & -2 & 2 \end{bmatrix} + \begin{bmatrix} 0 & -\frac{5}{2} & -\frac{3}{2} \\ \frac{5}{2} & 0 & -3 \\ \frac{3}{2} & 3 & 0 \end{bmatrix}$$

Thus A has been expressed as the sum of a symmetric matrix and a skew-symmetric matrix.

35. Here, $f: A \rightarrow B$ is given by $f(x) = \frac{x-1}{x-2}$, where

$A = R - \{2\}$ and $B = R - \{1\}$

Let $f(x_1) = f(x_2)$, where $x_1, x_2 \in A$ (i.e., $x_1 \neq 2, x_2 \neq 2$)

$$\Rightarrow \frac{x_1-1}{x_1-2} = \frac{x_2-1}{x_2-2}$$

$$\Rightarrow (x_1-1)(x_2-2) = (x_1-2)(x_2-1)$$

$$\Rightarrow x_1x_2 - 2x_1 - x_2 + 2 = x_1x_2 - x_1 - 2x_2 + 2$$

$$\Rightarrow -2x_1 - x_2 = -x_1 - 2x_2 \Rightarrow x_1 = x_2 \Rightarrow f \text{ is one-one.}$$

Let $y \in B = R - \{1\}$ i.e., $y \in R$ and $y \neq 1$

such that $f(x) = y$

$$\Leftrightarrow \frac{x-1}{x-2} = y \Leftrightarrow (x-2)y = x-1$$

$$\Leftrightarrow xy - 2y = x - 1 \Leftrightarrow x(y-1) = 2y-1 \Leftrightarrow x = \frac{2y-1}{y-1}$$

$$\therefore f(x) = y \text{ when } x = \frac{2y-1}{y-1} \in A \text{ (as } y \neq 1)$$

Hence, f is onto. Thus, f is one-one and onto.

OR

Given relation is $R = \{(a, b): a \leq b^2\}$

Reflexivity: Let $a \in \text{real numbers.}$

$$aRa \Rightarrow a \leq a^2$$

but if $a < 1$, then $a \not\leq a^2$

For example, $a = \frac{1}{2} \Rightarrow a^2 = \frac{1}{4}$ so, $\frac{1}{2} \not\leq \frac{1}{4}$

UNSCRAMBLED WORDS

FEBRUARY 2023

1-i- EXPANSION

2-h- DEVIATION

3-g- FRACTION

4-f- PARALLELEPIPED

5-a- SLOPE

6-j- PREDECESSOR

7-e- TRAPEZIUM

8-d- ALGEBRA

9-c- STATISTICS

10-b- SPHERE

Hence, R is not reflexive.

Symmetry : $aRb \Rightarrow a \leq b^2$

But then $b \leq a^2$ is not true $\therefore aRb \not\Rightarrow bRa$

For example, $a = 2, b = 5$ then $2 \leq 5^2$ but $5 \leq 2^2$ is not true.

Hence, R is not symmetric.

Transitivity : Let $a, b, c \in$ real numbers

Considering aRb and bRc

$$aRb \Rightarrow a \leq b^2 \text{ and } bRc \Rightarrow b \leq c^2 \Rightarrow a \leq c^4 \not\Rightarrow aRc$$

For example, if $a = 2, b = -3, c = 1$

$$aRb \Rightarrow 2 \leq 9; bRc \Rightarrow -3 \leq 1; aRc \Rightarrow 2 \leq 1 \text{ is not true.}$$

Hence, R is not transitive

36. Let E_1 be the event that Ravi solved the puzzle and E_2 be the event that Priya solved the puzzle.

Then, $P(E_1) = 1/4$ and $P(E_2) = 1/5$

(i) Since, E_1 and E_2 are independent events.

$$\therefore P(\text{both solved the puzzle}) = P(E_1 \cap E_2)$$

$$= P(E_1) \cdot P(E_2) = \frac{1}{4} \times \frac{1}{5} = \frac{1}{20} = \frac{1}{20} \times 100\% = 5\%$$

(ii) $P(\text{puzzle is solved by Ravi but not by Priya})$

$$= P(\bar{E}_2)P(E_1) = \left(1 - \frac{1}{5}\right) \cdot \frac{1}{4} = \frac{4}{5} \cdot \frac{1}{4} = \frac{1}{5}.$$

(iii) $P(\text{puzzle is solved}) = P(E_1 \cup E_2)$

$$= P(E_1 \cup E_2) = P(E_1) + P(E_2) - P(E_1 \cap E_2)$$

$$= \frac{1}{4} + \frac{1}{5} - \frac{1}{20} = \frac{8}{20} = \frac{2}{5}.$$

OR

$P(\text{Exactly one of them solved the puzzle})$

$$= P[(E_1 \text{ and } \bar{E}_2) \text{ or } (E_2 \text{ and } \bar{E}_1)]$$

$$= P(E_1 \cap \bar{E}_2) + P(E_2 \cap \bar{E}_1)$$

$$= P(E_1) \times P(\bar{E}_2) + P(E_2) \times P(\bar{E}_1)$$

$$= \frac{1}{4} \times \frac{4}{5} + \frac{1}{5} \times \frac{3}{4} = \frac{4}{20} + \frac{3}{20} = \frac{7}{20}. \quad [\because P(\bar{E}_1) = 1 - P(E_1)]$$

37. (i) Let F be the fuel cost per hour and v be the speed of train in km/hr.

According to question, we have,

$F \propto v^2 \Rightarrow F = kv^2$, where k is proportionality constant

$$\Rightarrow 48 = k(16)^2 \Rightarrow k = \frac{3}{16}$$

(ii) Let total cost of running the train be C .

$$\text{Then, } C = \frac{3}{16}v^2t + 1200t$$

$$\text{Now, distance covered} = 500 \text{ km} \Rightarrow \text{Time} = \frac{500}{v} \text{ hrs}$$

$\therefore$ Total cost of running the train for 500 km

$$= \frac{3}{16}v^2\left(\frac{500}{v}\right) + 1200\left(\frac{500}{v}\right) \Rightarrow C = \frac{375}{4}v + \frac{600000}{v}$$

(iii) We have, $\frac{dC}{dv} = \frac{375}{4} - \frac{600000}{v^2}$

$$\text{Put } \frac{dC}{dv} = 0 \Rightarrow v^2 = \frac{600000 \times 4}{375} = 6400 \Rightarrow v = 80 \text{ km/h}$$

$$\frac{d^2C}{dv^2} = \frac{2 \times 600000}{v^3} > 0, \text{ for } v = 80$$

$\therefore$ Most economical speed is 80 km/h.

OR

Fuel cost for running the train for 500 km

$$= \frac{3}{16}v^2\left(\frac{500}{v}\right) = \frac{375}{4}v = \frac{375}{4} \times 80 = ₹ 7500$$

38. (i) Given, perimeter of window = 10 m

$\therefore x + y + y + \text{perimeter of semicircle} = 10$

$$\Rightarrow x + 2y + \pi \frac{x}{2} = 10 \Rightarrow y = 5 - \frac{x}{2} - \frac{\pi x}{4} \quad \dots(i)$$

$$A = x \cdot y + \frac{1}{2} \pi \left(\frac{x}{2}\right)^2 = x \left(5 - \frac{x}{2} - \frac{\pi x}{4}\right) + \frac{1}{2} \frac{\pi x^2}{4} \quad [\because \text{From (i)}]$$

$$= 5x - \frac{x^2}{2} - \frac{\pi x^2}{4} + \frac{\pi x^2}{8} = 5x - \frac{x^2}{2} - \frac{\pi x^2}{8}$$

(ii) We have, $A = 5x - \frac{x^2}{2} - \frac{\pi x^2}{8} \Rightarrow \frac{dA}{dx} = 5 - x - \frac{\pi x}{4}$

$$\text{Now, } \frac{dA}{dx} = 0 \Rightarrow 5 = x + \frac{\pi x}{4}$$

$$\Rightarrow x(4 + \pi) = 20 \Rightarrow x = \frac{20}{4 + \pi}$$

$$\left[\text{Clearly, } \frac{d^2A}{dx^2} < 0 \text{ at } x = \frac{20}{4 + \pi} \right]$$

$$\text{At } x = \frac{20}{4 + \pi}$$

$$A = 5 \left(\frac{20}{4 + \pi} \right) - \left(\frac{20}{4 + \pi} \right)^2 \frac{1}{2} - \frac{\pi}{8} \left(\frac{20}{4 + \pi} \right)^2$$

$$= \frac{100}{4 + \pi} - \frac{200}{(4 + \pi)^2} - \frac{50\pi}{(4 + \pi)^2}$$

$$= \frac{(4 + \pi)(100) - 200 - 50\pi}{(4 + \pi)^2} = \frac{400 + 100\pi - 200 - 50\pi}{(4 + \pi)^2}$$

$$= \frac{200 + 50\pi}{(4 + \pi)^2} = \frac{50(4 + \pi)}{(4 + \pi)^2} = \frac{50}{4 + \pi}$$





Unlock Your Knowledge!




1. Find the projection of $\vec{a}$ on $\vec{b}$, if $\vec{a} \cdot \vec{b} = 8$ and $\vec{b} = 2\hat{i} + 6\hat{j} + 3\hat{k}$.
2. Write the number of all the one-one functions from $A = \{1, 2, 3\}$ to itself.
3. If $\begin{bmatrix} a+b & 2 \\ 5 & ab \end{bmatrix} = \begin{pmatrix} 6 & 2 \\ 5 & 8 \end{pmatrix}$, then find the value of $a + b$.
4. If A and B are disjoint sets and A' is the complement of A , then find the value of $B \cap A'$.
5. Find the value of x if $\cos 2x = \cos x$ for $0 \leq x \leq \pi$.
6. Find the value of $7! - 5!$.
7. The arithmetic mean of two numbers a and b is 3 and geometric mean is 2. Find the value of $a^2 + b^2$.
8. Find the last digit of $(1! + 2! + 3! + \dots + 2005!)^{502}$.
9. Write the derivative of $f(x) = \cos x$ at $x = 0$.
10. If A and B are square matrices of the same order and if $A = A^T$, $B = B^T$, then find the value of the transpose of (ABA) .
11. What is the equation of tangent to $y = -2x^2 + 3$ at $x = 1$?
12. Find the value of $\int_e^{e^2} \frac{1}{x} dx$.
13. If $0 \leq A$, $B \leq \frac{\pi}{2}$ and $\sin A + \cos B = 2$, then find the value of $\tan\left(\frac{A+B}{2}\right)$.
14. If $\tan(x+y) = 33$ and $x = \tan^{-1}3$, then what is the value of y ?
15. Find the value of $f[f(x)]$ if $f(x) = \frac{x+2}{3x-1}$.
16. Find the value of $2b + c$, for the function $f(x) = 2x^3 + bx^2 + cx$, which holds Rolle's theorem for $x \in [-1, 1]$ at $x = 1/2$.
17. Find the probability of showing at least one head if a pair of coin is tossed once.
18. What will be value of $2y(2) - y(1)$ for $\frac{dy}{dx} + \frac{y}{x} = x^2$?
19. Find standard deviation of four observations if their mean is 3 and sum of the squares of these observations is 48.
20. If 3 out of 6 vertices of a regular hexagon are chosen at a time at random, then what will be the probability that the triangle with three vertices is an equilateral triangle?

Readers can send their responses at editor@mtg.in or post us with complete address by 10th of every month. Winners' names and answers will be published in next issue.

SOLUTIONS TO FEBRUARY 2023 QUIZ CLUB

- | | |
|-----------------------------------|--|
| 1. $10C^3$ | 13. $\frac{1}{2}$ |
| 2. $6 + 11i$ | 14. A |
| 3. 7 units | 15. $\frac{-4x}{1-x^4}$ |
| 4. $2 \cos a$ | 16. $\tan^{-1} \sqrt{\frac{p}{q}}$ |
| 5. 671 | 17. $\frac{7}{3}$ |
| 6. $2 + \sqrt{3}$ | 18. $\tan(y/x) = cx$ |
| 7. 0.6 | 19. $\frac{5\sqrt{3}}{3}(\hat{i} + \hat{j} + \hat{k})$ |
| 8. $\frac{4}{\sqrt{5}-1}$ | 20. 90° |
| 9. A | |
| 10. $x = y$ | |
| 11. $3f(x)$ | |
| 12. $2(x^2y^2 + y^2z^2 + z^2x^2)$ | |

MONTHLY TEST DRIVE



This specially designed column enables students to self analyse their extent of understanding of specified chapters. Give yourself four marks for correct answer and deduct one mark for wrong answer. Self check table given at the end will help you to check your readiness.

Total Marks : 80

Series 8 : Differential Equations

Time Taken : 60 Min.

Only One Option Correct Type

1. If p and q are order and degree of differential equation

$$y^2 \left(\frac{d^2 y}{dx^2} \right)^2 + 3x \left(\frac{dy}{dx} \right)^{1/3} + x^2 y^2 = \sin x, \text{ then}$$

- (a) $p > q$ (b) $\frac{p}{q} = \frac{1}{2}$
(c) $p = q$ (d) $p < q$

2. Solve : $\left(x \sin \frac{y}{x} \right) dy = \left(y \sin \frac{y}{x} - x \right) dx$.

- (a) $\cos \frac{y}{x} = \log x + c$ (b) $\sin \frac{y}{x} = \log x + c$
(c) $\cot x = \log y + c$ (d) None of these

3. The differential equation of all non-vertical lines in a plane, is

- (a) $\frac{d^2 y}{dx^2} = 0$ (b) $\frac{d^2 x}{dy^2} = 0$
(c) $\frac{dy}{dx} = 0$ (d) $\frac{dx}{dy} = 0$

4. The solution of the differential equation $e^{-x}(y+1)dy + (\cos^2 x - \sin 2x)y dx = 0$ subjected to the condition that $y = 1$ when $x = 0$ is

- (a) $(y+1) + e^x \cos^2 x = 2$
(b) $y + \log y = e^x \cos^2 x$
(c) $\log(y+1) + e^x \cos^2 x = 1$
(d) $y + \log y + e^x \cos^2 x = 2$

5. The equation of a curve passing through $(0, 1)$ and

having gradient $\frac{-(y+y^3)}{1+x+xy^2}$ at (x, y) is

- (a) $xy + \tan^{-1} y = \frac{\pi}{2}$ (b) $xy + \tan^{-1} y = \frac{\pi}{4}$

(c) $xy - \tan^{-1} y = \frac{\pi}{2}$ (d) $xy - \tan^{-1} y = \frac{\pi}{4}$

6. If $ye^y dx = (y^3 + 2xe^y) dy$, $y(0) = 1$, then the value of x when $y = 0$ is

- (a) -1 (b) 0 (c) 1 (d) 2

One or More Than One Option(s) Correct Type

7. If the differential equation of the family of all circles whose centre lies on the straight line $y = x$ is represented by the $P \frac{d^2 y}{dx^2} + Q \frac{dy}{dx} + 1 = 0$, where P & Q are functions of x & y , then which of the following is/are true?

- (a) $P = y + x$ (b) $P = y - x$

(c) $P + Q = 1 - x + y + \frac{dy}{dx} + \left(\frac{dy}{dx} \right)^2$

(d) $P - Q = x + y - \frac{dy}{dx} - \left(\frac{dy}{dx} \right)^2$

8. The differential equation whose solution represents the family $y = Ae^{3x} + Be^{-7x}$ is $y_2 + \lambda y_1 + my = 0$, then

- (a) $\lambda + |m| = 25$
(b) Number of prime factors of $|m|$ is 2
(c) $\lambda = 16$
(d) If λ is represented as exponent of a prime number then base is smallest prime.

9. If the solution of the equation $\frac{d^2 x}{dt^2} + 4 \frac{dx}{dt} + 13x = 0$,

given that for $t = 0$, $x = 1$ and $\frac{dx}{dt} = 10$ is in the form

$x = e^{-2t}(A \cos 3t + B \sin 3t)$ then

- (a) $A + B = 5$ (b) $AB = 4$
(c) $AB = 5$ (d) None of these

10. Which of the following statement(s) is/are true?

- (a) $f(x, y) = e^{y/x} + \tan \frac{y}{x}$ is a homogeneous equation.
 (b) $x \ln \frac{y}{x} dx + \frac{y^2}{x} \sin^{-1} \frac{y}{x} dy = 0$ is a homogeneous differential equation.
 (c) $f(x, y) = x^2 + \sin x \cos y$ is a non-homogeneous equation.
 (d) $(x^2 + y^2)dx - (xy^2 - y^3)dy = 0$ is a homogeneous differential equation.

11. The equation of a curve passing through $(2, 7/2)$ and having gradient $1 - 1/x^2$ at (x, y) is $xy = ax^2 + bx + c$, then

- (a) $a = b = c$ (b) $abc = 1$
 (c) $a + b + c = 3$ (d) $2a + b = 4$

12. Which of the following equation(s) is/are linear?

- (a) $\frac{dy}{dx} + \frac{y}{x} = \log x$ (b) $y \left(\frac{dy}{dx} \right) + 4x = 0$
 (c) $(2x + y^3) \left(\frac{dy}{dx} \right) = 3y$ (d) None of these

13. If $y = a \sin(bx + c)$ be the solution of a differential equation, then

- (a) $y \frac{d^3 y}{dx^3} = \left(\frac{dy}{dx} \right) \left(\frac{d^2 y}{dx^2} \right)$
 (b) order of differential equation is 3.
 (c) degree of differential equation is 1.
 (d) $y^2 \frac{d^2 y}{dx^2} = \left(\frac{dy}{dx} \right)^3$

Comprehension Type

Paragraph for Q. No. 14 and 15

Differential equation $\frac{dy}{dx} = f(x)g(y)$ can be solved by separating the variables i.e., $\frac{dy}{g(y)} = f(x)dx$.

14. Solution of the differential equation

$$\frac{dy}{dx} + \frac{1+y^2}{\sqrt{1-x^2}} = 0 \text{ is}$$

- (a) $\tan^{-1} y + \sin^{-1} x = C$ (b) $\tan^{-1} x + \sin^{-1} y = C$
 (c) $\tan^{-1} y \sin^{-1} x = C$ (d) $\tan^{-1} y - \sin^{-1} x = C$

15. If $dy/dx = 1 + x + y + xy$ and $y(-1) = 0$, then $y =$

- (a) $e^{\frac{(1-x)^2}{2}}$ (b) $e^{\frac{(1+x)^2}{2}} - 1$
 (c) $\log(1+x) - 1$ (d) $1+x$

Matrix Match Type

16. Match the following :

	Column - I		Column - II
(P)	The solution of $\frac{dy}{dx} = 2^{y-x}$ is	(1)	$x^2(y+3)^3 = e^{y+2}$
(Q)	The solution of $\frac{dy}{dx} = 1 + x + y^2 + xy^2$, when $y = 0, x = 0$ is	(2)	$2^{-x} - 2^{-y} = k$
(R)	Solution of $x^2 \frac{dy}{dx} = x^2 + xy + y^2$ is	(3)	$y = \tan \left(x + \frac{x^2}{2} \right)$
(S)	The solution of $2(y+3) - xy \frac{dy}{dx} = 0$, given that $y(1) = -2$ is	(4)	$\tan^{-1} \left(\frac{y}{x} \right) = \log x + C$

	P	Q	R	S
(a)	2	3	1	4
(b)	3	2	4	1
(c)	3	2	1	4
(d)	2	3	4	1

Numerical Answer Type

17. If $x \frac{dy}{dx} = x^2 + y - 2$, $y(1) = 1$, then $y(-1)$ equals ____.

18. If $\frac{dy}{dx} = e^{-2y}$ and $y = 0$, when $x = e$, then value of x , when $y = \frac{1}{2}$ is $\frac{3e-1}{2k}$. Find the value of k .

19. Order of the differential equation whose solution is $xy = ae^x + be^{-x} + c$ is ____.

20. Number of straight lines which satisfy the differential equation $\frac{dy}{dx} + x \left(\frac{dy}{dx} \right)^2 - y = 0$ is ____.



Keys are published in this issue. Search now! ☺

SELF CHECK

No. of questions attempted
 No. of questions correct
 Marks scored in percentage

Check your score! If your score is

> 90%	EXCELLENT WORK !	You are well prepared to take the challenge of final exam.
90-75%	GOOD WORK !	You can score good in the final exam.
74-60%	SATISFACTORY !	You need to score more next time.
< 60%	NOT SATISFACTORY!	Revise thoroughly and strengthen your concepts.

mtg

Add Little 'Extra' to your Knowledge
to Become 'Extraordinary'

INDIA'S
NO.1

MONTHLY MAGAZINE FOR

NEET | JEE | CUET | CBSE Boards

Get Crystal Clear Concepts and
Best of Practice Material Every Month



@75/- each
100 pages in each



SUBSCRIBE TO GET UP TO 64% DISCOUNT

Subscription Form

Yes! I wish to Subscribe 'MTG Magazine'.

Please tick your subscription scheme and send this form to **MTG Learning Media (P) Ltd.**,
Plot 99, Sector 44, Gurugram-122 003 (HR). Enclose Demand Draft favouring.

SELECT SUBSCRIPTION PLAN

Subject	Format	☐ 1 Year	☐ 2 Years	☐ 3 Years
Individual ☐ Physics ☐ Chemistry ☐ Mathematics ☐ Biology	☐ Print issue	900 ₹ 749/- (Save 17%)	1800 ₹ 1349/- (Save 25%)	2700 ₹ 1799/- (Save 33%)
	☐ Print + Digital	1000 ₹ 1150/- (Save 36%)	3600 ₹ 1850/- (Save 49%)	5400 ₹ 2400/- (Save 56%)
Combo of 3 Magazines ☐ Physics, Chemistry, Biology ☐ Physics, Chemistry, Mathematics	☐ Print issue	2700 ₹ 2200/- (Save 19%)	5400 ₹ 4000/- (Save 26%)	8100 ₹ 5400/- (Save 33%)
	☐ Print + Digital	5400 ₹ 3000/- (Save 44%)	10800 ₹ 5000/- (Save 54%)	16200 ₹ 6600/- (Save 59%)
Combo of 4 Magazines ☐ Physics, Chemistry, Mathematics, Biology	☐ Print issue	3600 ₹ 2800/- (Save 22%)	7200 ₹ 5200/- (Save 28%)	10800 ₹ 6300/- (Save 42%)
	☐ Print + Digital	7200 ₹ 3800/- (Save 47%)	14400 ₹ 6399/- (Save 56%)	21600 ₹ 7800/- (Save 64%)
Courier Charges Add to your subscription amount for quicker & traceable delivery		₹ 800/-	₹ 960/-	₹ 1440/-



Scan QR CODE or
visit this link
bit.ly_3VUBMH
**SUBSCRIBE MAGAZINE
ONLINE**

Note: Magazines are dispatched by Book-Post on 4th of every month (each magazine separately)

- ☐ Student
☐ Teacher
☐ Coaching
☐ School Library

Name: _____

Complete Postal Address: _____

Pin Code: _____

Mobile No.: _____ Email: _____

For more details about the offer, you may contact us:

0124-6601200

8800255335


subscription@mtg.in

www.mtg.in




The Best tool for success in JEE Main


21 Years' JEE Main Chapterwise-Topicwise Solutions Physics, Chemistry and Mathematics contain not only chapterwise - topicwise questions that have appeared over the last 21 years in JEE Main / AIEEE but also their complete solutions. Needless to say these books are essential for any student to compete successfully in JEE Main.



₹ 800/-



₹ 800/-



₹ 800/-

HIGHLIGHTS:

- Chapterwise -Topicwise Questions (Online & Offline) of last 21 years' (2022-2002) of JEE Main / AIEEE
- Chapterwise -Topicwise segregation of questions to help you assess the level of effort required to succeed
- An unmatched series with more than 1740 pages filled with detailed solutions by experts

Visit
www.mtg.in
for latest offers
and to buy
online!



MTG Learning Media (P) Ltd.
Plot #99, Sector 44, Gurugram – 122 003 (HR)

Available at all leading book shops throughout India.
For more information or for help in placing your order,
Call 0124-6601200 or e-mail: info@mtg.in




Scan to explore & buy more
JEE Books on mtg.in


Find all your
QUESTIONS here,
before getting them
in your exam.

By not studying
MORE but
ACCURATE...

Our Books



₹ 900/-



₹ 900/-



₹ 900/-

JEE 2023 exams are nearing and all the students must be in the stage of preparation with full dedication and hard work. But have you put your knowledge to test? With our 41 years of experience, we would suggest you to put yourself through MTG's JEE Champion Series so that you can be sure about your level of preparation and find your core strengths and capabilities. The best part is you study at a pace you're comfortable with. Because it's all chapters, topicwise.




Scan to
explore & buy more
JEE Books on mtg.in




Some of the best lessons are learnt from history!


Introducing Chapterwise -Topicwise Solutions to JEE.
Crack the past to see if you will last.



₹ 900



₹ 900



₹ 900

Tune. Fine tune. Reach the peak of your readiness for JEE with MTG's 45+21 Years Chapterwise-Topicwise Solutions. It is undoubtedly the most comprehensive 'real' question bank, complete with detailed solutions by experts.

Studies have shown that successful JEE aspirants begin by familiarising themselves with the problems that have appeared in past JEEs as early as 2 years in advance.

Making it one of the key ingredients for their success. How about you then?
Get 45+21 Years Chapterwise-Topicwise Solutions to start your rehearsals early.



Available at all leading book shops throughout the country.
For more information or for help in placing your order:
Visit www.mtg.in to buy online!
Call 0124-6601200 or e-mail: info@mtg.in



Scan to explore & buy more
JEE Books on mtg.in



100 PERCENT

*Science, Mathematics,
 English & Social Science*

*Physics, Chemistry,
 Mathematics & Biology*

THE SMART STUDY SYSTEM

100% CBSE

MTG's 100 Percent Series is based on the most current CBSE guidelines and curriculum

100% SCIENTIFIC

MTG's editorial team knows what it takes to score more in exams - it keeps a close watch on changing examination patterns and has been able to put all ingredients critical to success together in a package aptly titled 100 Percent

100% CONTENT

Be it easy-to-comprehend text, graphics, illustrations or concept maps that complement, understanding theory and concepts is so straightforward with 100 Percent

100% CHECKS

After each topic, in every chapter, 100 Percent presents students with a variety of Q&A, including "Try Yourself", "NCERT Focus" & "CBSE Focus"

100% PRACTICE

100 Percent goes all out to ensure students are prepared for a diverse set of future challenges through a variety like Competition Focus, Exam Drills and Practice Papers, even Viva Voce Q&A for lab-based experiments

Series
 Available
 For CBSE
 Classes
9 to 12

**THE
 MISSION
 TO MARKS**



SCAN QR CODE



Available at all leading booksellers. To buy online, Scan QR code
 For more information, call 0124-6601200 (Lines open Mon to Sat, 9 am to 5:30 pm).
 Or e-mail at info@mtg.in now.

EXCEL ACADAMICS®

Bengaluru

A Premier Institute in KARNATAKA

Integrated PUC | PUC+NEET | PUC+IIT-JEE | PUC+CET



FOR
CBSE 12th
17.03.2023
FRIDAY

FOR
2nd PUC
24.03.2023
FRIDAY

NEET / IIT-JEE / CET LONG TERM COURSE

PCB / PCM (Repeaters Course)

REGISTRATIONS STARTED FOR

NEET, IIT-JEE, K-CET

CRASH COURSE & LONG TERM (REPEATERS) 2023-24

Note : Registrations Started for
Integrated PUC + CET / NEET / IIT-JEE

NEET / CET/ IIT-JEE

#326, 3rd A-Cross, Opp. People's Tree Hospital, Sheshadripuram College Road, Yelahanka New Town, Bengaluru - 560 064, KARNATAKA

7676-91-7777, 95136 57041, 9535656277.

★★★ Postal Coaching For PUC + CET / NEET / IIT-JEE / AIIMS / JIPMER Contact 9036357499 ★★★



Separate Deluxe Hostel for Boys and Girls





Outstanding Results

JEE (MAIN) 2023

Session-1



100

Percentile
Overall



Dhruv Sanjay Jain

4 Year
Classroom

43 100
Percentilers
(Physics or Chemistry or Maths)

625 **2606**

99+ Percentilers 95+ Percentilers
(Including Classroom + Distance & Digital) & Counting

➤ Scan the code
to check all the results
& watch the stories
of toppers



➤ Subject-wise 100 Percentilers from Classroom Programs

 100 Percentile in Maths 99.99 Percentile Kamyak Channa 4 Year Classroom	 100 Percentile in Chemistry 99.99 Percentile Aakash D Reja 4 Year Classroom	 100 Percentile in Physics 99.99 Percentile Mozeel Vanwani 1 Year Classroom	 100 Percentile in Maths 99.99 Percentile Tanishq Mandhane 4 Year Classroom	 100 Percentile in Physics 99.99 Percentile Shivanshu Kumar 4 Year Classroom
 100 Percentile in Physics 99.99 Percentile Adit Jindal 4 Year Classroom	 100 Percentile in Physics 99.98 Percentile Anoop Singh 4 Year Classroom	 100 Percentile in Physics 99.98 Percentile Shreyansh Solanki 2 Year Classroom	 100 Percentile in Physics 99.98 Percentile Shiv Narang 4 Year Classroom	 100 Percentile in Maths 99.98 Percentile Md Sahil Akhtar 4 Year Classroom
 100 Percentile in Physics 99.98 Percentile Raktim Kundu 2 Year Classroom	 100 Percentile in Physics 99.95 Percentile Siddharth Miglani 2 Year Classroom	 100 Percentile in Physics 99.95 Percentile Hardik Goyal 2 Year Classroom	 100 Percentile in Physics 99.95 Percentile Utkarsh Gupta 2 Year Classroom	 100 Percentile in Physics 99.94 Percentile Gaurang Gupta 3 Year Classroom

and many more top performers across 300+ branches in India

Though every care has been taken to publish the result correctly, yet Aakash Educational Services Ltd. shall not be responsible for inadvertent error, if any.

Admissions Open

➤ NEET | JEE

Foundation (Class 8-10)

Get up to **90% Scholarship***
Appear for Instant Admission &
Scholarship Test. Register now for FREE
visit: icst.aakash.ac.in or scan the code

Scan the code
to apply



➤ Crash Courses

for NEET / JEE 2023

Complete coverage of syllabus
Tests, Study Material & more
Hurry! Batches begin soon

Scan the code
to apply



*Terms & Conditions apply.




NEET Mock Test Paper
To Download,
Scan this QR Code



**JEE (Main & Advanced) & Other Engineering
Entrance Exams Mock Test Paper**
To Download,
Scan this QR Code

Download **Aakash App**



CALL (TOLL-FREE)
1800-102-2727

VISIT
aakash.ac.in

